THE PHILIPPINE

JOURNAL OF SCIENCE

B. MEDICAL SCIENCES

VOL. III DECEMBER, 1908. No. 6

STREPTOTHRICOSIS WITH SPECIAL REFERENCE TO THE ETIOLOGY AND CLASSIFICATION OF MYCETOMA.[1]

By W. E. MUSGRAVE and M. T. CLEGG, with a bibliography by MARY POLK.

(*From the Biological Laboratory and the Library, Bureau of Science, Manila, P. I.*

OUTLINE.

INTRODUCTION.

PART I.

ORIGINAL WORK.
- Material (cultures of streptothricæ).
  - General consideration.
  - Morphology.
  - Biology.
  - Pathogenesis.
    - Animal experiments.

REVIEW OF THE LITERATURE.
- General.
- As to genus determination.
- As to species and varieties.

GENERAL DISCUSSION.

CONCLUSIONS.

PART II.

STREPTOTHRICOSIS.
- Synonyms (actinomycosis—nocardiosis).
- Definition.
- History.
- Etiology.
  - Contributing causes.
  - Streptothricæ.
    - Synonyms.
    - Diagnosis.
    - Morphology.
    - Biology.
    - Pathogenicity.
    - Transmission; infection.
    - Species or varieties.
      - S. Actinomyces, types of: Bostrœm; Wolff and Israel; Wright.
      - S. Nocardii.
      - S. Eppingeri.
      - S. Maduræ.
      - S. Canis.
      - S. Capræ.

PATHOLOGY.
- General.
- Special.

SYMPTOMATOLOGY.
- General.

CLINICAL TYPES.
- Generalized streptothricosis.
- Thoracic streptothricosis.
  - Lungs; pleuræ; heart.
- Abdominal streptothricosis.
  - Liver; intestine; mesentery; kidneys; pelvic viscera; appendix, etc.
- Cerebral streptothricosis.
  - Brain; cord.
- External streptothricosis.
  - Streptothricosis of the jaw, mouth, etc. (Lumpy jaw.)
  - Streptothricosis pedis (mycetoma, Madura foot).
  - Miscellaneous streptothricosis.

DIAGNOSIS OF STREPTOTHRICOSIS.

PROGNOSIS.

COMPLICATIONS.

PROPHYLAXIS.

TREATMENT.

BIBLIOGRAPHY.

ILLUSTRATIONS.

[1] To be read in abstract at the Bombay Medical Congress, February 23, 1909.

INTRODUCTION.

Before entering on a study of the etiology of mycetoma, it is necessary to define the disease. This is exceedingly difficult, either from a clinical or an etiologic standpoint.

Formerly the diagnosis was based upon a clinical picture the essentials of which were a chronically enlarged foot with sinusses from which were discharged small granules of various colors and consistency, accompanied by a peculiar, oily degeneration of the tissues. Further classification was made principally upon the color of the granules; black, ochroid, white, red and mixed varieties being recognized. However, when bacteriologic studies began to show the multiplicity of the organisms concerned in the production of Madura foot it became necessary either to consider it a clinical entity of multiple etiology or to attempt an etiologic classification. This was made still more imperative by the discovery that the organisms causing this symptom complex were found also producing lesions in other parts of the body. A number of authors have continued to use a clinical classification, while others have attempted to give one based upon etiologic findings. The result is great confusion in the nomenclature of the disease.

As a prerequisite to an etiologic classification and definition of mycetoma it is necessary first to know and to be able to classify the etiologic factors concerned. So far this has not been done satisfactorily. The botanists are hopelessly confused in their grouping of the organisms and in consequence medical men have been unable properly to identify them. In the first place, authorities differ as to where the group of organisms belongs in the vegetable kingdom, some placing it among the bacteria and others higher up among the fungi. Practically all the latest writers upon the subject agree as to the great similarity between the various species of these organisms and the majority have placed them as species of a single genus, or as parts of two or even three closely allied genera. In order to establish a uniform and clear conception, and after carefully studying all phases of the question we have decided tentatively to accept *Streptothrix* Cohn 1875 as the generic name of the group of organisms under discussion.

The result of this decision is that the other names given to the tentative genus, such as *Actinomyces, Nocardia, Oöspora,* etc., become synonyms, the disease streptothricosis being limited to infection with parasites properly belonging to the genus, the various species of which will be considered presently.

Adopting such an interpretation, we have a fairly definite group of organisms with reasonably uniform pathologic and clinical manifestations, connected with other groups of closely related organisms giving somewhat similar manifestations. For example, the *Oïdia* or *Blastomyces* on the one hand and the tubercle bacilli on the other, may some-

what closely resemble the *Streptothricæ* in their action in certain parts of the body.

Mycetoma may be defined and classified in one of three ways, if *Streptothrix* is accepted as the most available generic name.

(1) It may be considered as a clinical disease of multiple etiology (which may or may not be limited to *Streptothrix* infections; (2) the term may appropriately be made synonymous with streptothricosis, actinomycosis, nocardiosis, etc.; (3) it may be made a clinical type (foot infection) by a *Streptothrix* (*Actinomyces* or *Nocardia*). Each of these classifications has something in its favor and also may be criticised.

The first would make diagnosis easy, for all forms of foot enlargement giving certain clinical manifestations regardless of the etiology, known or unknown, might be included, but it is open to the objection of not being definite and does not conform to our present methods of etiologic classification.

The second method has much in its favor and is practically adopted by Manson. The objections to it are that it adds a new significance to a term which does not clearly express the conditions and also adds a further synonym to a group of infections already rendered confusing by the number of its synonyms. The difficulty can more satisfactorily be met by the third method.

We have therefore decided to follow the third classification making mycetoma a *clinical* type or variety of streptothricosis—*Streptothricosis pedis*—and to define it as follows:

A disease consisting of a *Streptothrix* infection of the foot (*Streptothricosis pedis, Actinomycosis pedis*), characterized by a chronic course, swelling and deformity of the part, a peculiar, oily degeneration of the tissues with cavity and sinus formations and the discharge through the fistulous openings of mycotic aggregations containing the microörganisms.

Mycetoma, with this definition, becomes so intimate a part of streptothricosis that a balanced conception of it can only be obtained by a study of the whole subject of streptothricosis and a classification of the organisms concerned.

For this reason, the scope of this paper has been enlarged to include a discussion in two parts of the whole subject of *Streptothrix* or *Nocarida* infections.

Part I.

ORIGINAL WORK.

MATERIAL.

The material for this paper consists of the following cultures:

1. *S. freeri* Musgrave & Clegg, *This Journal* Sec. B. (1907), 2, 477.

2. *S. maduræ* Vincent. (This culture was sent to us by Professor Foulerton who writes as follows concerning it: "This culture has been in my hands for the last ten years; it corresponds with Vincent's description of his organism, and

is apparently of the same stock as that from which all the cultures in the London laboratories are derived.")

3. *S. maduræ* Vincent. (This culture was sent to us by Professor Binot of the Pasteur Institute in Paris.)

4. *S. maduræ* Vincent. (This organism was isolated by Dr. R. P. Strong from a case of the pink variety of mycetoma in India, and the culture has been given by him to us, for study.

5. *Streptothrix* of human actinomyces. (From Binot of the Pasteur Institute in Paris.)

6. *Streptothrix* of *"farcin de boeuf"* "A". (From the late Professor Nocard through Binot.)

7. *S. nocardii.* (From Foulerton who writes that "this culture was given to me by Nocard some six years ago.")

8. *S. eppingeri* "A".

9. *S. eppingeri* "B". (Both of these cultures are from Foulerton who writes as follows: " 'A' is a descendant of what I believe to have been Eppinger's original culture, 'B' a culture isolated by myself from a case which has not yet been published in detail. I have examined also a third culture of this species obtained by Dr. McDonald from a third case (*The Scottish Medical and Surgical Journal*, 1904). All three strains evidently belong to the same species."

10. *S. caprœ* Silberschmidt. (Received from Professor Silberschmidt of Zürich through Binot.)

11. *S. canis* Levy. (From Professor Levy of Strassburg through Binot.)

12. *S. chalcea* Foulerton. (Received from Foulerton who writes that this is a saprophytic species isolated by him from the air.)

The following cultures have also been received, but failed to develop on transplantation:

S. maduræ Vincent. (Furnished by Professor Vincent of the Val du Grace, Paris.)

S. maduræ. (From Legrain de Bougie through Binot.)

Streptothrix of bovine Actinomyces. (From Binot.)

Streptothrix of *"farcin de boeuf"* "N". (From the late Professor Nocard through Binot.)

Streptothrix of Eppinger ("A") and (Kl). (From Binot.)

Streptothrix of Deci. (From Binot who writes regarding this culture that "it was isolated by Deci in the Argentine Republic from a human case resembling tuberculous meningitis." Binot further states in his letter that all the cultures which he has forwarded to us were received by him directly from their authors).

Streptothrix "32". (Received from Foulerton who writes that this organism was isolated by him from a kidney removed from the living patient by operation; the organism was demonstrable in sections of the kidney and was present in large quantity. It is briefly referred to as *S. hominis III* in the *Lancet* (1906), 1, 970. "This organism resembles closely some of the apparently purely saprophytic species, and the infection in the case was probably primary in the lungs."

Streptothrix isolated from a case of mycetoma by Dr. Chaterjee of India.

Dr. Homer Wright states, in reply to a letter regarding cultures of *Actinomyces* and his *"Hyphomycete,"* that he no longer has cultures of these organisms.

A comparative study under conditions as nearly alike as possible has been made of all these strains, with due consideration of their morphology, biology, and pathogenic character. All of them, to obviate as

much as possible any influence of long cultural environment were, where feasible, first passed through animals and the studies then made from the reclaimed cultures taken from the experimental lesions. All the cultural and other biologic properties have so far as we have been able to do so, been studied under the same conditions in regard to media, temperature, and generation from animal lesions. Our results and consequent descriptions show some variation from previous descriptions of some of these organisms, but in general the picture is very much as given by the original authors.

We have failed entirely to obtain a culture of the Wolf and Israel and Wright type of *Actinomyces,* but, because of the great importance of this organism in the discussion, we have, for comparative purposes, taken James Homer Wright's description of it from his classical monograph. We have also failed to secure a definite culture of Bostroem's *Actinomyces,* although the one sent by Professor Binot from the Pasteur Institute is probably of this species. However, in view of a possible mistake we have used both the description given by Bostroem as well as the Pasteur Institute culture in our comparative work. The important characteristics of our cultures are shown in the following descriptions and are summarized in the attached table.

No. 1. *Streptothrix freeri.* This organism was isolated by us from a case of the ochroid variety of mycetoma and has already been fully described (*This Journal, Sec. B* (1907), **2**, 477). It is identical with Eppinger's organism which will be discussed subsequently in this paper, and the name given by us in the first publication should fall as a synonym for *S. eppingeri,* the latter having priority. Other described organisms of the group which are probably identical with this are those of Stokes, Aoyama and Miyamoto, MacCallum, Sabrezes and Rivière, and possibly several others.

No. 2. *Streptothrix maduræ.* This organism has been carefully described by Vincent, Foulerton, and others. A summary of its characteristics are given both in the text and table and in our first report and are again shown in the table attached to this paper. Our work with cultures of this species has been in the main confirmatory of the published results of others. We have worked with three cultures received respectively from Foulerton, Binot, and Strong. These strains are identical in every particular except that the one from Binot is less pathogenic for monkeys than either of the other two.

Morphologically this species gives very constant results. The colonies are made up of intertwining, long filaments which may, or may not, be radially placed at the periphery. There is true branching, but no club formation in cultures and none in experimental lesions in monkeys up to fourteen days. The organism takes the usual aniline stains; it is Gram positive, but is neither acid nor alcohol fast.

Culturally it is a facultative aërobe with strong oxygen requirements. It grows well in the incubator and at tropical room temperature. Positive transplants on artificial media are readily made, but some difficulty is often encountered in securing growth from experimental lesions. This difficulty is greatest when large quantities of the material are transferred to the media, probably because of an inhibiting material in the contents of the lesions. This hypothesis is supported by the fact that development is much more easily secured if the

granules from the lesions are washed in sterile salt solution before transferring them to artificial media. Growth on satisfactory media becomes apparent in from two to four days; development gradually continues for several weeks, depending somewhat upon the quantity of medium present. The principal characteristics of growth on various types of media are shown in the table.

Pathogenicity: Several monkeys have been successfully infected with the Foulerton and Strong strains of this organism, but results with the one from the Pasteur Institute were all negative. Intra-abdominal inoculation has proved the most satisfactory and under similar conditions the character of the lesions produced varies but little. The lesions consist of nodular, tubercle-like processes in the omentum, abdominal wall, and contiguous tissues. These lesions break down and become surrounded by pockets and channels containing a peculiar, pus-like material in which may be found the microörganisms. (See also animal experiments.)

No. 3. *Streptothrix maduræ.* (Pasteur Institute culture.) This is identical in all essentials with *S. Vincent* except that it is not pathogenic for monkeys. This may be due in part at least to its long cultivation on artificial media.

No. 4. *Streptothrix maduræ.* (Strong's culture.) This organism is identical with Vincent's species and its description has been included in the above.

No. 5. *Streptothrix actinomyces.* This culture was received from Professor Binot of the Pasteur Institute as human *Actinomyces.* It is quite different from the Wright type and it also appears to have slight distinctions from the Bostroem type of *Actinomyces.*

Morphologically it appears in the tissues as a fungus of the ray type, but without clubs in young experimental lesions. In colonies the cultures are dark, almost black in color, are made up of the branching filaments and transformation products, generally with radial arrangement of the terminals at the margins of the colonies.

The organism stains poorly by the usual laboratory methods; it is Gram positive and neither acid nor alcohol fast.

Culturally it is a facultative aërobe and grows on the majority of the usual media. In reclaiming cultures from experimental lesions this organism acts very much like *S. madurae* mentioned above; however, if the colonies are first washed in distilled water and then transplanted to artificial media, growth takes place without difficulty.

This type, like the majority of the species examined, grows better in media containing sugar or glycerine. Pigment is produced in all media and the body of the medium is generally darkened to a considerable extent by the culture. The pigment varies from an ochre-yellow in ordinary agar to a dark, almost black color in glycerine-agar and on potato.

The characteristics on varous media are shown in the table.

Pathogenicity: This species is pathogenic for monkeys by intraperitoneal inoculation. It produces the usual actinomycotic lesion like those seen in the clinical disease in man and animals. The organisms are found in the broken-down mass discharged from the sinusses.

No. 6. *Streptothrix* of *farcin du boeuf* "A," and

No. 7. *Streptothrix nocardii.* These two strains are identical in all essential particulars and may therefore be described together under the correct name of *S. nocardii.*

Nocard's original description of this organism was not very complete, but several other observers have studied his and other strains and a number of good descriptions have been given. Foulerton has devoted careful study to this species and our work is in general confirmatory of his description. We have

already noted the principal characteristics of this organism in both the text and table in our first report and a summary is also shown in the table in this one. Photomicrographic illustrations of its appearance in cultures are also submitted.

Morphologically this parasite shows marked variation under different environment. In young cultures and early lesions the hyphal forms are prevalent and quite uniform in appearance, but in older cultures and in colonies from lesions, irregularities are encountered. Coccus and bacillus-like forms predominate and very old cultures may almost have the appearance of mixed culture of bacteria. The filaments may or may not be radially placed at the periphery of the colonies. There is true branching, but no true clubs have been seen in cultures or in experimental lesions up to fifteen days of age. The Gram-Weigert method of staining gives good results and portions acid fast to the Ziehl-Neelsen-Gabbett method of staining are very numerous. The organism is also alcohol fast.

Culturally this species is a facultative aërobe. It grows on the majority of laboratory media at incubator or tropical room temperature; development becomes apparent in from two to four days after inoculation. The difficulty of securing transplants from experimental lesions, mentioned in connection with *S. maduræ* does not obtain with this organism. The principal cultural characteristics on various media are shown in the table and may briefly be summarized as follows:

Small, white, irregular, round, raised, opaque colonies develop on *ordinary agar* after two to three days. Growth is slow and not abundant and no pigment is produced.

Glycerine-agar after three days shows discrete colonies which coalesce later and become heaped up, presenting a moist, meal-like growth. No pigment is produced.

On *potato*, growth appears after 48 hours as a buff-yellow, granular, raised mass without pigmentation or erosion of this medium.

In *bouillon* growth appears after 48 hours as a grayish, flocculent mass at the bottom of the tube, later a few grayish granules appear on the surface of the medium and in some instances "puffball" formations occur at the bottom of the tube. The fluid remains clear.

In *ascitic fluid* a very slight growth slowly develops at the bottom of the tube.

In litmus-milk growth appears only at the bottom of the tube. The milk is not coagulated and its color is not changed.

Pathogenicity: This species is an exquisite tissue parasite for monkeys, and is pathogenic to a less extent for some other animals. Monkeys inoculated in the abdominal cavity and killed after five to fifteen days show characteristic lesions. In the majority of instances the early lesions are tubercle-like formations scattered through the mesentery and other tissues contiguous to the point of inoculation. In some cases pus pockets and channels are formed, surrounded by adhesions and containing granules made up of the microörganisms in the pus-like material. Guinea pigs inoculated in a manner similar to the monkeys give parallel results.

No. 8. *Streptothrix eppingeri* "A," and

No. 9. *S. eppingeri* "B." These strains are identical in all essential particulars and may be described under the correct name of *S. eppingeri.* This is one of the most important species of this group of organisms and has been well described several times. It has been isolated from anatomical lesions of such varying character as those from meningitis, by Eppinger and others, to Madura foot, as has been published in our first report. Foulerton has described this species and we have fully discussed its characteristics in our first report and summarized them in the table in this paper. Our work has demonstrated fully

the identity of *S. freeri* with this species and it is probable that several of the varieties described in the literature also must be classified with it. Its characteristics may be summarized as follows:

Morphologically a variety of forms exist varying from small, oval, coccus and bacillus-like bodies, to long branching filaments, depending upon the lesion or the age of the culture and the media used. In tissues the branching filaments are not seen by the ordinary method of staining, the growth occurring as a skein of thread-like filaments and with colony formation. The threads present a beaded appearance by using Gram-Weigert's stain. In cultures portions of the filaments are acid-fast by the Ziehl-Neelsen-Gabbett method, and they are also Gram positive. The entire mass from experimental lesions is often acid-fast.

Culturally the species is a facultative aërobe and grows on the majority of media. Growth becomes apparent on ordinary agar after two days, consisting of a smooth, adherent membrane which at first is porcelain-white, later it develops a delicate, orange-pink color. The medium remains moist and is not colored in its depth.

Glycerine-agar, after about three days, shows a growth consisting of small, whitish colonies, which gradually develop a delicate pink color. Later, as the colonies progress, they become umbilicated and coalesce, forming a heaped-up growth and producing a bright, orange color. The medium becomes slightly darkened.

On *potato* the growth appears after about four days as a granular layer, at first white, later becoming yellowish and gradually assuming a brick-red color. As the colonies develop they become umbilicated then coalesce and finally produce a moist, meal-like growth on the surface of the medium. The colonies can be lifted in heaps from the medium by the platinum loop, and when immersed in salt solution or distilled water readily disintegrate into fine, flat particles which float on the surface.

In *bouillon* growth appears after 48 hours in the form of small, flat particles, which at first are white, but later assume an orange tint on the surface of the medium. As growth progresses, the colonies coalesce and produce a membrane on the surface of the liquid. Portions of this membrane fall to the bottom of the tube.

In *ascitic fluid* a very slight growth occurs after six day's incubation.

Litmus-milk shows growth on the surface of the medium in the form of fine, white particles which in time becomes pink in color. A film is gradually formed on the surface of the milk, which is neither coagulated nor changed in color.

Pathogenicity: The three cultures of this species, including the two strains from Foulerton as well as our own, are pathogenic for monkeys, producing on inoculation characteristic progressive lesions. (See animal experiments.) Typical Madura foot may be produced by inoculation into the feet of monkeys. Subcutaneous inoculation produces abscess-like pockets, containing a thick, viscid pus in which the colonies of the organism may be observed as small, white granules. Intra-abdominal inoculation results in tubercle-like formations with adhesions, breaking down of tissue with abscess and sinus formation. Granules made up of the colonies are present in the abscess contents.

No. 10. *S. capræ.* A study of this species from the culture sent us by Foulerton confirms in general the findings of Silberschmidt and of Foulerton. It shows characteristics which differentiate it specifically from any of the others with which we have worked.

Morphologically this organism more closely resembles the bacteria than the others and in its action on monkeys it also resembles the tubercle bacillus quite closely. The filaments show true branching without club formation and they are shorter than with most of the other species. Bacillus- and coccus-like forms

predominate in cultures and are also very prevalent in experimental lesions. It stains by the stronger aniline dyes in a somewhat irregular manner, is Gram positive and largely and strongly acid and alcohol fast.

Culturally it is a facultative aërobe and grows on most of the laboratory media. Growth first becomes apparent in from two to four days and progresses slowly. The cultural characteristics are shown in the accompanying table and need not be repeated here.

Pathogenicity: This organism is pathogenic for monkeys and guinea pigs and by intraperitoneal injection for rabbits. The lesions differ somewhat from some of the other members of the group in that there is much less tendency to suppuration or other form of tissue necrosis. The lesion is essentially a tubercle-like formation which spreads from the point of inoculation by the formation of other small granules. When studied microscopically these lesions resemble the tubercle quite closely, always contain the microörganism and occasionally giant cells are encountered. The tubercle-like granules occasionally break down and form small pockets of thick granular pus-like material, but this is the exception and granules made up of colonies of the parasite have not been encountered as in the other species.

No. 11. *S. canis.* This organism is probably identical with *S. capræ* of Silberschmidt. The only important points of difference in the cultures which we have are shown in the table, and consist principally in the difference in reaction in litmus milk. Morphologically and in its pathogenic action *S. canis* gives very similar results to *S. capræ.* The same tubercle-like lesions are formed in experimental animals and the close similarity in other respects has led to the conclusion that the two belong to a single species.

No. 12. *S. chalcea.* This is a nonpathogenic species reported by Foulerton and carried along in this work principally as a control. It is distinguished specifically from the other species by morphologic and cultural characteristics, and particularly by its being nonpathogenic for monkeys.

ANIMAL EXPERIMENTS.

The following are summaries of our animal experiments. The general results are placed under the discussion of the characteristics of the various species of the microörganisms. A complete summary will be found in the chapter on general discussion and conclusions.

S. nocardi:

Monkey No. 3635, inoculated in the abdominal cavity with one loop of a two months' culture and the animal killed 12 days later. Autopsy shows numerous miliary tubercle-like granules extending over the entire surface of the omentum and mesentery with an occasional suppurative process. The pus when pressed out contains small, yellowish-white granules made up almost entirely of the microörganism.

Monkeys numbered 3722, 3726, 3714 were inoculated in a similar manner to the above. All showed similar lesions with the exception of monkey No. 3722 which gave a negative result.

Guinea pigs and rabbits inoculated in the abdonimal cavity for the greater part developed similar lesions. Intravenous inoculation in rabbits gave negative results.

S. eppingeri:

Monkey No. 3665 was inoculated in the abdominal cavity with one loop of a ten days' culture and the animal was killed thirteen days later. Autopsy shows numerous, tubercle-like granules in the lower portion of the mesentery,

with extensive adhesions of the mesentery to the abdominal wall. On separating the adhesions, or on section through them, numerous sinusses are exposed; these hold a thick, viscid, pus-like substance containing numerous small grayish-white granules composed principally of *Streptothrix*.

Monkeys numbered 3664, 3633, 3622 were inoculated in a similar manner to monkey No. 3665 and the lesions produced were in all essentials similar to those found in that animal.

Foulerton's strain of S. eppingeri.

Monkey No. 3725 was inoculated in the abdominal cavity with one loop of a ten days' culture and the lesions produced are similar to those found in animals inoculated with *S. eppingeri* of the original strain.

S. maduræ.

Monkey No. 3854 was inoculated by opening the abdominal cavity and burying a loop of the material from a two weeks' culture of Strong's strain of the organism. Ten days later the animal presented a distended abdomen with a marked tympanites and was killed on this date. Autopsy shows numerous miliary tubercle-like processes in the omentum and mesentery and adhesions surrounding the inoculated material. The adhesions are easily separated and show small, pinkish-white granules adhering to the mesentery.

Rabbits and guinea pigs inoculated subcutaneously, intravenously, and intraperitoneally with this organism gave negative results.

Monkey No. 3856 was inoculated in the abdominal cavity with one loop of a two weeks' culture of Fullerton's strain of this organism and killed two weeks later. Autopsy shows a small, tumor-like mass made up largely of adhesions confined to the coils of the intestine, omentum, and abdominal wall. On section the tumor shows several small abscesses from which can be expressed a thick pus containing small, white granules composed almost entirely of the organisms. The latter are present in large numbers.

Monkey No. 3724 was inoculated in a manner similar to the above with negative results.

Guinea pig No. 3856 was inoculated into the abdominal cavity with one loop of a two weeks' culture and killed after 10 days. Autopsy shows slight adhesions of the omentum to the abdominal wall and intestine, but no other lesions are present.

Subcutaneous and intravenous inoculations of monkeys, rabbits, and guinea pigs gave negative results in every case.

Monkeys numbered 3719, 3627, 3668 were inoculated with one loop of a two weeks' culture of Vincent's original strain of this organism. The inoculations were made subcutaneously and intraperitoneally and in no instance were progressive lesions produced.

Guinea pigs and rabbits inoculated subcutaneously, intravenously, and by intraperitoneal injections also gave negative results.

Actinomyces (from the Pasteur Institute):

Monkey No. 3864 was inoculated into the abdominal cavity with one loop of a two weeks' culture. The animal died three weeks later. Autopsy shows small abscesses in the abdominal wall at the site of inoculation and numerous miliary tubercle-like granules in the omentum and mesentery. The pus from the abscesses contains many minute, hard, black granules which by microscopic examination are seen to be made up of a net-work of mycelia and transformation products of the microörganism.

Monkeys numbered 3622 and 3623 were inoculated in a similar manner and presented similar lesions to the above. Intravenous, subcutaneous, and intraperitoneal inoculation of rabbits and guinea pigs gave negative results in each case.

S. capræ:

Monkey No. 3858 was inoculated into the abdominal cavity with one loop of a two weeks' culture and ten days later the animal was killed. Autopsy shows numerous miliary, tubercle-like bodies on the mesentery, diaphragm, and on the surface of the liver. There is slight adhesion of the omentum to the abdominal wall at the site of the inoculation.

Monkey No. 3625 was inoculated into the subcutaneous tissue and into the peritoneal cavity with a suspension of this organism. The animal died three weeks later. Autopsy shows extensive adhesions around the site of inoculation and there is a large abcess in the abdominal wall. On section the abcess contains a heavy, creamy, pus-like material in which are numerous, minute, grayish-white granules. These granules are composed of microörganisms and a pure culture is easily obtained by transplanting the granules to artificial media.

Guinea pig No. 3859 was inoculated in a similar manner as were monkeys numbered 3858 and 3625 and the lesions produced were similar to those described for the monkeys. Intravenous inoculations of rabbits gave negative results in the animals used.

S. canis:

Monkeys numbered 3633 and 3626 were inoculated in the abdominal cavity with one loop of a two weeks' culture and a two months' culture of this organism' respectively, and both animals were killed two weeks later. Autopsy shows slight adhesions near the site of inoculation. The omentum and mesentery are intensely congested and contain numerous miliary, tubercle-like granules.

Monkeys, guinea pigs, and rabbits inoculated subcutaneously showed no progressive lesions at autopsy. Intravenous inoculation of rabbits gave negative results. Intraperitoneal inoculation of guinea pigs in two instances produced lesions similar to those described for intraabdominal inoculations in monkeys.

S. chalcea.

Monkeys, guinea pigs, and rabbits when inoculated subcutaneously and intraperitoneally gave negative results.

LITERATURE.

A review of the literature of this subject may for convenience be divided into two divisions (1) *genus* and (2) *species* determinations.

Several of the generic names which have been introduced are clearly descriptive of organisms belonging to other genera such as *Cladothrix* or *Leptothrix,* and some refer to organisms properly belonging to the classes *Oïdia* or *Bacteria.*

GENUS.

The generic names which appear to have some claim for consideration are as follows:

STREPTOTHRIX.

Corda named this genus, giving, according to Saccardo (Sylloge Fungorum (1886), 4, 282) the following generic diagnosis:

"Hyphæ fertiles erectæ, virgato-ramosæ, ramis ramulisque spiraliter tortis. Conidia globosa v. ovoidea, in ramulis solitarie acrogena, subinde pleurogena, sessilia v. stipitellata, fusca."

Cohn (Untersuchungen über Bacterien, II.—Beiträge zur Biologie der Pflanzen (1875), 1, 186) used the name *Streptothrix* in a new sense in describing an

organism *Streptothrix fosteri*, which he found in the lachrymal ducts of a patient. The definition for this organism as given by De Toni and Trevisan (Saccardo's Sylloge Fungorum (1889), **8**, 928) is as follows:

"Filamentis tenuissimis, hyalinis, parallele insimul stratiformi-coalitis vel fasciculatis, rectis vel incurvis, sparse irregulariterque ramosis, in fragmenta inaequilia secedentibus."

CLADOTHRIX.

Cohn (*loc. cit.*, 185) under the name *Cladothrix dichotama* described an organism which was later included as the type species of a genus established by Marchand (Botanique cryptogamique, Paris (1883)) of which De Toni and Trevisan (Saccardo, (1889), **8**, 927) give the following generic diagnosis:

"Filamenta basi ab apice superiore distincta, vagina crassa obducta cylindrica, ætate provecta a basi ad apicem magis magisque incrassata, articulata, pseudo-ramosa. Arthrosporae binae in singulis microbaculis ellipsoideis ortae."

However Mace (*Compt. rend. acad. sci.* (1888), **106**) in a careful study of *Cladothrix dichotoma* Cohn showed that it did not have the supposed property of dichotomous division of the filaments, but that this appearance was simulated by a lateral branching process.

ACTINOMYCES.

Meyen (Linnæa (1827) **2**, 433) described a fungus under the name *Actinomyces horkelii*. Bollinger (*Centrabl. f. d. med. Wiss.* (1877), **15**, 481) published an account of a disease in cattle caused by a vegetable organism which was carefully described and named *Actinomyces* by Harz (*Jahresb. d. k. Central-Thierarzneischule*, München (1877–78), who at this time probably did not know of the previous occupation of the name by Meyen.

The work of Bollinger and Harz attracted much attention and was rapidly followed by a number of important contributions to the subject. For example, James Israel (*Virch. Arch.* (1878), **74**, 15; 50) described a similar organism as the cause of a similar disease in man and Ponfick (*Breslauer ärtzliche Zeitschrift*, May, 1879, and in subsequent publications) established the generic identity of the microörganisms in man and animals. The work of several writers showed that there existed varieties of this organism, but for a considerable time *Actinomyces* remained unquestioned as a generic name for the organisms.

Botanists classified the organisms in various ways, the most noteworthy being that of De Toni and Trevisan who created a new genus *Nocardia* (Saccardo, **8**, 927) and placed *Actinomyces* (Saccardo, **8**, 928) as a species with the following diagnosis:

"Glomerulis subglobulosis e filamentis densissime intertextis, fasciculos innumeros apice valde incrassato clavatos, a puncto centrali communi undique irradiantes constitutis."

OÖSPORA.

This generic name was introduced by Sauvageau and Radais (*Ann. Inst. Pasteur* (1892), **6**) to include the organisms previously described as *Actinomyces*, *Cladothrix*, and *Streptothrix*. This term never was seriously considered in the literature and needs no further notice.

NOCARDIA.

De Toni and Trevisan (Saccardo, **8**, 927) introduced *Nocardia* as the generic name and included as synonyms, *Streptothrix Actinomyces*, and in part *Discomyces*. The generic diagnosis given by them is as follows:

"Filamenta tenuissima, evaginata, articulata, Cladotricis more pseudoramosa, nunc e nucleo firmo radialiter expansa, nunc varie coalita. Arthrosporae in filamentis normalibus obvenientes, transformatione cocci singuli ortæ.—Est *Cladothrix* sine vaginis.

SPHÆROTILUS.

Engler (Syllabus der Pflanzenfamilien (1907), 5 ed., 5) places all the above-described organisms in the family *Chlamydobacteriaceae* of the *Schizomycetes*. He reduces the above-mentioned genera including *Actinomyces* to *Sphaerotilus* Kuetz (1833) the generic definition of which as given by De Toni and Trevisan (Saccardo, **8**, 926) is as follows:

"Filamenta premitus affixa, basi ab apice superiori distincta, initio simplicia, dein Cladotricis more pseudoramosa a basi ad apicem subæquilata, articulata, vagina gelatinosa obducta, in fasciculos crassos floccosos varie divisos consociata. Multiplicatio fragmentis filamentorum secedentibus, quae filamenta et fasciculos novos efficiunt. Arthrosporae numerosissimae, articulorum divisiones in tres directiones ortæ."

The above review shows the confusion which exists regarding the botany of the group of organisms which are variously recognized as *Streptothrix, Cladothrix, Actinomyces, Nocardia,* etc., and this seems to have been still further emphasized by workers in medicine and by other recent writers upon the subject. Some of the more important contributions are as follows:

Petruschky (Handbuch der Microörganismen, Kolle and Wassermann (1903), **2**, 832) classifies *Actinomyces*, *Streptothrix, Cladothrix* and *Leptothrix* as species of *Trichomyceta* of the *hyphomycetes*. He recognizes *Streptothrix* as the generic name of the parasites of mycetoma.

From the standpoint of an investigator in medicine, whatever may be the criticisms of botanists, the work of L. Gedoelst (Les Champignons parasites de l'homme et des animaux domestique, Paris, 1902) is particularly satisfactory, because of its simplicity of arrangement and attention to details. His classification places the organism under discussion among the *Fungi imperfecti* and in the genus *Discomyces* Rivolta (1878), with *Streptothrix* Cohn (1875); *Actinomyces* Harz (1877); *Nocardia* Trevisan (1889); *Oöspora* Sauvageau et Radais (1892) and *Discomyces* Blanchard (1900) as synonyms.

Foulerton (Allbutt's System of Medicine (1906), **2**, pt. 1, 302) who has been interested in this subject and has made many most important contributions to it, maintains *Streptothrix* Cohn to be the generic name. Foulerton gives four reasons for using *Streptothrix* in preference to any of the other terms.

1. He does not consider that Corda's indefinite use of the term in 1833 should make *Streptothrix* Cohn (1875) untenable.
2. This being acceptable, *Streptothrix* becomes the correct botanical name by priority of description.
3. *Streptothrix* was adopted by the committee of the Pathological Society.
4. It is by far the most generally used name.

James Homer Wright (Osler's Modern Medicine (1907), 1, 327; 340) who has made valuable contributions to the study of *Actinomycosis* recognized *Nocardia* as the correct generic name, but raises Actinomyces from species, as given by Trevisan, to generic rank and thus converts what most authors have included in a single genus, into two genera.

Sir Patrick Manson (Tropical Diseases (1907), 4 ed., 759) in his recent discussion of mycetoma has adopted a liberal etiologic classification, both by his definition and his classification of the varieties of mycetoma. He classes actinomycosis as one variety of mycetoma, designating it as actinomycotic mycetoma, and the other names under discussion here might also be included in this category, if we adopt this definition. This eminent author in his definition of mycetoma terms it "a fungus disease of warm climates, affecting principally the foot, occasionally the hand, rarely the internal organs or other parts of the body."

Manson in this classification follows Brumpt (*Arch. Parasit.* (1906), **10**, 489) who in turn apparently adopts his classification of the parasites from Gedoelst (*loc. cit.*). The diverse forms of the disease recognized are:

I. Actinomycotic mycetoma caused by *Discomyces bovis* (Harz 1877). II. Vincent's white mycetoma produced by *Discomyces maduræ* (Vincent 1894). III. Nicolle's white mycetoma the result of *Aspergillus nidulans* (Eidam 1883). IV. Buffard's black mycetoma caused by *Aspergillus bouffardi* (Brumpt 1906). V. Classic black mycetoma from *Madurella mycetomi* (Laveran 1902). VI. Brumpt's white mycetoma from *Indiella mansoni* (Brumpt 1906). VII. Reynier's white mycetoma resulting from *Indiella reynieri* (Brumpt 1906). VIII. Bouffard's white mycetoma from *Indiella somaliensis* (Brumpt 1906).

SPECIES CONSIDERATION.

Following the generic name of the entire group of organisms under consideration, the next logical step of the problem is a discussion of the species properly belonging to the genus.

The classification of species is a more difficult problem than the generic determination, largely because of the imperfect description of many of the so-called species or varieties. In fact, but few have been described with sufficient clearness to make determination possible.

It is not possible to state who first recognized this group of organisms. It may have been Corda, 1842; Mayen, 1827; Von Langenbeck, 1845 (Acland in Allbutt's System (1906), **2**, pt. 1, 325); Sir T. Smith, 1855 (*ibid*); Lebert (1857) (Traité d'anatomie pathologique, **1**, 54); Ballingal, 1855; or any one of several other writers, some of whom undoubtedly saw these parasites in lesions of the human body.

However, Henry Vandyke Carter, working from 1859 to 1874, first established the pathogenic rôle of the streptothrices in human pathology. While Carter did not cultivate his organism on artificial media and his descriptions are insufficient for species determination, a study of his publications leaves no doubt but that he was working with a clinical type of *Streptothrix* infection and his observations as to the specific etiology were quite satisfactory.

Erye 1860, Collas 1861, Biddie 1862, Coquerel 1866 Moxon and Hogg (*Trans. Path. Soc.*, London (1870)), Bristowe (*Trans. Path. Soc.*, London (1871)), Hogg (*Med. Times & Gaz.*) (1871), all recognized fungus-like bodies in cases of foot infection, which were clinically mycetoma or Madura foot.

Berkeley (*Med. Press & Circ.*, (1876)) states that he had cultivated one of

these organisms from a case of Madura foot which had been forwarded to him by Vandyke Carter. He named the organism *Chionyphe carteri.*

Rivolta (*Med. Vet.* (1868)) probably saw these organisms in pus taken from an abscess on the jaw of a cow. Rivolta (1868–1875) also found disc-like bodies in nodular masses from the tongue of a cow. He named the disease sarcomycosis.

Robin (Traité de Microscopie, Paris (1871), 575) gives a brief description of yellowish granules from two abscesses which were probably due to *Streptothrix* infection.

Heller (about 1872, completed a work which he published in *Deutsch. Arch f. klin. Med* (1885), **37,** 372) and from his description and drawing probably saw these organisms in the discharges from a case which he later considered to be acute actinomycosis.

Perroncito (1875) evidently saw and described species of this organism taken from a pseudosarcoma of the jaw of a cow.

Cohn (*Beitr. Biol. d. Pflanzen.* (1875)) **1,** 148, described an organism which he obtained from concretions in the lachrymal canals of a patient, terming it *Cladothrix fosteri.*

Bollinger (*Deutsche Ztsch. f. Thiermed.*, Leipz. (1877), **3,** 334) noted the constant occurrence of a branching organism in lumpy-jaw of cattle; Harz (*ibid,* (1878)) described it, naming the organism *Actinomyces bovis* and Bollinger termed the disease actinomycosis.

Israel (*Virch. Arch* (1878), **74,** 15) described a disease in man similar to actinomycosis in cattle and detailed the microscopic appearances of the causative *Streptothrix.* Ponfick (*Breslau. ärztl. Ztschr.* (1879) furnished strong evidence of the etiologic identity of the diseases of man as described by Israel and of cattle as given by Bollinger and Harz.

Corre (*Arch. méd. Nav., Par.* (1883)) in reporting the notes of Dr. Collas called attention to the probable identity of actinomycosis with the earlier described mycetoma, and many other authors since that time have come to the conclusion that the diseases are probably the same.

Mandareau (*Bull. acad. méd.* (1887), **18,** 555) gave a morphologic description of a species of *Streptothrix.* Nocard (*Ann. Inst. Pasteur* (1888), **2,** 293) described and cultivated a *Streptothrix* from the lesions of a disease known as *farcin du boeuf* which was very prevalent among cattle in Guadeloupe. Organisms which agree with Nocard's description have repeatedly been isolated from pathologic lesions by other authors. Nocard's work was epoch making in that he was the first satisfactorily to cultivate a *Streptothrix* in pure culture.

Affanassiew and Schultz (*Münchener med. Wchnsch.* (1889), **36,** 418) briefly described a *Streptothrix* obtained from pus and sputum. Cultures were pathogenic for guinea pigs. Protopopoff and Hammer (*Ztschr. f. Heilk.* (1890, **11,** 255) reported incomplete studies of the cultures of Affanassiew's and Schultz's organism. Kischensky (*Arch. f. exp. Path. u. Pharm.* (1889), **26,** 79) partially described and probably cultivated a *Streptothrix* which may have been identical with that cultivated later by Wolff and Israel from a human case of actinomycosis of the lungs. Bujwid (*Centrbl. f. Bakt., etc. Abth.* 1 (1889), **6,** 630) cultivated and briefly described a *Streptothrix* which was a facultative anaërobe taken from a human case of actinomycosis.

Eppinger (*Wien klin. Wchnsch.* (1890), **3,** 321) cultivated a *Streptothrix* from a brain abscess and a "pseudo tuberculosis of the lungs", the patient having manifested the clinical symptoms of meningitis during life. This organism was Gram positive, acid-fast and a facultative aërobe. Growth occurred on the ordinary laboratory media and the organism was pathogenic for laboratory

animals. Eppinger's publication, like that of Nocard, forms a landmark in the history of the disease. His work has been confirmed by MacCallum and others and his organism, which is a different species from that of Nocard, will be more fully considered in this paper.

Almquist (*Ztschr. f. Hyg.* (1890), **8**, 189) cultivated species of *Streptothrix* from three different sources. The first was discovered as a contamination in a gelatin culture tube, the second was cultivated from pus taken from the base of the brain from a human case and the third culture was isolated from water. Liebman (*Arch. per le sci. med.* (1890), **14**, 361) discussed the cultivation of *Streptothrix* in pure culture.

Bostroem (*Beitr. z. path. Anat., etc.* (1891), **9**, 1 to 240) cultivated a single species of *Streptothrix* from eleven cases of actinomycosis. His organism was Gram positive and in part acid-fast and was a facultative aërobe in cultures. Growth was obtained with difficulty directly from the lesions, but the later transplants were more satisfactory in artificial media. Bostroem's article was the first published exhaustive consideration of the cultivation of *Streptothrix*. Wolff and Israel 1891 (*Virch Arch.* (1891), **126**, 11) cultivated a *Streptothrix* from two cases of clinical actinomycosis in man. In one case the lesion was a retromaxillary tumor and the other was actinomycosis of the lungs. This organism was acid-fast and a facultative anaërobe. Gruber (*Tr. VII Intern. Cong. Hyg. & Demog.*, Lond. (1891), **2**, 65) discovered (as an accidental contamination of culture media) an organism which he described as *Micromyces hofmanni*. This organism produced pyæmia in animals and "ray fungus" types were present in the lesions. The articles by Doyen (*Tr. VII Intern. Cong. Hyg. & Demog.*, Lond. (1891), **3**) and Roussel (These, Paris, 1891) are not available to us.

Hesse (*Deutsche Ztschr. f. Chir.* (1892), **34**, 274) obtained a *Streptothrix* by culture from an abscess in the groin, which when opened was found to communicate by a sinus with the intestine. Autopsy on the patient, who died after more than a year's illness, showed a rectal fistula communicating with multiple abscesses in and around the pelvis. There was bronchitis, but the organisms were not found in the sputum. The organism was Gram positive and aërobic in culture media. Growth occurred on the majority of media and serum media were liquefied. No positive results were obtained from intravenous, intraperitoneal, or subcutaneous injections of cultures into rabbits, guinea pigs or white mice. Sauvageau and Radais (*Ann. Inst. Pasteur* (1892), **6**, 242) studied a number of streptothrices which had already been isolated by other authors. They made a classification of *Streptothricae* in general, based upon cultural characteristics.

The articles by Lanz (*Cor. Bl. f. schweiz. Aerzte* (1892), 307; 339); Ebermann (Inaug. Diss. St. Petersburg (1893); Dor (*Gaz. hebd. de med.* (1893) 2 s., **30**, 40) and of Dor and Bérard (*Prov. méd.* (1893) are not available to us.

Vincent (*Ann. Inst. Pasteur* (1894), **8**, 129) cultivated a *Streptothrix* which was pathogenic for laboratory animals from a case of ochroid variety of mycetoma. We have already noticed this organism somewhat in detail and it will be considered further. Vincent's work is very important because he was the first satisfactorily to cultivate a *Streptothrix* from the Madura foot type of the disease and because his organism is specifically different from any of those described before his paper appeared. Boyce and Surveyor (see bibliography) in a series of articles, described streptothrices which they were able to isolate from cultures made by friends in India, taken from cases of mycetoma and forwarded to the authors in England. The brief description given by these authors renders it probable that this organism was identical with that of Vincent. Karg (Ref. Düms' *Deutsche mil-ärztl. Ztschr.* (1894), **23**, 145) cultivated a *Streptothrix*

from a case of actinomycosis in man. Karg considered his organism to be identical with that of Bostroem. This author also reported a second case of the disease with results similar to those obtained with the first one. Leith (*Edinb. Hosp. Rep.* (1894), **2**, 121) and Newjadomsky (Ref. Lubarsch and Ostertag Ergb. (1898), **5**, 665) each isolated by culture a *Streptothrix* from a human case of actinomycosis. The article of Kitchensky (*Arch. f. exper. Path. u. Pharmakol.* (1890), **26**, 79) is not available to us. Sabrazès and Rivière (*Presse méd.* (1894), Sept. 22) reported the cultivation of a *Streptothrix* from abscesses in the brain, lungs, and kidneys of a human case. This organism only grew anaërobically. In their second case (*Mercredi méd.* (1895), 485) an aërobic *Streptothrix* was cultivated from the sputum and subcutaneous abscesses of a patient. Both these organisms are reported as being pathogenic for small animals, the latter one, however, only producing lesions when injected together with dilute acetic acid.

Aschoff (*Ber. klin. Wchnsch.* (1895), **32**, 738) cultivated a *Streptothrix* from a case of human actinomycosis and stated that he had produced actinomycotic lesions in animals by inoculating cultures of the organism.

Du Bois Saint-Sevrin (*Semaine méd.* (1895)) cultivated a *Streptothrix* from a case of ulcer of the cornea in a human patient with a pseudomembranous conjunctivitis. Ferré and Faguet (*Mercredi méd.* (1895), 441) did the same from an abscess in the *centrum ovale* of the brain. The authors considered this organism to be identical with that of Eppinger. Rullmann (Ref. *Centralbl. f. Bakt., etc.* (1895), **17**, 884) cultivated a *Streptothrix* from soil and again in 1898 the same author (*Münchener med. Wchnsch.* (1898), **45**, 919) cultivated what he apparently considered to be the same organism from a case of disease of the lung. It was a *Streptothrix* positive to Gram's stain, which grew on ordinary media as a facultative aërobe. Positive local lesions were produced in rabbits, guinea pigs, and mice when these animals were inoculated with cultures. Garten (*Deutsche Ztschr. f. Chir.* (1895), **41**, 257) cultivated a *Streptothrix* from lesions in a human case. There was necrosis of the vertebrae and ribs, with abscesses, sinus formation and empyæma. This organism grew on laboratory media and was pathogenic for three out of thirty-seven rabbits and guinea pigs.

Dor (*Gaz. hebd. d. méd.* (1896), **43**, 553) obtained a *Streptothrix* from an actinomycosis-like abscess at the angle of the jaw. The pus contained granules with long, beaded filaments which did not stain by Gram's method. The organism grew aërobically on liquid media only. Sawtschenko (*Baumgarten's Jahresb.* (1896), **12**, 613) cultivated an organism from a case which he termed pseudoactinomycosis. It formed long threads which did not stain by Gram's method and which grew only under anaërobic conditions. Jurinka (*Mitth. a. d. Grenzgeb. d. Med. u. Chir.* (1896), **1**, 139) cultivated a *Streptothrix* from a human case of actinomycosis which he considered identical with that of Wolff and Israel. The article by Kozerski (Ref. *Jahresb. d. ges. Med.* (1896), **2**, 223) is not available to us.

Scheele and Petruschky (*Verhandl. d. Con. f. innere Med.* (1897), **15**, 550) cultivated a *Streptothrix* which was acid-fast by the Ziehl-Neelsen method from the sputum and subcutaneous abscesses of a case of pyæmia, pleuro-pneumonia, cystitis, and pyelitis. It grew on agar, gelatin, and broth. Its pathogenic character for animals apparently was not determined. Buchholtz (*Ztschr. f. Hyg.* (1897), **24**, 470) described a *Streptothrix* found in the lung of a patient suffering from a disease which, both clinically and pathologically, very closely resembled tuberculosis of the lungs. The organism was Gram positive and acid-fast, but did not grow on artificial media. Urban (*Münchener med.*

Wchnsch. (1897), **44**, 124) cultivated a *Streptothrix* from a case of human actinomycosis. Berestnew (Aktinomykose und ihre Erreger (Dissert), Moskau (1897) reported several cases. A *Streptothrix* was cultivated from a subperiostial abscess of the lower jaw of a woman (Case I). This clinically was a case of "lumpy jaw." This organism was a *Streptothrix*, facultative anaërobic, Gram positive, dying after three generations on artificial media. It was not pathogenic for rabbits, guinea pigs, or tame mice. The organism from his Case II was cultivated from an abscess under the right nipple of a 16-year-old patient who had symptoms of disease of both lungs. The organism differed from that of Case I in that it grew somewhat better in the presence of oxygen. The cultures in Case III were from sputum and chest sinuses in a patient forty-four years of age, who died after an illness of about one year. The organism was Gram positive, facultative aërobic, and grew best in bouillon. It was neither fatal for rabbits nor guinea pigs when injected either intraperitoneally or subcutaneously; a temporary infiltration of the tissues resulted by the latter method. Abel (*Münchener med. Wchnsch.* (1897), **44**, 124; 149) made brief reports of a pathogenic *Streptothrix* from sputum. Van Neissen (*Virch. Arch.* (1897), **150**, 482) worked with an organism which was not differentiated from that of Wolff and Israel.

Berestnew in another case (*Ztschr. f. Hyg. u. Infectionskrankh.* (1898), **29**, 94) obtained a *Streptothrix* in pure culture from the lesions in a child suffering from an abscess of the lung and pyæmia. This organism was negative to Gram, grew anaërobically on media and was pathogenic for animals. Petruschky (*Deutsche med. Wchnsch.* (1898), **24**, Ver.-Beil. 78) cultivated an organism similar to that of Scheele and Petruschky (*loc. cit.*) from the sputum obtained from a second case of streptothricosis. Alissow and Skworzow (ref. Lubarsch and Ostertag (1898), **5**, 664) studied two cases of streptothricosis in man. They obtained two organisms in bouillon cultures, a bacillus and a *Streptothrix* growing in symbiosis. Pure cultures of the *Streptothrix* developed only clubs and rods, but when bacteria were again mixed in the cultures, long fungus threads were produced. The authors consider the lesions to be due to the mixed infection. Schtscheglow (1898) (*ibid.*), also secured mixed cultures and the bacteria present influenced the morphology of the *Actinomyces*. Newjadomsky (1898) (*ibid*) reported one case in man in which a *Streptothrix* was obtained in pure culture from a subdiaphragmatic abscess. The author considered the organism to belong to a new species. It was not pathogenic for guinea pigs. Rullman (*Münchener med. Wchnsch.* (1898), **45**, 919) cultivated a *Steptothrix* from the sputum of a patient with a localized lesion near the root of the right lung. This organism was Gram positive and grew on all media. It was pathogenic for mice, rabbits and guinea pigs, and could be reclaimed by culture from lesions in these animals. Grillo (*Reforma Medica* (1898), **14**, 301) cultivated a *Streptothrix* from a case of human actinomycosis and from the bovine infection and reproduced the disease by animal inoculation with cultures of the organism. Harbitz (*Norsk Mag. f. Lægevidensk* (1898), **13**, 1) obtained a *Streptothrix* by culture from five cases of human actinomycosis. Flexner (*Journ. Exp. Med.* (1898), **3**, 435) reports a case of human infection with a *Streptothrix*, but his organism did not grow on artificial media. The article by Marwedel (*Beit. z. klin. Chir.* (1898), **21**, 561) is not available to us.

Foulerton (*Lancet* (1899), **2**, 779) cultivated his *Streptothrix hominis* from abscesses of the chest with deep sinuses in a patient showing symptoms of a severe lung involvent with pyæmic symptoms. This author (see bibliography) has also made numerous other contributions to the morphology of the subject and has probably done more than any other toward clearing up the confusion which

exists regarding this group of organisms. Levy (*Centralbl. f. Bakt., etc.* (1899), **26,** 1) cultivated a *Streptothrix* from five cases of clinical actinomycosis in man. He considers his organism to be identical with that of Wolf and Israel. Paul Krause (*Centralbl. f. Bakt., etc.* (1899), **26,** 209) obtained a *Streptothrix* from an abscess of the lower jaw. It was Gram positive but not acid-fast, a facultative aërobe and grew on ordinary laboratory media. No positive results were obtained by intraperitoneal or subcutaneous injection of cultures into rabbits, guinea pigs, or white mice. Hayo Bruns (*Centralbl. f. Bakt., etc.* (1899), **26,** 11) cultivated a *Streptothrix* from a lesion which according to the diagnosis was an actinomycotic abscess of the abdominal wall. The organism was Gram positive, but not acid-fast and did not stain well with the ordinary aniline dyes. Growth occurred on ordinary media, best under aërobic conditions. No lesions were produced by intravenous, intraperitoneal, nor subcutaneous injections of cultures in rabbits, guinea pigs, or mice. Silberschmidt (*Ann. Inst. Pasteur* (1899), **13,** 841) obtained a *Streptothrix* by culture from the lungs of a goat which had a disease closely resembling tuberculosis. This organism was carefully described by the author and later by Foulerton and Jones, (*Tr. Path. Soc.* (1902), **53,** 99) who considered it to be a new species and our work is in accord with this conclusion.

Brault (*Arch. d. parasit.* (1899), **2,** 535) observed two human cases of actinomycosis. He cultivated a *Streptothrix*, at first in symbiosis with bacteria and later in pure cultures. His organism grew on ordinary media and was pathogenic for animals.

Norris and Larkin (*Journ. Exp. Med.* (1900), **5,** 155) obtained a *Streptothrix* by culture from two cases of bronchopneumonia and bronchiectiasis. Both cases proved fatal. This organism was Gram positive, but not acid-fast by the Ziehl-Neelsen method. The authors consider it probably to be identical with Israel's *Streptothrix*. Local lesions were produced in both rabbits and guinea pigs by injection of the cultures. Cozzolino (*Ztschr. f. Hyg. u. Infectionskranfh.* (1900), **33,** 36) reports a human case of *Streptothrix* infection of the ear. He cultivated an organism from the lesion which was pathogenic for mice and guinea pigs and described it very well. It was a facultative aërobe, but branching was not observed.

Dean (*Tr. Path. Soc.* (1900), **51,** 25) cultivated a *Streptothrix* from the lesions in a horse which clinically was suffering from actinomycosis. He considered his organism to be a new species, differing in certain particulars from any described previously. This organism was a facultative aërobe, Gram positive, and grew very poorly on the majority of media and not at all on others. It was pathogenic for animals, producing lesions which contained organisms resembling *Actinomyces*, with club-shaped ends. However, in cultures no club formations were observed. Tusini (*Arch. f. klin. Chir.* (1900), **62,** 249) observed four cases of *Streptothrix* infection in man. The one reported in detail was an infection which clinically was Madura foot. From this case, he cultivated a *Streptothrix* which was pathogenic for guinea pigs and which in cultural characteristics closely resembled the organism of Vincent. Sternburg (*Wien. klin Wchnsch.* (1900), **13,** 548) cultivated a *Streptothrix* resembling that of Wolff and Israel from three cases of human actinomycosis.

Aoyoma and Miyamoto (*Mitth. a. d. med. Fac. d. k. jap. Univ. a. Tokio* (1901), **4,** 231 to 276) obtained a *Streptothrix* in pure culture from an abscess of the lung. This was Gram positive, acid-fast, and a facultative aërobe; pathogenic for rabbits, white mice, and hens. The authors believed their organism to be very closely related to or identical with *S. eppingeri.* Mertens (*Centralbl. f. Bakt., etc.* (1901), **29,** 649) reported one human case and cultivated a *Streptothrix*

from the lesions. Cultures at first only grew in bouillon, but gradually growth on other media was successful. The anaërobic tendencies of his organism were changed so that it grew aërobically after a considerable time on artificial media. This organism produced lesions in the eyes of rabbits. Silberschmidt (*Ztschr. f. Hyg. u. Infectionskrankh.* (1901), **37**, 345) studied eight cases of *Streptothrix* infections in man. He cultivated the organism and believed he had isolated three different species. In six of his eight cases the organisms closely resembled those of Wolff and Israel. The others more closely approximate Bostroem's organism. Jelenewski (Ref. *Baumgarten's Jahresb.* (1901), **17**, 495) studied fifty cases of actinomycosis occurring in the mouths of animals. He cultivated the fungus first with *B. subtilis.* Later he obtained pure cultures both aërobically and anaërobically. Cultures of the organisms were pathogenic for calves, but not for rabbits and guinea pigs. The author concluded that his *Streptothrix* was a new species, basing his decision upon both its cultural and pathogenic properties.

Foulerton and Jones (*Tr. Path. Soc.* (1902), **53**, 56) studied the group extensively and described some new species. The tentative classification by these authors was the most satisfactory which had been made up to the time of its publication.

They also cultivated a *Streptothrix* ((ibid), **75**, 93) which they believed to be the cause of the disease, from a conjunctivitis and keratitis in a twelve year old girl. The organism was Gram positive, but not acid-fast; a facultative aërobe growing on all media. It was not pathogenic for rabbits, guinea pigs, or white mice. The same authors (*ibid,* 100) isolated a *Streptothrix* from "a case of pulmonary infection in a woman." This organism was Gram positive, not acid-fast, grew on the usual laboratory media, and was not pathogenic for rabbits. MacCallum (*Centralbl. f. Bakt., etc. orig.* (1902), **31**, 529) cultivated a *Streptothrix* which he considered to be identical with that of Eppinger from the peritoneal exudate in the body of a child dead of subacute purulent peritonitis which followed a surgical operation for relief of stricture of the oesophagus. Birt and Leishman (*Journ. Hyg.* (1902), **2**, 120) studied a *Streptothrix* which was obtained from a fatal human case. Empyæma, pericarditis, and "cirrhotic nodules in the lungs" were present at autopsy. The organism was isolated from the pleuræ, pericardium, and lung. It was Gram positive and acid-fast and showed strong aërobic tendencies. It grew on the majority of media, producing a coral-pink color. This organism was pathogenic for guinea pigs and is probably identical with Eppinger's *Streptothrix.* Schukewitsch (Ref. *Jahresb. d. ges. Med.* (1902), **1**, 711) cultivated a *Streptothrix* somewhat resembling that of Wolff and Israel from twenty-two out of twenty-five cases of human actinomycosis. Von Baracz (*Arch. f. klin. Chir.* (1902), **68**, 1050) reports on fifty-two cases of actinomycosis. The article is a splendid monograph, but does not go extensively into the question of the etiology of the disease. Welleminsky (*Deutsche med. Wchnsch.* (1902), **28**, Ver.-Beil. 196) cultivated a *Streptothrix*-like organism, nonpathogenic for animals, from lesions in one case.

Trolldenier (*Ztschr. f. Thiermed.* (1903), n. s. **7**, 81) reports a case of actinomycosis in a dog. He cultivated a *Streptothrix* which he considered to be a new species from the lesions. The organism was pathogenic for mice and guinea pigs, rabbits, dogs, and calves. Doepke (*Münchener med. Wchnsch.* (1903), **50**, II, 2245) studied several human cases and cultivated organisms which where similar to that of Wolff and Israel. Lignières and Spitz (*Arch. d. parasitol.* (1903), **7**, 428; *Centralbl. f. Bakt., etc.* (1904), **35**, 294; 452) cultivated a *Streptothrix* from a calf suffering from actinomycosis which they considered similar to that of Wolff and Israel. The article by Horst (*Ztschr. f. Heilk., Abt. f. path. anat.* (1903), **24**, 157) is not available to us.

James Homer Wright (*Journ. Med. Research* (1904), **13**, 349) cultivated what he believes to be a single species of *Actinomyces* from 13 cases of human actinomycosis and two cases of the bovine disease. He considers his organism to be of the same species, if not identical, with that of Wolff and Israel. This article is the most thorough study of one species of this group of organisms that has yet been made. Stokes (*Am. Journ. Med. Sci.* (1904), n. s. **128**, 861) obtained a *Streptothrix* from an abscess of the lung. This organism was acid-fast, grew aërobically and was pathogenic for laboratory animals. Schabad (*Ztschr. f. Hyg.* (1904), **47**, 41) considers the true *Actinomyces* and the organisms of Bostroem or Eppinger to be different species. He cultivated from the sputum and abscess of the breast in one human case an organism which he considered to be identical with that of Eppinger in its morphology, biology, and pathogenicity. Hartman (*Tr. Chicago Path. Soc.* (1904), **6**, 157) cultivated a *Streptothrix* from one human case. The articles by Tuttle (*Med. and Surg. Rep. Presbyterian Hosp.* N. Y. (1904), **6**, 147) and McDonald (*Scot. Med. & Surg. Journ.* (1904), **14**, 305) are not available to us.

Pinoy (*Comp. rend. acad. sci.* (1906), **143**, 1175) isolated a *Streptothrix* and claimed to have experimentally reproduced the black granules by inoculation of cultures into a pigeon's foot. His organism was a facultative anaërobe. Nicolle and Pinoy (*Arch. d. parasitol.* (1906), **10**, 437) cultivated a *Streptothrix* from a case of mycetoma in an Arab.

Musgrave and Clegg (*This Jurnal, Sec. B.* (1907), **2**, 477) obtained a culture from the sinuses in a case of the ochroid variety of mycetoma occurring in a Filipino woman in Manila. The cultural characteristics have already been described and will be further discussed. Poppenheimer and Satchwell (*Journ. Infect. Dis.* (1907), **4**, 617) report a case of endocarditis and isolation of a pleomorphic bacterium resembling somewhat closely the *Streptothricæ*. They review the literature of *Streptothrix* infections in man and tentatively classify their organism as a *Cladothrix*. The report of these authors is thorough and complete and from their description it does not seem as if their organism could properly be considered under *Streptothricæ*.

Strong cultivated a *Streptothrix* from a case of mycetoma in India. This organism is undoubtedly identical with that of Vincent.

OTHER STREPTOTHRICÆ.

In addition to the pathogenic organisms of the group under discussion and which have been isolated from pathologic lesions in man and animals, others have been cultivated from sputum, pus, or other human excreta under circumstances which make their exact source questionable; there are still others which have only been described as being obtained from animals; and some have been cultivated only from air, water, soil, and other outside sources. The following may be mentioned: *S. rubra* of Ruiz-Casabó (ref. *Centralbl. f. Bakt., etc.* (1895), **17**, 466) isolated from sputum; *Cladothrix mordoré* of Thiry (*Arch. d. physiol.* (1897), **29**, 284) obtained from an anginal exudate; the *Streptothrix* of Dean (*Tr. Path. Soc.* (1900), **51**, 26) isolated from an abscess of the jaw in a horse; and *S. lacertæ* of Terni (ref. *Centralbl. f. Bakt. etc.* (1896), **19**, 953) isolated from nodules in the liver of a lizard. Rossi-Doria (*Ann. d. l'Ist. d'ig. sper. d. Univ. di Roma* (1891), **1**) cultivated a *Streptothrix* from a case of bovine actinomycosis. Sanfelice (*Centralbl. f. Bakt. etc., orig.* (1904), **36**, 355; and *Arch. f. wiss. u. prakt. Thierheil.* (1896), **22**, 153) obtained *Streptothricæ* from several minor cases of bovine actinomycosis. He considered his organisms to be similar to those of Gasperini and Doria. Gasperini (*Proc.-verb. Soc. Tosc. di Sci. Nat.* (1895), **9**, 64; 292) isolated and described two strains of *Streptothrix* from actinomycosis in cattle. He (*Ann. d. l'Ist. d'ig. sper. d. Univ. di Roma* (1892), **2**, 167; *Proc.-verb. Soc.*

Tosc. de. Sci. Nat., July 5, (1896) also worked with a number of cultures obtained from other authors including that of Dessy (*loc. cit.*). With this latter organism, he stated that he had produced typical actinomycosis by injection into the jaw of a cow.

Lange and Manasse (ref. Levy. *Centralbl. f. Bakt. etc.* (1899), **25,** 5) isolated a *Streptothrix* somewhat resembling that of Wolff and Israel from actinomycosis in a dog. Bahr (*Ztschr. f. Thiermed.* (1904), n. s. **8,** 47) cultivated a *Streptothrix* from an actinomycosis-like disease in a dog. Dean (*Tr. Jenner Inst. Prevent. Med.* (1899), **2,** 17) did the same with an organism resembling that of Wolff and Israel, obtained from an abscess in the jaw of a horse and this *Streptothrix* was pathogenic for animals. Mosselmann and Liénaux (*Ann. d. méd. vet.* (1890), **39,** 409) obtained a *Streptothrix* from a case of bovine actinomycosis.

Lignières and Spitz (*Bull. Soc. centr. d. méd. vet.*, Paris (1902), **56,** 487) cultivated a bacillus from cases of clinical actinomycosis in cattle. They state they produced true actinomycosis, the organism showing the club-shaped *Streptothrix* granules, with cultures of this organism, by inoculation of their cultures into cattle.

The *Streptothrix* of Caminiti (*Centralbl. f. Bakt. etc., orig.* (1907), **44,** 193) is one of the most important of the organisms cultivated from the air, because of its pathogenic properties and close cultural resemblances to the *Streptothricæ* which produce clinical actinomycosis. This organism is Gram positive, acid-fast in part, a facultative aërobe with anaërobic preferences, liquefies gelatin and produces pigment in varying amount and color in artificial media. It is pathogenic for all animals experimented upon, producing circumscribed local lesions by injection of cultures.

Other organisms of this class which may be mentioned are *S. leucea* Foulerton (*Tr. Path. Soc.* (1902), **53,** 101) isolated from sewage and water; *S. alpha* Jones (*ibid*, 102); *S. erythrea* Foulerton (*Ibid*, 107) from culture tubes prepared with horse serum; *S. beta* Jones (*ibid*, 109); *Cladothrix invulnerabilis* of Acosta and Grande Rossi (ref. *Centralbl. f. Bakt, etc.* (1893), **14,** 14); the *Streptothrix* of Kedzior (*Arch. f. Hyg.* (1896), **27,** 328) isolated from sewage and water; *S. albido-flava*, *S. carnea*, and *S. aurantiaca* of Rossi-Doria (*Ann. d'Ist. d'ig. d. Roma* (1891), **1**) isolated from air; *Micromyces hoffmanni* described by Gruber (ref. *Centralbl. f. Bakt., etc.* (1891), **10,** 648) isolated from air.

Several other brief descriptions of *Streptothricæ* from pathologic lesions might be noticed, but the principal ones have been mentioned above. The technical descriptions have for the greater part been omitted in this review of the literature, but it may be stated that a classification of this material which will be satisfactory to all is scarcely possible. However, several authors have attemped an arrangement of species and varieties.

Foulerton worked with a large collection of cultures obtained from various sources. He redescribed and renamed several species, both pathogenic and non-pathogenic, the classification being based largely upon morphologic and cultural characteristics. Rossi-Doria occupied himself for the greater part with *Streptothricæ* cultivated from extraneous sources and his classification is based upon the character of the cultures obtained. Gaspereni gave a classification based upon cultural characteristics. Brumpt (*Arch. d. parasitol.* (1906), **10,** 489) recognized several species and more than one genus. Wright worked with one species and gave the most complete description of that organism to be found in literature. His "*Hyphomycete*" cultivated from a case of the "black variety" of mycetoma evidently was not a *Streptothrix* and therefore is not included in

the discussion in this paper. Caminiti recognizes forty-one species based upon his own work and a review of the literature. Manson's classification refers particularly to mycetoma and has already been noticed.

If we carefully study the literature of species determination, it will be found in greater part to be only of historic interest and value. The work is generally too incomplete to be final for species determinations, and in cases even the generic position of the described parasites is doubtful. For this reason we have considered only the technical descriptions of the organisms which have been cultivated and described with sufficient accuracy and completeness to make them available for comparative purposes. Much of the work of the older writers was well done when due consideration is given to the technique developed at the time of publication, and this work should always be kept before us in giving historical summaries. However, for purposes of further study and for comparison, having in view the harmonizing of our ideas regarding these infections, it seems to us that the following are the principal publications regarding *Streptothricæ,* which have been isolated from pathologic lesions of man or animals.

1. Nocard (1888); 2. Eppinger (1890); 3. Almquist (1890); 4. Bostroem (1891); 5. Wolff and Israel (1891); 6. Hesse (1892); 7. Vincent (1894); 8. Boyce and Surveyor (1894); 9. Aschoff (1895); 10. Du Bois Saint-Sevrin (1895); 11. Sabrazès and Rivière (1894) (A and B); 12. Rullmann (1895); 13. Dor (1896); 14. Sawtschenko (1896); 15. Scheele and Petruschky (1897); 16. Bucholtz (1897); 17. Berestnew (1897); 18. Rullmann (1898); 19. Silberschmidt (1899); 20. Brault (1899); 21. Cozzolino (1900); 22. Dean (1900); 23. Aoyoma and Miyamoto (1901); 24. Jelenewski (1901); 25. Foulerton (1902) (3 or more species); 26. Trolldenier (1903); 27. Pinoy (1906). The work of Musgrave and Clegg (1907) might also be included in this list.

The twenty-eight species or varieties to be taken from this list are considered with due regard to the manner in which they are described by the authors. It is certain that they do not all represent different species and it is probable that some of the other organisms not included do belong to new species. Many of the articles in the literature are incomplete, and a careful study of such descriptions as are given convinces us that not more than ten different species are represented in the group. Several of the varieties may show slight morphologic and biologic differences, but these are not sufficient to warrant their being classed as separate species.

GENERAL DISCUSSION AND CONCLUSIONS.

The subject of streptothricosis or nocardiosis becomes somewhat simplified if the results of our own work are combined with those taken from the literature, but we realize fully that after considering all possible evidence, points must still remain open for discussion, and complete harmony regarding this subject can only be obtained by common adoption of a somewhat arbitrary classification.

DETERMINATION OF GENUS.

It is difficult to determine upon the correct name of the diseases, because of the botanical confusion regarding the position and designation of the group of organisms concerned, and of the lack of clearness of definition specifying definite limitations for the organisms to be included as the etiologic factor.

Both *Streptothrix* and *Actinomyces* are untenable as generic names if we strictly follow the rules of nomenclature, because they are not entitled to priority in the literature which belongs to *Nocardia;* besides both *Streptothrix* and *Actinomyces* are open to the objection that they have not received a sufficiently clear botanical definition and both names have been used in too uncertain a sense in the past. *Actinomyces* has also, unfortunately, been taken into recent literature with practically two definitions. The objections to *Nocardia* are as follows: (1) The published definition erroneously states that the branching is "false branching;" (2) the name has not been sufficiently accepted by botanists to insure permanency and (3) usage is very much against it, particularly as in some of the recent writings its meaning has been limited to include only a portion of the organisms which surely belong in the genus. Considering the confusion in botanical literature and the uncertainty, and therefore the liability to change which pertains to this nomenclature, we have, as stated in the introduction, chiefly because of usage, and therefore somewhat arbitrarily, tentatively accepted *Streptothrix* as the generic name of the organisms concerned, and streptothricosis as the designation of the disease caused by this group of branching, filamentous organisms. In making this decision we are fully aware of the rights of those who favor *Actinomyces* or *Nocardia,* and under the circumstances are tempted to introduce a new name (*Carterii*) for the genus, together with a full and complete definition.

Whatever the nomenclature, it is a fact that we have here a group of closely allied vegetable parasites of man and animals, which have the following principal characteristics:

Branching, filamentous organisms which develop into colonies made up of the organisms and "transformation products." The terminal hyphæ may or may not be radially placed on the surface of the colony and they may or may not develop "clubs." The group in general take Gram's stain and several members show acid-fast properties in a varying degree.

The organisms grow on artifical media, differing in their requirements for oxygen and in pigment production. To a less degree they show other variations in appearance on artificial media.

The majority of the organisms produce lesions in monkeys, which histologically resemble those found in the human infections and in those of other animal diseases caused by members of this genus.

This group or genus is closely allied to other genera of somewhat

similar characters, the latter ranging from the branching bacteria through the nonbranching filamentous *Leptothrix* and the pseudo-branching *Cladothrix,* to the budding *Oïdia* or *Blastomyces.*

The only question of doubt regarding the expediency of our generic classification is raised by the very thorough and exhaustive work of James Homer Wright, who has brought forward some strong arguments in favor of making two genera out of the group of organisms under discussion.

As may be remembered, Wright separates this group into two genera; *Actinomyces* and *Nocardia.* He defines *Actinomyces* accurately and places all the organisms which he thinks do not come within this definition into the genus *Nocardia.*

Actinomycosis is defined as:

"A suppurative process combined with granulation tissue formation, the pus of which contains characteristic granules of 'drusen' composed of dense aggregates of branched filamentous microörganisms and their transformation of degeneration products. In the term transformation products are included the characteristic, refrangent, club-shaped bodies radially disposed at the periphery of the granule, for these bodies have long since been clearly shown to arise by a transformation of the peripheral filaments. They may or may not be present at the periphery of the granule."

Wright, in his further discussion of the organisms, emphasizes the following points as being characteristic of *Actinomyces.*

Club formation in tissues, anaërobism, granule formation, peculiarities in culture media, morphology and action when injected into laboratory animals. However, some of these distinctions were not constant even in the author's series of cases.

If we accept Wright's definition of actinomycosis, some of the organisms termed *Nocardia* by the author must be included as etiologic factors in actinomycosis, because the requirements are fulfilled by microörganisms not included in his description of the organism which he considers the specific cause of the disease, and furthermore, if his definition and description of *Actinomyces* be accepted as sufficient for genus determination, other genera could be removed from the group with almost, if not with equally good cause. However, some of the principal characteristics given for *Actinomyces* are not peculiar to his organisms. Most of these are differences in *degree only* from other members of the group, and are not specific, and in some instances, as we have demonstrated by work with original cultures, Wright's conclusions, based upon a study of the literature, are wrong. His observations regarding *S. caprœ* Silberschmidt, and *S. maduræ* Vincent are examples of this. Wright states that Vincent's case was probably one of actinomycosis, and that Silberschmidt worked with mixed cultures. We have had cultures from the original strains of both these organisms in our hands and are able to state positively that the bacillus- and coccus-like

forms of *S. capræ* described by Silberschmidt are forms of the pure culture and not bacterial contamination. As to Vincent's case, the organism is distinctive and can not possibly be included in Wright's definition of *Actinomyces;* indeed, it forms one end of the *Streptothrix* group, while the *Actinomyces* of Wright forms the other.

SPECIES DETERMINATION.

Species determination in this genus is made extremely difficult, and classification of all the forms which have been given in the literature is impossible, because of the incomplete and imperfect descriptions which are given. Several of these descriptions are old, have not been repeated or confirmed, and are lacking in the elucidation of important points which are now recognized as essential for diagnosis. Our correspondence has elicited the fact that cultures of several of these species are no longer obtainable, and therefore, no further study of them can be made. Fortunately, we have been able to secure cultures of the most important members of the group and by comparative morphologic and biologic studies, particularly with reference to their pathogenic character, we have been able somewhat to simplify the classification by showing that some of the described species are identical. This includes *S. freeri,* which from a study of the literature we thought to be a new species, but after comparative study found to be identical with *S. eppingeri* in all essential particulars.

Another difficulty in establishing species or varieties is encountered in attempting to fix the amount of variation necessary to constitute a new species. The differences between various members of the group are practically in every instances those of degree only. For example; in the requirements for oxygen we have variations from facultative anaërobism with strong tendencies toward being negative to oxygen, as exemplified by the Wolf, Israel and Wright type of *Actinomyces,* to facultative aërobism with strong oxygen requirements, as exemplified in *S. eppingeri;* while other strains show intermediate stages of aërobiosis. Although strict anaërobes and equally strict aërobes have been mentioned, the conclusions probably are based upon errors in technique. Certainly this is true with the organisms with which we have worked, for in no instance have we encountered either a strict anaërobic, or an equally strict aërobic *Streptothrix,* although some of the cultures under our hands have had such properties ascribed to them by others.

Staining reactions.—The terminal filaments in lesions and young cultures usually stain very well by the usual aniline dyes, but in older colonies the central detritus does not stain and the filaments give irregular staining reactions often showing irregular, including coccus- and bacillus-like, forms. The reaction to Gram's solution is somewhat variable, depending upon the age of the colonies and there is some variation

between the different species or varieties. However, in general, all strains are Gram positive to a greater or less degree.

Differences in *acid fast* properties have been used as being diagnostic in a number of described species, while unfortunately in others they have not been mentioned. These are important points in differentiation, but unfortunately such properties are a somewhat variable manifestation, depending in part upon the age and environment of the parasite and perhaps also upon other conditions, the nature of which is not clear. As with the oxygen requirements, degrees from the marked acid fast properties of *S. eppingeri* through those organisms showing the tendency only in certain portions, to *S. maduræ* which has no such properties are noted. *S. eppingeri,* often when taken from animal lesions and sometimes for example in cultures, will show acid fast properties which involve the entire organism, while in other cultures only portions of the organism will retain the fuchsin when decolorized and counter stained with Gabbet's stain.

Pigment production under certain circumstances is a property of most of the organisms of this group, as the cause of this is unknown and the result is variable, its diagnostic importance is lessened. Under like conditions, the color and general appearance of the pigment in any of these strains is usually fairly constant, but sometimes, particularly in animal tissues, it may show a more intense color. For example, *S. maduræ,* which usually produces a reddish-pink pigment in cultures, may, when inoculated into monkeys, at one time give a similarly colored pigment or again one considerable darker; so dark in fact as to appear almost black to the naked eye. The pigment of *Actinomyces* in our cultures shows but little color other than black. On media the color develops slowly and gradually to a dark gray, almost black. Some of the phenomena of pigment production and color may be explained, but in general the process is still obscure. For example, a darker pigment usually results in animals from the inoculation of fresh material from another animal than is the case if cultures are used. The next lighter shade is produced by inoculation with old, pigmented cultures, but color production is slower when young, unpigmented cultures are used for inoculation. Early pathologic lesions due to these parasites, like young cultures, rarely show pigment. The later in both instances is produced and grows more marked with age. Concentration undoubtedly has something to do with the apparent color of the pigment, for the harder the granule the darker the pigment appears to be.

Granule formation both in tissues and to a less extent, in cultures, is a property of all the pathogenic *Streptothricæ* with which we have worked. These granules vary considerably in consistency and color in the several species, but in the majority of instances they are made up of branching filaments with transformation products, which consist in irregular forms,

crystals and unstainable detritus. The arrangement of the peripheral filaments may or may not be radial and there are various degrees of density of the mass. In two of our cultures (*S. canis* and *S. caprae*) there is less tendency to suppuration after inoculation in animals and the individual colonies have more the appearance of tubercles than in our other cultures. This difference has been pointed out by other observers. However, the lesions in our experimental animals were small; in the older lesions the type of the granule is not known, if we except the knowledge gained from the statements of the authors who described the parasites, the description being taken from the original lesions. The granules or colonies produced by the other cultures were in the majority of cases free in the necrosed, liguefied substance found in the various channels produced by these infections.

Club formation, in the lesions, in tissues and less frequently in cultures, is used as one of the points of differentiation between species of the organisms of this group. No doubt there is considerable variation in this respect under certain conditions, but the circumstances reported as surrounding club development are sufficiently varied and confusing to detract somewhat from what would otherwise be valuable for purposes of distinction.

Wright found club formation to be rather constant in original lesions, and observed it occasionally in certain cultures. In experimental lesions, it was also fairly constant, except in early lesions where clubs were sometimes absent.

Bostroem and others, working with species differing somewhat from Wright's organism in other particulars, also observed clubs both in lesions in animals and occasionally in cultures.

None of our cultures show definite clubs in culture or in experimental lesions in animals, although in several the terminal branches show some enlargement suggesting club formation (see illustrations). Our experimental animals were all killed within a shorter time after inoculation than were those of Wright, in which clubs were found.

Branching is similar in all of our cultures and is also similar to that recorded by most observers who have worked with this group of microörganisms.

The cultural characteristics of all of our strains have already been described. While there are variations in the reactions between some of these strains, they all appear to belong to one group of organisms, and if we analyze the literature carefully others, including the organisms of Wright and of Bostroem, may also be included in the group.

The following may be noticed if we take up the principal cultural characteristics comparatively as well as somewhat more in detail.

Surface growth is considered to be poor in Wright's organism and is not very profuse in some of our strains, but in Bostroem's organism and in most of ours, while it may be slow, it does occur in a satisfactory manner and in some strains

it may be luxuriant. There is but little tendency for surface growth to spread in any of the cultures, but on the other hand, it heaps up, as it were, giving raised colonies sometimes 1 centimeter above the surface of the media. (See plates.) This is also true with Wright's organism, where surface growth appears.

Wright first called particular attention to the character of growth in glucose-agar suspension cultures. In this medium with his organism, there is a narrow zone 5 to 10 millimeters below the surface where the colonies are very numerous but small, while lower down they are less numerous, but of larger growth. Surface growth is the rule in our cultures in this medium, and only occassionally a colony may develop in its depth.

Stab cultures in sugar-agar.—Growth with Wright's organism occurred in small nodules along the course of the needle, it did not penetrate the medium to any extent and did not grow on or near its surface.

The growth in our cultures occurs mainly upon the surface and upper portion of the track of the needle.

Anaërobiosis.—Wright considers his organism as an obligate anaërobe, a statement which is somewhat qualified in the details of his work. For example, in discussing the growth upon glucose-agar suspension media he states that the most profuse growth of the organism is in a narrow zone 5 to 10 millimeters below the surface of the medium and that this way may be "explained as the result of a stimulating action exerted upon the growth of the microörganisms by the presence of a small amount of oxygen which has penetrated into the medium from the surface." This oxygen requirement is further illustrated by the author when he states that "in sugar-agar suspension cultures, placed under anaërobic conditions, there is little or no tendency to form colonies on the surface."

Growth occurred along the course of the needle only in stab cultures on glucose-agar. It did not penetrate the medium and, on the other hand, did not grow on the surface. However, when discussing growth upon surface inoculations in slanted solid media, Wright found that surface growth was not luxuriant and in some of his strains no growth occurred even under anaërobic conditions. Again, Wright observed that "in general, growth appears to be as luxuriant in bouillon either under aërobic or anaërobic conditions." To judge from the above and other statements which might be taken from Wright's excellent monograph and *according to the standards used in this report* for the other members of the group, Wright's organism must be considered to be a facultative anaërobe.

In *bouillon* media there is some variation between the manner of growth of our cultures, but little which may be considered of value in the differentiation of species. The tendency in some of our cultures is toward a surface growth which may gradually separate and fall to the bottom, or settle on the sides of the tube. In other instances, the majority or all of the growth may occur at the bottom of the tube. The medium is not clouded, but in some cases it becomes darker, apparently because of the absorption of pigment from the organisms. There are no essential differences between our cultures in bouillon and those reported by Wright.

Potato, as is well recognized, is not a very satisfactory medium for comparative studies, excepting where the same set of media is used. Fairly luxuriant growth was obtained in every instance with our cultures on potato slants.[2]

[2] The potato slants employed in this laboratory are made from selected potatoes treated and prepared according to the rules laid down by the committee of the American Public Health Association.

There was not much variation between the different cultures except in the color of the pigment and the appearances which have already been described. Potato proved itself to be a very unsatisfactory medium for Wright's organism. Most of his strains grow on it very poorly, and some of them not at all.

Litmus-milk is not a favorable medium for the differentiation of species. Growth takes place slowly, and with two strains the color is slowly discharged. No other appreciable changes occur in the medium excepting with cultures of "chalcea," in which a slightly acid reaction is produced.

The favorable influence of *sugar* and *glycerine* upon cultures of all strains of the group of organisms has been noted repeatedly. Sugar or glycerine containing media not only are the best in securing growth, but with all our species pigment production is more active in these media.

Pathogenic characters.—The pathogenic character of many varieties of this group of organisms in the human and animal economy is well established. However, experimental inoculations in animals have not given constant results in the hands of other investigators, and our experiments have been uncertain with all animals except monkeys. The details have already been described and it only remains to state here that lesions similar in their gross and histologic appearances to the original ones produced by these organisms, have resulted with each of the strains which we have studied, except with one of the strains of *S. Madurae* which has already been mentioned in this discussion.

CONCLUSIONS.

There is, in human pathology, a very important group of branching, filamentous microörganisms which logically belong to a single genus. The generic name is variously given as *Streptothrix, Actinomyces* or *Nocardia;* the last of these names is probably scientifically the most correct, but because of the present botanical confusion and uncertainty the first is here employed, because of its more general acceptance.

The following species to judge from our work and from a study of the literature are the most important and may be recognized as established. There are probably a number of others but the description of many of them are too imperfect to allow of their recognition.

S. actinomyces Bostroem, 1890.
S. actinomyces Wolff & Israel, 1891 and Wright, 1905.
S. nocardii.
S. eppingeri.
S. maduræ Vincent.
S. capræ Silberschmidt.

The disease caused by infection with these parasites is properly named streptothricosis, with actinomycosis and nocardiosis as synonyms. Other names, such as lumpy-jaw, madura foot, mycetoma, etc., should be considered more as describing anatomic location rather than as designations relating to any special or specific cause of infection. Mycetoma might well be taken as the correct name for the group of infections if a strict interpretation of rules of nomenclature is followed, but usage renders it perhaps more desirable to retain the name as representing

Streptothricosis pedis. If mycetoma is to be considered in any more comprehensive light than this, it should become another synonym for streptothricosis. It should not be considered a disease caused by organisms other than *Streptothricæ*.

PART II.

STREPTOTHRICOSIS.

SYNONYMS.—Actinomycosis, Nocardiosis.

DEFINITION.—An infectious disease of man and animals caused by one or more species of *Streptothrix*. It is characterized anatomically by a peculiar, low grade of inflammation, usually confined to one part of the body, but in rare instances assuming the proportions of a general infection. The inflammatory process usually is accompanied by suppuration of a certain kind; the discharges contain granules made up principally of colonies of *Streptothrix*. The general picture of chronic inflammatory disturbance is seen clinically, and enlargement, suppuration and the presence of the causative organism in the lesions is observed locally.

HISTORY.—The history of *Streptothrix* infection is shown in the review of the literature given in Part I. In general it may be stated that the disease has probably long been recognized and frequent references are made in older medical literature to conditions which presumably were due to infection with these microörganisms. According to Rivolta, Trutto discussed the condition among cattle in Italy in 1785 under the name of "Krotenkrankheit." The colonies of the organisms were more or less accurately described by Langenbeck (1845), Sir T. Smith (1855), Lebert (1848), Rivolta (1868), Robin (1871), Heller (1872), Perroncito (1875), and others (see Ackland). Mention of the anatomical form of the disease generally described as mycetoma or Madura foot are also quite old, the principal authors being: Kämpfer (1712), Heynes (1806), Brett (1840), Gill (1842), Godfrey (1844), Colebrook (1844), Gunther (1844), Eyre (1848), Bollingol (1855), Eyre (1860), Collos (1861), Bidie (1862), Hirsch (1868 and 1886), Coquerel (1866), and many others (see *This Journal, Sec. B.* (1907), **2**, 487). Of these observers Colebrook (1850), Bollingol (1855), Coquerel (1866), Maxon and Hogg (1870), Bristowe (1871), Berkeley (1876), and others probably saw and in a few instances partially described the organisms present in the lesions. However, the really important history of *Streptothrix* infections dates from Vandyke Carter's work (1859–1874). This author described and illustrated the parasites in many cases of the foot (Madura foot) type of the infection. Bollinger and Marz (1876) first accurately described and named one of the parasites causing the disease known as lumpy-jaw in cattle. The next epoch in the history was due to an article by J. Israel (1878) who was the first to describe the disease and parasite in man in locations other than the foot (Madura foot) and

hand, the infection in these places already having been described by Carter and others. Israel's work was followed by that of Ponfick (1879) with strong evidence as to the identity of the disease in man and animals. From the year 1876 the literature accumulated rapidly and the infection was reported for man and animals from widely varying locations and countries.

Bostroem (1890) contributed the first thorough and exhaustive laboratory study of the microörganisms and this work was followed a year later by an equally exhaustive treatise on the same subject by Wolf and Israel. The organisms described by Bostroem on the one hand and by Wolf and Israel on the other, although isolated from apparently similar diseases, showed certain morphologic and biologic differences which led to much discussion during the next few years. Organisms corresponding to the descriptions of both these pioneer observers continued to be mentioned in the literature, each investigator maintaining his organism to be the true cause of the disease.

In 1888 Nocard cultivated and quite accurately described a *Streptothrix* as the cause of a disease locally termed *farcin de boeuf* of cattle in Guadalupe; Nocard's organism seemed to be of a different species from either Bostroem's or Wolf and Israel's. Eppinger (1890) cultivated and accurately described another species of *Streptothrix* as the cause of a brain abscess and "psuedotuberculosis" in a man. Vincent (1894) first satisfactorily cultivated and described a *Streptothrix* as the cause of mycetoma or Madura foot and his organism seemed to have certain characteristics showing it to be specifically different from any of those previously described.

Foulerton (1899–1907) gave descriptions of several new species of the parasites and has brought out the most thorough and exhaustive consideration of the whole subject with which we are familiar. Wright (1905) contributed a most complete and exhaustive study of the biology of the Wolf and Israel variety of this organism. The work of bringing all this group of similar organisms together in a single genus is largely due to the researches of Rossi Doria, Petruschky, Foulerton and several other recent writers, particularly of the German and French observers.

In this article we have shown conclusively that Eppinger's organism isolated from clinical actinomyces on several occasions is identical with *S. freeri*, which is one cause of Madura foot in the Philippine Islands. Very little is known regarding *Streptothrix* infections in this Archipelago. Our records show two cases in which *Actinomyces* was the etiologic factor in human disease, and several infections of cattle have been observed in the islands. One case of the Madura foot type of the disease has been seen.

Etiology.—*Predisposing causes*, for this disease, except in a few instances, have not been sufficiently studied to make a satisfactory estimate of their influence.

The *geographic distribution* is very wide, but no very large series of cases have been studied by any one observer. Collections of such series have been made in America, Germany, France, England, Russia, Switzerland, and in the British colonial possessions. The most frequent incidence is found in certain parts of the Tropics, particularly in India, but, as has been pointed out by Ackland, the disease may be overlooked in a community for a long time until special attention is called to its prevalence.

According to available statistics the infection is more frequent during young, adult life, but cases have been reported in children and in old people. Males are much more frequently affected than females—295 males to 110 females being given in Leith's statistics, and 65 males to 36 females in Ackland's.

Race, nationality, residence, overcrowding, climate, occupation, and physical condition, to judge from available statistics, seem to exert but a minor influence on the spread of the disease, although a more careful study of a large material may show some of these to have greater significance.

General environment and *mode of living* would naturally be expected to have considerable influence as predisposing factors, particularly in determining the part of the body involved. For example, the Madura foot type of infection is most common among people who go barefoot, it being largely a wound infection.

Streptothricæ are the specific, causative agents of the infection. They have already been considered in Part I of this report, and only a brief summary will here be given.

Synonyms.—*Actinomyces, Nocardia, Oöspona, Chenaphe carteri.*

Diagnosis of these organisms is made from morphologic and biologic considerations. They are branching, filamentous organisms, which develop slowly into colonies made up of the branches and their "transformation products." These colonies vary in color, size and consistency, and when stained show various changes in different portions. The filaments at the periphery are usually intact, with or without club formation, and the terminals may or may not be radially placed. Toward the center of the colony, or granule, irregular forms, such as coccus- and bacillus-like ones, are found, together with crystals and nonstaining detritus.

The majority of these organisms may be cultivated on artificial media, where they show various but characteristic biologic properties. That some of them are pathogenic tissue parasites is shown by their action on laboratory animals.

Morphologically these parasites are rather closely related to some of the branching bacteria. The young filaments vary in width from 0.5 to 1 μ and in length from 5 to 20 μ or more. They usually stain homogeneously, and in some strains taken from lesions they are acid-fast to the

Ziehl-Neelsen-Gabbett method. In older forms the segments are often broken, the sheath-like substance not showing either in fresh or stained specimens. The breaking up of the filaments produces great variety and size of irregular forms; some of these may appear to be very much like bacteria, but their true nature is easily established by laboratory methods. The majority of these organisms may be stained by the aniline dyes, all show Gram positive characteristics to a greater or less degree, and several are as acid-fast as is the tubercle bacillus.

Biologically certain species—both saprophytic and pathogenic—are widely distributed in nature. They have been found on food stuffs—particularly cereals—in water, air, soil, etc. Zoölogically they are found in some insects and several animals, and in man. Cattle are the most frequently infected of the animals, but rarer instances of the disease have been reported for other species.

Cultivation of these organisms is probably possible with all species, although several authors have reported failures, and all observers have noted the difficulty of obtaining a culture from the lesions in animals. However, when adaptability to artificial media has once been secured, no very great difficulty is found in keeping the culture. In general, it may be stated that these organisms grow slowly on artificial media, but that the cultures show most positive characteristics in the majority of instances. In all there is a tendency to "pile up" on the medium rather than spread over the surface. Pigment production of various shades is a very common property, and this is more marked as a rule in glycerine or glucose containing media.

Resistance to physical, chemical, and other agents by these organisms is rather great if we consider them to be non-spore bearing parasites, but this resistance is not sufficient to permit us to consider that true spore formation is present as it is understood for bacteria.

Pathogenic character.—The pathogenicity of several species of *Streptothrix* has recently been demonstrated by different workers, according to the accepted bacteriologic rules governing such decisions. Other members of the group appear to be saprophytes in that they do not produce lesions in animals by the usual methods of procedure. There is also considerable lack of uniformity in the results of animal experiments with nearly all the pathogenic strains. However, working with monkeys, our results have been more uniformly successful than those reported by other observers, or than our own results with other laboratory animals. The pathogenicity of this group of organisms seems to be influenced to a certain extent by the same conditions which produce changes in the virulence of bacteria. Not only may the virulence of some of these species be increased by passage through susceptible animals, but with the increased virulence differences in cultural results may be noticed. These are shown principally in greater difficulty in securing growth, in

a slower growth on artificial media and sometimes in slight changes in the color and quantity of pigment production.

Some of the so-called saprophytic species may in reality be found to be pathogenic, when improved technique is used.

Modes of transmission.—Although experimental evidence is still far from satisfactory regarding modes of transmission, it seems to be entirely reasonable to assume from the known biology and the available evidence, that transmission may take place both directly and indirectly. So far as we are aware, no positive examples of direct transmission from person to person have been reported, but that such infection does not take place in localities with high incidence of the disease, seems probable. Rather convincing evidence of indirect transmission through the agency of food-stuffs, water, etc., has been furnished by several observers. Several authors whose work has already been reviewed, have shown that pathogenic species may be isolated from water, air, soil, and food-stuffs and the history of many of the reported cases indicates infection from some such source. The frequent presence of the infection in the lower animals must not be forgotten in considering the manner of transmission of the parasites.

Infection probably takes place in two general ways, first, by direct or wound infection, as is shown in many cases by the history of injury before the development of the symptoms of the *external* forms; and secondly, infection by way of the respiratory or gastro-intestinal tracts, which is shown in many of the *internal* forms of the disease.

Species.—The number of species important in this disease is probably a considerable one. Those which we have been able to recognize from the literature and our own work are given in Part I.

It seems probable that all the species have not, as yet, been discovered and that some, which have been described, are specifically different from the ones mentioned here.

Pathology.—In general the morbid changes found in this disease are those of a peculiar, low-grade, chronic infective process. Anæmia, atrophy of tissues, and mild, chronic degenerations of parenchymatous organs are present. The special pathology depends somewhat on the part of the body involved and on the extent of this involvement. The streptothritic unit, as it were, is a granule which is usually surrounded by a zone of peculiarly appearing suppuration, and this in turn by an area of inflammation characterized by connective tissue proliferation and cell infiltration. Cells of the usual character are found, with a relative increase of fixed tissue cells and occasionally a giant cell. In other instances the suppurative zone is absent and the streptothritic unit takes on more the general and histologic appearance of a tubercle. In fact, the whole morbid process of *Streptothrix* infections resembles more or less closely that produced by *Bacillus tuberculosis*. Tissue destruction often spreads

by means of continuity, but more usually by intercommunicating channels running in various directions. These channels contain the broken down tissue and granules made up of colonies of the infecting organism. Few tissues escape the destructiveness of the *Streptothrix,* but it appears to be more active in the connective tissues, bones, and mucous membranes. However, practically every organ and tissue of the body has been involved. The disease spreads through the body in at least two ways, by directly continuing along the tissues, in which case the area involved is rarely great, and by metastases through the blood vessels, when the infection may be so great as fairly to be considered a general infection.

The character of the morbid process suggests that the minute action is toxic in character, although if such is the case, the toxin must act very slowly. The lymphatics near a diseased process may be enlarged, but are rarely found to be broken town or to contain the microörganism.

Symptoms.—The clinical manifestations of *Streptothrix* infections are essentially those of a slowly developing, chronic inflammatory process. They vary much with the location of the lesions and further complications are secured by the presence of mixed infections which are frequent when the disease attacks regions of the body exposed to bacterial invasion. In many instances, where the lesions are located in the internal organs such as the liver, definite clinical manifestations do not occur and unless the infection spreads, its nature in all likelihood is not recognized during life.

When the lesions are located externally, or in places where the discharges reach the surface of the body, the nature of the infection should be suspected from the appearance of the wound and the character of the discharges; on the other hand, it may be generalized, manifest itself in an acute course and be difficult to distinguish from a pyæmia of other etiology.

The *incubation* period varies between wide limits, and as given by authors may be from a few days to more than two years. The onset is usually gradual, by the slow development of the lesions.

The following clinical varieties may be recognized:

Generalized streptothricosis.—This form of infection has been noted by several observers. It occurred 9 times in Ackland's 109 cases. This type usually begins in a local lesion and is transmitted through the blood vessels. In the majority of instances, it occurs in mixed infections with bacteria, but cases in which no bacteria were present have also been reported. Several species of the organisms causing these general infections have been found. This type is generally acute or subacute clinically and the symptoms as given by various authors are similar to septicæmia or pyæmia from other causes. Extensive lesions may occur in practically every organ and tissue of the body. The diagnosis is only made by laboratory methods, the prognosis is bad and treatment unsatisfactory.

Thoracic streptothricosis.—The organs of the chest, particularly the

lungs and pleuræ are most frequently involved in this disease. It occurred 65 times in 257 cases reported by Duvan and 29 times in Ackland's 109 cases. The clinical symptoms in the lungs and the physical signs may resemble a chronic tuberculosis or severe bronchitis and the pleural involvement may consist of chronic, adhesive pleuritis or more often of empyæma. Unless there is perforation of the chest wall or metastases, the clinical manifestations in these types show nothing characteristic of the disease and the true nature of the infection can only be determined by laboratory methods. Streptothricosis of the heart and mediastinum have been reported, and Ackland notes nine cases involving the œsophagus.

Abdominal streptothricosis, including all the organs of the abdominal and pelvic cavities, have been reported by various observers. The liver and gall bladder suffer most often. Involvement of these viscera occurred forty times in Duvan's series and thirty-three in Ackland's. The appendix is quite frequently involved and in some instances is the primary seat of the lesions.

Other lesions are found in the mesentery, intestine, rectum, spleen, bladder, prostate, testicle, kidneys, and abdominal wall. In none of these types are there any characteristic clinical manifestations which differentiate this infection from other chronic inflammatory conditions of tuberculous or other etiology in the same locations.

Cerebral streptothricosis includes involvement of the brain, cord, meninges, and nerves, as well as the other tissues in these organs. The central nervous system was involved 19 times in Duvan's 257 and 5 in Ackland's 109 cases. Several different species of the organisms have been found in these lesions, that of Eppinger being the principal one. This species, as has been shown in Part I of this report, is identical with *S. freeri* which caused the mycetoma in our case. The reported cases show no characteristic symptoms in these types of infection.

External streptothricosis may involve any of the external or contiguous tissues, including the mouth and jaw, neck, skin, and subcutaneous tissues, conjunctiva, nose, and extremities. "Lumpy-jaw" or "big-jaw" is one of the most frequent of these various types; mycetoma, Madura foot or *streptothricosis* pedis is a very common tropical type and others involving many anatomical locations are frequently encountered. Practically all described species of the genus of the organism have been found at one time or another in external streptothricosis, and there does not appear to be much variation in the clinical picture because of any special species or variety. The clinical manifestations of external streptothricosis are rather characteristic. The peculiar, crater-like skin lesions of a chronic character leading through ramifying channels, through which is discharged the peculiar, oily, pus-like substance containing granules, makes a rather constant and characteristic picture. As we have already stated, mycetoma or Madura foot is best classified here as the clinical

type of *Streptothricosis pedis,* because it may be caused by several if not any of the species of this genus and the clinical manifestations are practically the same in all cases. The special symptoms of this type are well known and have already been discussed in this report.

Miscellaneous types of streptothricosis may be mentioned, such as involvement of the conjunctiva, bones, ear, nose, intercostal spaces, etc. Their only peculiarities are in the anatomical locations of the lesions.

Diagnosis.—The diagnosis of *Streptothrix* infections is made clinically by the character of the exposed lesions and discharges and by laboratory methods. Obviously, the common and frequent involvement of internal organs is rarely recognized except by microscopic examinations at operations or at autopsy. The thoracic types may be determined by microscopic and bacteriologic studies of the sputum or aspirated fluid from the pleural cavities. It is generally stated by competent observers that these types are frequently overlooked, even in sputum examinations and it is not improbable because of the acid-fast properties of some of the organisms that they may occasionally be mistaken for the tubercle bacilli. The resemblance may, at times, be quite close. In some of our experimental lesions the similarity has been striking. The microscopic examination of material from exposed lesions often requires some patience and care to enable the observer to find the organisms; unless a granule is encountered, when, of course, the determination is easy. Inasmuch as some other closely related organisms such as *Oïdia, Leptothrix,* and *Cladothrix* may cause somewhat similar lesions, *Streptothricæ* should not be diagnosed positively without careful microscopic study of the organisms present in the lesions.

Prophylaxis.—From what we know of the distribution of the organisms of this disease and the mode of transmission and the prevalence of certain anatomical types, prophylaxis should consist in guarding against wound infection and generally by care in food and drink. In the Tropics most of the types consist in local foot or other skin wound infections, which should be guarded against by wearing shoes and by promptly treating skin wounds and abrasions according to antiseptic methods.

Prognosis.—The prognosis depends to a considerable extent upon the location and extent of the lesions. The general infections and those of the internal organs almost always end fatally. The external types are much more amenable to treatment and the mortality is small.

According to the statistics of Duvan, and Poncet and Berard who analyzed 257 cases, the mortality was as follows: Skin, 2.3 per cent; face and neck, 10; jaw and temporal region, 30; abdominal cavity and intestine, 65; thoracic, 85; liver, 100, and brain and spinal cord, 100 per cent. The mortality in mycetoma, under proper treatment, is very small. The course of the disease is usually chronic, but it may be general

	Streptothrix eppingeri from Foulerton "A."		Streptothrix eppingeri from Foulerton "B."		Streptothrix nocardi.		Streptothrix maduræ from Strong.		
	Literature.	Our culture.	Literature.	Our culture.	Literature.	Our culture.	Literature.	Our culture.	
Morphology	Mycelium with branching threads ending in spores of an elongated appearance and also showing clubbed ends.	Various forms noticed, depending upon the age of the culture and media, from small oval to long branching filaments. The diameter is fairly constant. No sheath observed and no clubs.	Foulerton has not yet published the details regarding this organism.	Similar to Eppinger's original culture.	Long, thin, hyphæ twisted in masses showing many branching forms.	Many coccoid forms; short plump filaments with short branching hyphæ occur, in young cultures; in older cultures ovoid forms predominate.	This oganism has not previously been described.	Long, thin, filaments with true branching; no club forms.	Many twiste forms; th at the per of the hyp courses.
Staining characteristics.	Cultures from 2 to 3 months old stained by Ziehl-Neelsen method show many portions which are acid-fast but not alcohol-fast.	Portions of filament acid-fast by Ziehl-Neelsen method; ovoid forms were decolorized; spore-like bodies and portions of mycelium are Gram-positive.	.....	Similar to Eppinger's original.	Cultures from 9 days to 3 months old stained by Ziehl-Neelsen method show many acid-fast portions; these are also alcohol-fast.	Cultures stained by Ziehl-Neelsen method show many acid-fast portions. The organism is Gram-positive.	.....	Cultures stained by Ziehl-Neelsen method showed no acid-fast portions. Gram-positive with irregular staining.	Culture 1 we thod showe
Growth on ordinary agar.	Small whitish colonies which coalesce into a thick adherent membrane with elevations and depressions of an orange color.	Growth appears after 2 days as smooth, adherent membrane at first a porcelain-white, later a delicate orange color; medium remains moist.	.....	.....do.....	Small, white, irregular, round, opaque colonies which form an adherent, thick, wrinkled membrane-like lichen.	Small, white, irregular, round, opaque colonies; growth slow and not abundant on this medium; no pigment.	.....	Grows very slowly and very little pigment is produced.	Small projec which slow umbilicati and adher
Glycerine-agar.	.....	Growth appears after 3 to 4 days, at first as small whitish colonies which gradually take on a delicate pink color. Later as the colonies develop they become umbilicated and coalesce forming a heaped up growth and producing an orange color.	.....	.....do.....	.....	Appears after 3 days as discrete colonies; later the colonies coalesce and become heaped up, presenting a moist, mealy growth. This organism does not produce pigment.	.....	Growth appears after 5 days as small pinkish-white colonies; later the colonies develop a heaped up growth and assume a wrinkled appearance with a deep-pink color.	.....
Maltose	.....	Similar to glycerine-agar.....	.....	.....	.....	Similar to glycerine-agar.....	.....	Similar to glycerine-agar.....	Growth appe yellowish- surface; l
Potato	Growth appears after 48 hours as a granular layer, at first white, becoming yellow to brick-red. This is later covered by a fine, white, powdery efflorescence, culture resembling a sugar almond.	Growth appears after 48 hours as a granular layer at first white, becoming yellow to brick-red.	.....	Growth appears after 48 hours, similar to Eppinger's original.	Growth appears after 48 hours as a buff-yellow granular growth; no pigmentation or erosion of medium.	Growth appears after 48 hours as a buff-yellow granular growth.	.....	Growth appears after 3 days as small, pinkish-white colonies; these later develop a dark pink, raised growth.	.....
Boullion	Growth appears after 24 hours as a small, white granule, forming a pellicle and adhering to sides of tube; some portions fall to the bottom. No pigmentation of medium.	Similar to literature.....	.....	Similar to Eppinger's original.	Growth appears after 24 hours as a grayish, flocculent mass at the bottom of the tube.	Growth appears after 24 hours as a grayish, flocculent mass at the bottom of the tube; later a few grayish granules appear on the surface and in some instances puffball formation occurs at the bottom of the medium.	.....	Growth appears after 5 days as a coherent mass at the bottom of the tube; surface growth rare.	Growth app or less c tube; sur medium.
Ascitic fluid	.....	After 6 days very slight growth occurs on the surface of the medium; no pigment; no clubs.	.....	.....do.....	.....	Very slight growth at the bottom of the tube; no pigment; no clubs.	.....	Very slight growth; no pigment.....	.....
Pathogenicity for lower animals.	Not pathogenic for white mice. 3 rabbits inoculated subcutaneously recovered after developing temporary local swellings. 1 rabbit inoculated in anterior chamber of eye developed a nodule on the iris but no general infection. 1 rabbit inoculated intravenously and another in the kidney died on the fifth and seventh days respectively. Two guinea pigs inoculated intraperitoneally died on the eighteenth and twenty-fourth day.	Pathogenic for rabbits, guinea pigs and monkeys by intraperitoneal injection.	.....	.....do.....	Rabbits, dogs, cats, horses, and oxen are not affected by intraperitoneal or intravenous injections. In guinea pigs intraperitoneal or intravenous injections cause constantly within 20 days, a miliary tuberculosis similar to that produced by *Bacillus tuberculosis*.	Pathogenic for guinea pigs and monkeys by intraperitoneal injections.	.....	Pathogenic for monkey by intraperitoneal injection.	The inocula mice proc at the si (Vincent intraveno dogs, sh pigeons; case.

ultural characteristics of various types of Streptothrix and of Actinomyces.

Streptothrix maduræ from Vincent.		Streptothrix maduræ from Foulerton.		Streptothrix capræ Silberschmidt.		Streptothrix canis, our culture.	Streptothrix chalea Foulerton, our culture.	Actinomyces from Pasteur Institute.		Streptothrix freeri.	
Literature.	Our culture.	Literature.	Our culture.	Literature.	Our culture.			Literature.	Our culture.	Literature.	Our culture.
laments showing true branching ycelium shows a ray-like growth ry but no true club forms: some show an enlargement along their	Long, thin filaments with true branching; no club forms. These hyphæ may or may not be radially placed at the periphery of the granule.	Similar to *Streptothrix maduræ* of Vincent.	Similar to *Streptothrix maduræ* of Vincent.	Twisted, branching filaments which break off at the end into forms resembling bacilli or cocci. A few of the shorter forms show clubbed ends.	Short filaments resembling bacilli and many coccoid forms; occasionally longer filaments showing branching but no club forms.	Short, plump, filaments; many resembling bacilli; oval and coccoid forms.	Long, thin, filaments with branching hyphæ.	Ray fungus consisting of a mass of mycelia, containing many spores in center; branching forms present.	Radiating mycelium containing many spore-like bodies in center; true branching occurs.	Similar to morphology described.	Long, branching filaments, 2 to 7 μ in diameter. Transverse segments are shown of various length from coccoid to 10 μ in length. Branching occurs as lateral hyphæ developing from the segments. The filaments have a definite wall. Spores have not been observed. Coccoid and club-like forms present in certain medium.
ld stained by Ziehl-Neelsen me- acid-fast portions.	Cultures stained by Ziehl-Neelsen method showed no acid-fast portions. Gram-positive with irregular staining.	do	do	Irregular staining	Cultures from 2 weeks to 2 months old stained by Ziehl-Neelsen method show many acid-fast portions, giving the organism the appearance of short, plump bacilli. Gram-positive, irregular staining.	Culture stained by Ziehl-Neelsen method shows many acid-fast portions. The organism is Gram-positive in portions, giving it the appearance of cocci and short bacilli.	No acid-fast portions by Ziehl-Neelsen method of staining. Gram-positive for portions.	Culture stained by Ziehl-Neelsen method shows no acid-fast portions. This organism stains well by the Gram method.	Culture stained by Ziehl-Neelsen method shows no acid-fast portions. This organism stains well by the Gram method.		Culture from 5 days to 2 months stained by Ziehl-Neelsen method show many acid-fast portions. The organism is Gram-positive.
round, whitish-yellow colonies ake on a rose color; they show the center and become confluent the medium.	Grow very slowly on this medium; after 4 days small, whitish-yellow colonies develop.	do	do	In 2 or 3 days varicose, shriveled, brownish-white colonies appear which form an adherent, brown, irregular membrane in from 8 to 15 days.	Growth appears in 2 or 3 days as small brownish colonies; medium remains moist; does not grow so well nor does it produce the pigment on this medium that it does on the sugar-containing media.	Grows as a smooth, glistening, brownish-white colony; does not grow so well on this medium as on sugar-containing media.	Grows as dark-brown, adherent membrane.	Small, opaque colonies which later show a white, downy appearance and which coalesce into an adherent, wrinkled, lichen-like membrane.	Small, opaque colonies which later become black and in time produces a bright yellow membrane; the medium becomes dark.	Similar to morphology described.	From 2 to 3 days' growth appears as smooth, glistening colonies which later coalesce and produce a delicate pink pigment; medium remains moist.
	Growth appears after 5 days as small, pinkish granules and later assumes a dark-pink color.		Similar to *Streptothrix maduræ* of Vincent with the exception that its pigment is a more delicate pink with a pale periphery.		Growth appears after 2 days as small, brownish-white colonies which later coalesce giving the growth a moist, mealy appearance with a light-brownish color.	Appears after 2 days as small brownish-white colonies. These later assume an ocher color. The colonies soon coalesce giving culture a moist, mealy appearance.	Grows as a dark-brown, adhering membrane.		Small, opaque colonies which later coalesce, developing a tough, black membrane; as growth proceeds a bright yellow pigment is produced on the surface of the membrane; the medium also takes on a dark color.		Growth after 3 to 4 days at first as small, whitish colonies; these gradually take on a delicate pink color; later as the colonies develop they become umbilicated and coalesce forming a heaped-up growth and produce a burnt-ocher color.
ter 8 days as small, discrete, colonies with waxy, wrinkled ten developing pink pigment.	Similar to glycerine-agar		Similar to glycerine-agar.		Similar to glycerine-agar	Similar to glycerine-agar	Similar to glycerine-agar.		Similar to glycerine-agar		Similar to glycerine-agar.
	Growth appears after 3 days as small, pinkish-white colonies, which later develop a dark pink, round, granular growth.	Growth appears after 72 hours as a heaped-up, irregular mass of a creamy-yellowish-white color, often showing patches of pink coloration; no pigmentation or erosion of medium.	Similar to *Streptothrix maduræ* of Vincent.	Colonies in 4 to 8 days which form a dry, brown, raised growth.	Appear after 3 days as slightly raised colonies which soon assume a light-brown color.	Growth appears after 2 to 4 days as slightly raised, white colonies; these later assume a light brown color.	Grows as a dark-brown, adhering membrane.		Similar to glycerine-agar; medium becomes moist.		Do.
opaque-white, granular, more colonies at the bottom of the wth rare; no pigmentation of	Similar to literature	Similar to *Streptothrix maduræ* of Vincent.	do	Colonies develop on surface as concave, fine, dry, pale disks which latter form a surface membrane; fluid clear but a deposit forms in the bottom of the tube.	Colonies develop on the surface as fine, dry disks and later form a surface membrane; a deposit forms at the bottom of the tube; medium remains clear.	Growth occurs as a thin, white membrane on the surface of the medium; a deposit forms but the medium remains clear.	Growth from the planted material in the bottom of the tube; no surface growth.	Opaque, granular, like grape seeds, which coalesce to a loose, flaky sediment; no surface membrane forms.	Growth occurs in the bottom of the tube as a film containing many black granules; no growth on the surface of the medium.		Growth appears after 3 days as flat particles on the surface of the medium; these produce in time a delicate pink color; as growth proceeds a granular mass collects in the bottom of the tube; this mass is more or less coherent.
	Very slight growth; no pigment.		Very slight growth; no pigment.		Very slight growth; no pigment	Very slight growth; no pigment; no clubs.	Very slight growth; no pigment; no clubs.		Very slight growth; no pigment; no clubs.		Very slight growth; no pigment; no clubs.
ts, rabbits, guinea pigs, and ing more than a small nodule ction and these disappeared carried out intraperitoneal, ubcutaneous inoculations of ts, guinea pigs, fowls and ts were negative in every	Negative results in monkeys by intraperitoneal inoculation.	Similar to *Streptothrix maduræ* of Vincent.	Extensive lesions were produced by intraperitoneal injection in monkeys; lesions contain granules with organisms and are similar to those seen in madura foot.	Pathogenic for guinea pigs by subcutaneous and intraperitoneal injections.	Pathogenic for guinea pigs and monkeys by subcutaneous and intraperitoneal injections.	Intraperitoneal injections in monkeys produced numerous miliary tubercle-like bodies in the peritoneum and mesentery.	Negative		Intraperitoneal injection in monkeys produces extensive suppurative lesions containing many small, black granules.		Pathogenic for guinea pigs and monkeys by intraperitoneal injections.

and acute and general infection may take place during the course of a chronic localized type of the infection.

COMPLICATIONS.—While streptothricosis may be found together with a number of other diseases, or may develop its symptoms as sequelæ of other diseases, the principal direct complications are those due to mixed infections of bacteria. Several of the pathogenic bacteria have been found associated with *Streptothricæ* in lesions and the course and outcome of the disease is influenced accordingly.

Tuberculosis, because of its close clinical relation to streptothricosis and the frequent lung involvement of both organisms, has led to confusion in some of the reported cases.

TREATMENT.—The varieties of treatment most useful in this disease are general, as well as local surgical intervention if cause is indicated and a combination of all of these methods may be necessary. Potassium iodide administered in large doses over a long period of time is generally admitted to have a favorable influence on the course of the disease and complete cures have been reported from this method of treatment. Local measures consist in antiseptic dressings and the use of the Roentgen rays. Surgical measures consist in drainage, removal, or amputation according to the location and extent of the lesions. A combination of all three forms of treatment would suggest itself as offering the most favorable opportunity for recovery.

BIBLIOGRAPHY.[3]

* ABBOTT, A. C., and GILDERSLEEVE, N. The etiological significance of the acid-resisting group of bacteria, and the evidence in favour of their botanical relationship to bacillus tuberculosis. *Univ. Penna. Med. Bull.* (1902), **15**, 106–125.—On the actinomyces-like development of some of the acid-resisting bacilli (Streptothrices?). *Centralbl. f. Bakt. etc.*, (1902), Orig., **31**, 547–550, 1 pl.

ABBÉE, CONRAD. Drei Fälle von tödtlich verlaufener Actinomycose. *Beitr. z. path. Anat. u. z. allg. Path.*, (1897), **22**, 132–171.

ABEL. Demonstration von Sporozoen und pathogenen Hefen. *Münchener med. Wchnsch.*, (1897), **44**, 124.—Fettgehalt von Scolices und Coccidien. *Ibid*, 149–150.—Zur Bakteriologie der stomatitis und angina ulcerosa. *Centralbl. f. Bakt. etc.*, (1898), Abt. 1., **24**, 1–7.

* ACLAND, T. D. Actinomycosis hominis. *Trans. Path. Soc.* London, (1885–6), **37**, 545–549, 1 pl.—Actinomycosis hominis. *Brit. Med. Journ.*, (1886), **1**, 1159–60.—Etiology and pathology of actinomykosis. *Lancet*, (1886), **1**, 973–74.

† ACOSTA, E. Nueva propiedad del Cladothrix invulnerabilis. *Crónica médico-quirúrgica de la Habana*, (1894), no. 18; *Ref.: Centralbl. f. Bakt.* etc., (1895), Abt. 1. **17**, 465.—† and GRANDE ROSSI, F. Descripción de un nuevo cladothrix invulnerabilis. *Crónica médico-quirúrgica de la Habana*, (1893), no. 3; *Ref.: Centralbl. f. Bakt. etc.*, (1893), Abt. 1., **14**, 14–16.

[3] Unmarked references have been verified in the Library of the Bureau of Science; in references marked † in most cases the original is not available, but reviews and abstracts have been verified as cited; references marked * have not been verified. All symbols before names of authors refer to first title *only*.

Adami, J. G. Certain points in connexion with the development of our knowledge of actinomycosis and its causation. *Montreal Med. Journ.*, (1905), **34**, 88–93.—* and Kirkpatrick, R. C. A case of Madura foot disease (Mycetoma pedis, ochroid variety). *Trans. Assoc. Am. Phys.*, (1895), **10**, 92–100.—* Further observations upon Madura foot disease in America. *Montreal Med. Journ.*, (1895–6), **24**, 485–490.

* Äkerman, J. Actinomycosis hominis. *Hygiea*, (1891), **53**, 595–607.

* Afanassjeff, M. J. O klinicheskoi mikroskopii i bakteriologie aktinomikoza. St. Petersburg, (1888), 16 p.—† Ueber die klinische Mikroskopie und Bacteriologie der Actinomycosis. *St. Petersb. med. Wchnsch.*, (1888), n. s., **5**, 79; 83; *Also abstr.: Centralbl. f. Bakt. etc.*, (1888), Abt. 1., **4**, 207–208.—and Schultz, Fr. Ueber Actinomycosis. *Münchener med Wchnsch.*, (1889), **36**, 418.—Ueber die Aetiologie der Aktinomykosis (in der dritte Kongress der russischen Aerzte in St. Petersburg). *Centralbl. f. Bakt. etc.*, (1889), Abt. 1., **5**, 683–684; *also ref.: Jahresb. d. path. Mik.* (*Baumgarten*), (1889), **5**, 398.

* Agnew, D. H. Mycetoma of India. *In his:* Principles and practice of surgery, Phila., (1883), **3**, 489.

* Ahmed, Z. Mycetoma of the hand. *Indian Med. Gaz.*, (1882), **17**, 219; *Also: Trans. Calcutta Med. Soc.*, (1882), **3**, 103–105.

* Alamartine, H. Actinomycose sousmaxillaire. *Lyon méd.*, (1908), **110**, 140–142.

* Albert. Actinomykosefälle. *Wien. med. Presse*, (1889), No. 48.

* Albertini, Antonio Diaz, and Desvernine, Carlos M. Nota preliminar sobre dos casos de pié de Madura. *Rev. med. trop.* (1901), **2**, 73–87.—*See also* Desvernine, Albertini, Calnek, Debayle, and Moréno.

Alissow and Skworzow. *Ref.: Ergeb. d. allg. Path.*, (Lubarsch-Ostertag), (1898), **5**, 664–665.

* Ally, S. Case of fungus foot. *Indian Med. Gaz.*, (1876), **11**, 298.

Almquist, Ernst. Untersuchungen über einige Bacteriengattungen mit Mycelien. *Ztschr. f. Hyg. u. Infectionskrakh.*, (1890), **8**, 189–197, 1 pl.

* Ammentorp, L. Fyre Tilfälde af actinomycosis hominis. (Quatre cas d'actinomycose humaine.) *Nord, med Ark.*, (1888), **20**, no. 23, 1–19; no. 25, 4.—* Om Aktinomycose. Kliniske jagttagelser paa kommune hospitalet. *Bibliothek f. Laeger*, (1893), VII, **4**, 433–472; *ref.: Centralbl. f. Chir.*, (1894), 1074.—Zur Aetiologie der Actinomykose. *Wiener klin. Wchnsch.*, (1894), **7**, 514.

* Ancel and Thiry. Une observation d'actinomycose humaine avec étude bactériologique. *Rev. méd. de l'est*, (1898), **30**, 358–371.

* Anderson, A. R. A case of actinomycosis of the face and neck; operation; cure. *Med.-Chir. Trans.*, (1891–2), **75**, 103–107; *also abstr.: Proc. Roy. Med. & Chir. Soc.*, (1891–2), III., **4**, 15–20.

Angerstein, C. Kleine Mittheilungen aus der Praxis. *Berl. thierärzt. Wchnsch.*, (1901), 659–660.

* Anglin, W. G. A case of actinomycosis of the liver. *Kingston Med. Quart.*, (1901–2), **6**, 62–66.

* Antonelli, G. Sopra un caso di actinomicosi umana. *Policlin.*, (1907), **14**, 465–468.

Aoyama and Miyamoto. Ueber die menschenpathogene Streptothrix. *Mitt. a. d. med. Fac. d. kais. Univ.*, Tokyo, **4**, 231–276; *rev. in: Centralbl. f. Bakt. etc.*, (1901), Abt. 1., **29**, 262–263.

Archibald, E. W. Clinical bacteriology of actinomycosis. *Montreal Med. Journ.*, (1905), **34**, 103–105.

* Aribaud, G. Actinomycose du foie. Lyon, (1897).

Arnold, W. F. A non-fatal case of actinomycosis in man, with a reference to the prevalence of actinomycosis in northern China. *Southern Pract.*, (1896), **18**, 535–540, 1 pl.

* Arquembourg, and Colle, J. Un malade atteint de tumeur actinomycosique. *Écho méd. du Nord.* (1906), **10**, 405.—* Tumeur actinomycosique. *Nord. méd.*, (1906), **12**, 197.

* Artigalas. De l'actinomycose. *Rev. san. Bordeaux.* (1884–5), **2**, 75–77.

* Arustamoff. Sur Morphologie und Biologie der Leptothrix. *Vratch*, (1889).

Arwine, J. T. and Lamb, D. S. A fifth case of "fungous foot" in America. *Am. Journ. Med. Sci.*, (1899), n. s., **118**, 393–399.

* Von Arx, M. Leptothrixphlegmone, eine Phlegmone sui generis. *Correspondenzbl. f. Schweiz. Aerzte*, (1899), 161.

Aschoff, A. Ein Fall von primärer Lungenaktinomykose. *Berl. klin. Wchnsch.* (1895), **32**, 738–740; 765–768; 786–789.

Assmann, A. Generalisierte Aktinomykose. *Deutsche tierärzt. Wchnsch.*, (1904), **12**, 63–64.

* Atchinson. *Indian Ann. Med. Sci.*, (1861), **14**, 517.

* d'Audibert, Caille du Bourguet, and Legrain, E. Actinomycose de la face. *Ann. derm. et svph.*, (1893), III, **4**, 460.

† Audry, Ch. Sur les dégénérescences cellularies dans les parois de l'abcès actinomycosique sous-cutané. *Journ. mal. cut. et syph.*, (1896), **8**, 147–150; *transl. in: Monatsh. f. prak. Derm.*, (1896), **22**, 553–554.

*Aufrecht. Ein Fall von Aktinomykosis hominis. *In his:* Path. Mitt., Magdeburg, (1883), **2**, 50–53.

*Augier. Actinomycose de la région mammaire. *Nord méd.*, (1903), **9**, 154.

*Ausländer, M. I. Nĕskolko sluchayev aktinomikosa u shelovĕka. (Several cases of actinomycosis in man). *Yuzhno-russk. med. gaz.*, (1894), **3**, 509; 522.

*Auvray, M. Actinomycose du foie. *Rev. chir.*, (1903), **23**, pt. 2, 1–19.

*Axe, J. W. Observations and experiments tending to elucidate the pathology of a disease affecting the tongue of the ox. *Veterinarian*, (1877), **1**, 605; 759.

Babes, Victor. Ueber einige pathologisch-histologische Methoden und die durch dieselben erzielten Resultate. *Virch. Arch.*, (1886), **105**, 511–521, 1 pl.—Der Madurafuss (Aktinomyces des Fusses, Perical, Mycetom). *In:* Kolle und Wassermann's Handb. d. path. Mik., Jena, (1903), **3**, 454–470.—and Levaditi, C. Sur la forme actinomycosique du bacille de la tuberculose. *Arch. méd. exp.*, (1897), **9**, 1041–1048, 2 pl.—*Ibid. Compt. rend. Acad. sci.*, (1897), **124**, 791–793.

Bahr, Louis. Ueber Actinomycose beim Hunde. *Ztschr. f. Thiermed.*, (1904), **8**, n. s., 47–67, 1 pl.

Bail. Actinomykoseculturen. *Ref.: Deutsche med. Wchnsch.* (1899), **25**, Ver.-Beil., 263.

* Bainbridge, G. A case of mycetoma of the foot (pale variety) with implication of the lymphatics of the lower extremity, and, probably those of the abdomen. *Trans. Med. and Phys. Soc.* Calcutta, (1883), III, **2**, 32–38.

Bajardi, A. Die Streptothrix lingualis (Syn. Vibrio, Spirosoma linguale) im Munde der Gesunden und der Diptherischen. *Centralbl. f. Bakt. etc.*, (1904), Orig., **35**, 129–137, 1 pl.

* Balack, H. W. C. Ueber Lungenaktinomycose (im Auschluss an einen Fall Aktinomykose der Lungen und des Magens). *Inaug.-Dissert.*, Leipzig, (1893).

* Ballingal. *Trans. Med. and Phys. Soc.*, Calcutta, (1855), n. s. **2**, 273.

* Bang, B. Aktinomykosen eller Straalesvampsygdommen. (Actinomykosis or radiating fungoid disease.) *Tidsskr. f. Vet.* Kjobenh., (1883), II, **13**, 47–110.—* Aktinomykosen eller Straalesvampsygdommen. (Actinomykosis or radiating fungoid disease). *Hosp.-Tid.*, Kjobenh., (1883), III, **1**, 673; 697; 721.—Die Strahlenpilzerkrankung (Actinomykosis). *Deutsche Ztschr. f. Thiermed.*, (1884), **10**, 249–264.

* Baracz, Roman Von. Przenos'nos'c' promienicy z czlowieka na czlowieka. (Actinomycosis transferred from man to man.) *Przegl. lek.*, (1888), **27**, 201–203; *also transl.* (with additions): *Wien. med. Presse*, (1889), **30**, 6–11.—Ueber neun Fälle der menschlichen Actinomycose; ein Beitrag zur Lehre von der Actinomycose. *Wien. klin. Wchnsch.*, (1890), **3**, 497–499; 517–519; 540–542.—* Przypadek promienicy pluc i klatki piersiowéj. (Actinomycosis of lungs and thorax). *Przegl. lek.*, (1891), **30**, 313.—Ueber die Actinomykose der Menschen auf Grund einiger Beobachtungen. *Arch. f. klin. Chir.*, (1902), **68**, 1050–1161.—* Report of sixty cases of Actinomycosis. *Ann. Surgery*, Phila. (1903), **37**, 336–340; 463–468.

Baranski, A. Zur Färbung des Actinomyces. *Deutsche med. Wchnsch.*, (1887), **13**, 1065.—Ein Beitrag zum Vorkommen des Actinomyces beim Pferde. *Arch. f. wiss. u. prak. Thierhlknde.*, (1889), **15**, 242–247.

* Bargum, J. Ein Fall von Actinomykosis hominis unter dem Bilde einer acuten Infectionskrankheit verlaufend. *Inaug.-Diss.*, Kiel (1884).

* Barnard, C. E. Notes on actinomycosis, and its transmissibility to the human subject. *Papers & Proc. Roy. Soc. Tasmania*, (1890), 254–259.

Barth, Arthur. Ein Fall von geheilter Actinomycose des Unterliebs. *Berl. klin. Wchnsch.*, (1889), **26**, 122.—Ueber Bauchaktinomykose. *Deutsche med. Wchnsch.*, (1890), **16**, 742–745.

Barthelat, G. B.—Les mucorinées pathogenes et les mucormycoses chez les animaux et chez l'homme. *Arch. parasit.*, (1903), **7**, 5–116.

† Bary, A. de. Vergleichende Morphologie und Biologie der Pilze, Mycetozoen und Bakterien, Leipzig, (1884), 406–407; *also transl.* (1887), 377–379.

† Bassini, E. Un caso di micetoma el piede, o piede di Madura. *Arch. per le sci. med.*, (1888), **12**, 309–318, 3 pl.; *also abstr.: Gazz. osp.*, (1887), **8**, 331; *also abstr.: Centralbl. f. Bakt. etc.*, (1888), Abt. 1., **4**, 652–654.

* Batasheff, I. S., and Shtshegloff, M. A. Case of actinomycosis. *Vratch, Zapiski*, (1897), **4**, 191–199.

* Batut, L. De l'actinomycose dans l'armée a propos de quatre cas observés. *Journ. mal. cutan. et syph.*, (1902), VI, **14**, 648–657, 1 pl., 2 fig.

* Bauby. Actinomycose et botryomycose. *Arch. méd. Toulouse.*, (1902), **8**, 557–564.

* Bauer, A. K. Aktinomikoz u 11-liet. devochki. (Actinomycosis in a girl of 11 yrs). *Med. Obozr.*, (1891), **36**, 594–599.

Baum, Richard. Ein Beitrag zur Klinik der Aktinomykose. *Berl. klin. Wchnsch.*, (1908), **45**, 256–258.

Baumgarten, P. Der Actinomyces. *In his:* Lehrbuch der pathologischen Mykologie, Braunschweig, (1890), **2**, 869–894.

* Baylac, and Ferron, M. De l'actinomycose dans le département des landes. *Prov. méd.*, (1907), **20**, 250.

* Bayo, J. T. Madura foot in Cape Colony. *South African Med. Rec.*, (1905), **3**, 75.

* Beck, B. von. Ueber Aktinomykose des Verdauungsapparates. *Aerztliche Mitt. a. Baden*, (1901).

* Bécue, G. De l'actinomycose. *Thèse de Lille*, Paris, (1892).—* Un cas d'actinomycose chez l'homme. *Bull. Soc. anat.-clin. Lille*, (1892), **7**, 92–98; *also: Journ. sci. méd. Lille*, (1892), **2**, 12–18.

* Begrain. Sur un cas d'actinomycose de la face. *Ann. derm. et. syph.*, (1891), **2**, 772.—* Actinomycose du menton et du maxillaire infereur. *Ibid.*, (1895), **6**, 682.

Behala, Robert. Ueber die systematische Stellung des Erregers der Aktinomykose. *Centralbl. f. Bakt. etc.*, (1898), Abt. 1., **23**, 817–829.

Beijerinck, M. W. Ueber Chinonbildung durch Streptothrix chromogena und Lebensweise dieses Mikroben. *Centralbl. f. Bakt. etc.*, (1900), Abt. 2., **6**, 1–12.

Bell, C. E., and Harris, J. D. Two cases of actinomycosis. *Lancet*, (1904), **1**, 1656–1657.

Bell, James. Actinomycosis: Introduction and recital of cases. *Montreal Med. Journ.*, (1905), **34**, 81–88.

Bellisari, G. Sulla presenza e sulla patogenità di streptotricee nelle polveri, residui di cereali. *Ann. ig. sper.*, (1904), n. s., **14**, 467–483, 1 pl.

† Belski. Versuche mit Uebertragung der Actinomycose. *Arch. f. Veterinärm. rev.: Jahresb. d. gesammt. Med.*, (1891), **1**, 683.

* Benavente, D. Actinomicósis humana. *Rev. méd. Chile*, (1905), **33**, 321–325.

Benda, C. Die morphologische Bedeutung der Actinomyceskolben. *Deutsche med. Wchnsch.*, (1900), **26**, Ver-Beil., 132.—† Zwei Fälle von metastasirender Aktinomykose. *Verhandl. d. Ver. f. inn. Med. zu Berlin*, (1900), **19**, 485–490; *also: Münchener med. Wchnsch.*, (1900), 47, 372; *also: Deutsche med. Wchnsch.*, (1900), **26**, Ver.-Beil., 70–71.

* Bendig, Ernst. Ueber einen Fall primärer von Wurmfortsatz ausgehender Bauchaktinomykose. Leipzig, (1904), 34 pp.

* Bérard. De l'iodure de potassium dans l'actinomycose, Bordeaux, (1894).—* Actinomycose de la face. *Lyon méd.*, (1895), **78**, 261.—* De l'actinomycose humaine; sa frequence en France; nécessité et moyens de la reconnaître; données cliniques; diagnostic et traitement. *Gaz. hôp.*, (1896), **69**, 253; 289.—* Actinomycose cervico-faciale è forme chronique. *Lyon méd.*, (1898), **87**, 448.—* Actinomycose anale périrectale et pérrivéscale. *Bull. méd.*, (1898), **12**, 559.—* and Dor, L. Actinomycose expérimentale. *Mém. et Compt. rend. Soc. sci. méd.* Lyon, (1893), **33**, pt. 2, 41–43.—* and Nicolas, Joseph.—Sur l'étiologie de l'actinomycose. *Lyon méd.*, (1898), **87**, 22–24.—† Note sur la résistance des spores de l'actinomyces. *Compt. rend Soc. biol.*, (1900), XI, **2**, 835–836; *also: Prov. méd.*, (1900), **15**, 498–499; *ref.: Centralbl. f. Bakt. etc.*, (1901), Abt. 1., **29**, 66.—* and Thévenot, L. Á propos de l'actinomycose du foie. *Rev. chir.*, (1903), **23**, 397.

† Berestnew, N. Aktinomykose und ihre Erregers. *Inaug.-Dissert.*, Moskau, (1897), 206 pp., 5 pl.; *abstr.: Centralbl. f. Bakt, etc.*, (1898), Abt. 1. **24**, 706–708; *also: Virch. Arch.*, (1898), **152**, 399–400; *also: Ergeb. d. allg. Path.* (Lubarsch-Ostertag), (1898), **5**, 664–667.—Ueber Pseudoaktinomykose. *Ztschr. f. Hyg. u. Infectionskrankh.*, (1898), **29**, 94–116, 3 pl.—* Zur Aktinomykosefrage. *Prager med. Wchnsch.*, (1899), **24**, 619.—Zur Frage der Klassifikation und systematischen Stellung der Strahlenpilze. *Centralbl. f. Bakt. etc.*, (1899), Abt. 1., **26**, 390.

* Berg. Aktinomykose des Metatarsus bei einem Rind. *Maanedsskrift for Dyrlaeger*, **8**, 226.—* Aktinomykose bei Schafen. *Ibid.*, **10**, 9.

* Bergase. Sur un cas d'actinomycose. *Bull. soc. méd.-chir. Drôme etc.*, (1905), **6**, 114–116.

* von Bergmann. Forstellung von Fällen von Aktinomykose. *Tageblatt der 59. Versamml. deutscher Naturforscher u. Aerzte zu Berlin*, (1886), 113.

* Berkeley, M. J. On the peculiar mode of fructification in Chionyphe carteri. *Trans. Med. & Phys. Soc.*, Bombay, (1869), n. s., **9**, (app.), 34–37.—* On the so-called fungus foot disease of India. *Med. Press & Circ.*, (1876), Dec 6, 465.

DE BERNARDINIS, D. Ulcera corneale da streptothrix. *Ann. ig. sper.*, (1904), n. s., **14**, 461–466.

* BERNHARTH, A. Ueber Aktinomykose und Demonstration eines Fälle von Bauchaktinomykose. *Prager med. Wchnsch.*, (1895), **20**, 383–386.

* BERT, A. Un nouveau cas d'actinomycose temporo-faciale dans la region lyonnaise. *Prov. méd.*, (1895), **9**, 577–579.

BERTARELLI, E. Ueber einen ziemlich seltenen Tuberkelsputumbefund. *Centralbl. f. Bakt. etc.*, (1903), Abt. 1., **34**, 411–412.

† BERTHA, M. Ueber einige bemerkenswerthe Fälle von Aktinomykose. *Wien. med. Wchnsch*, (1888), **38**, 1181–1184; *also: Med. chir. Centralbl.*, (1889), **24**, 27–29.

* BETAGH, G. Studio sulla biologia del agente patogeno isolato in due casi di actinomicosi umana. *Arch. gén. méd.*, (1906), **1**, 595–628, 1 pl.

† BEVAN, ARTHUR DEAN. Treatment of actinomycosis and blastomycosis with copper salts. *Journ. Am. Med. Assoc.*, (1905), **45**, 1492–1493; *also: Ill. Med. Journ.*, (1905), **8**, 514–516.—* Actinomycosis. *Ann. Surgery*, (1905), **41**, 627–654; *also: Chic. Med. Rec.*, (1905), **27**, 697–708.

* BIAGI, NELLO. Contributo alla conoscenza del genere actinomyces. *Sperimentale*, (1904), **58**, 655–716.

† BIANCHI, A. Cited by Rivolta *in his*, "Sul cosi detto mal del rospo, etc.". *Clinica vet.; rev.: Jahresb. d. gesammt. Med.*, (1878), **1**, 619.—* L'actinomicosi o malattia del Rivolta. *Sperimentale*, (1883), **51**, 287–304.

* BIDIE, G. Notes on morbus pedis entophyticus. *Madras Quart. Journ. Med. Sci.*, (1862), **4**, 222–227, 1 pl.

* BĚLSKI, G. E. Opyty nad zarazheniem aktinomikozom. (Experiments on the contagion of actinomycosis). *Arch. vet. nauk.*, (1891), **1**, 117–128.

* BIGAIGNON, L. Un cas d'actinomycose. *Odontologie*, (1907), **37**, 360–362.

* BILLET, A. Contribution a l'étude de la morphologie et du developpement des bacteriacées. *Bull sci. France et Belgique*, (1890), 287, 9 pl.; Cladothrix dichotoma, 25–109, 4 pl.

* BILLINGS, F. S. A consideration of actinomycosis as to its nature and relation to the public health. *Journ. Comp. Med. & Vet. Arch.*, (1892), **13**, 269; 358; *also: Times & Reg.*, (1892), **24**, 484; 502.

* BILLROTH, T. Actinomycosis. *Allg. Wien. med. Ztg.*, (1886), **31**, 316; 328.—* Some remarks on actinomycosis. *Ala. Med. & Surg. Journ.*, (1887), **2**, 321–329.

*BIONDI, D. Due casi di actinomicosi umana. *Boll. clin.*, (1903), **20**, 116–123.

* BIRCH-HIRSCHFELD. Actinomykosis, Strahlenpilzkrankheit. *Real-Encycl. d. ges. Heilk.*, (1893–4), 3. ed., **1**, 220–242; *also transl.: Uchen. zapiski Kasan. Vet. Inst.*, (1893), **10**, 9–32.—* Allg. path. Anat. Leipzig, (1897), 391.

BISHOP, D. D. *See* Hyde, Senn, and Bishop.

* BIZZOZERO. L'actinomicosi una nouva malattia da parasiti vegetali. *Gazz. osp.*, (1882).

* BLACK, S. P. Actinomycosis. *South. Cal. Pract.*, (1898), **13**, 41–44.

* BLAIR, W. R. Actinomycosis in antelopes. *Am. Vet. Rev.*, (1903–4), **27**, 857.—Actinomycosis in a grizzly bear. *Ibid.*, (1905–6), **29**, 1344–1347.

* BLANCHARD, R. Parasites. *In:* Traité de pathologie générale de Bouchard, Paris, (1895), **2**, 649–932.—* Sur un mémoire de M. le Dr. E. Legrain intitulé: Notes sur un nouveau cas de pied de Madura observé en Algérie. *Bull. Acad. méd.*, (1896), III, **36**, 753–757.—Sur le champignon du mycétome a grains noirs. *Ibid.*, (1902), III., **48**, 57–60.—Sur un cas de mycétome d'origine aspergillaire observé en Tunisie par les Drs. Nicolle et Brunswic-Le Bihan. *Bull. Acad. méd.*, (1906), III., **55**, 132–135.

* Block. Zur Kenntniss der Pilzbildung in der geweben des thierische Organismus. *Inaug.-Dissert.*, (1870).

Blume, A. Die Vasogene in der thierärztlichen Praxis: Behandlung actinomycotischer Geschwülste bei Rindern mit Jodvasogen. *Berl. thierärzt. Wchnsch.*, (1901), 89.

* Boari, A. Un caso di actinomicosi umana primitiva del fegato. *Atti Accad. sci. med. e nat.* Ferrara, (1895–6), **70**, 247–256; *also: Policlin.*, (1897), **4–C.**, 19–25.

* Bocarro, J. E. Mycetoma. *Indian Med. Rec.*, (1893), **5**, 1; 74; 106; 148; 214.—An analysis of one hundred cases of mycetoma. *Lancet*, (1893), **2**, 797–798; *also rev.: Monatsh. f. prak. Derm.*, (1894), **19**, 461–462.—* Clinical notes on mycetoma; differentiating the pale from the black variety. *Indian Med.-Chir. Rev.*, (1894), **2**, 893–896; *also: Indian Med. Rec.*, (1894), **7**, 361; *also: Indian Med. Gaz.*, (1894), **29**, 451.—Mycetoma. *Lancet*, (1895), **1**, 70–71.—* Pied de Madura. *Arch. méd. nav.*, (1895), **64**, 149.—* Mycetoma. *Indian Lancet*, (1897), **10**, 318.

* Bodamer, G. A. The pathology of actinomycosis, with record of cases and experiments; and observations upon human actinomycosis. *Trans. Path. Soc.* Phila., 1887–9, (1891), **14**, 280–318; *also: Journ. Comp. Med. & Surg.*, (1889), **10**, 105–126; 195–209, 3 pl.—* Actinomycosis in man, with the report of a case. *Med. News*, (1889), **54**, 230–232.

* Boddaert, E., and Vercauteren, H. Note sur un cas d'actinomycose abdominale. *Ann. Soc. méd. Gand.*, (1903), **82**, 231–236.

Böhm, Ignaz. Primäre Actinomycosis cutis am Hinterhaupte. *Arch. f. Derm. u. Syph.*, (1902), **59**, 393–396.

* du Bois Saint-Severin. Note sur une streptothricée parasite (streptothrix aurea). *Arch. méd. nav.*, (1895), **63**, 252–262, 1 pl.—* Sur une conjonctivite streptothrique, curable par les applications de jus de citron. *Sem. med.*, (1895).

* Bollinger, O. Aktinomykose der Rachenschleimhaut in Form eines faustgrossen Tumors und fünf Fälle von Aktinomykose der Zunge beim Rind. *Jahresb. d. K. Centr.-Thierarznei-Schule in München*, (1876–7).—Ueber eine neue Pilzkrankheit beim Rinde. *Centralbl. f. med. Wiss.*, (1877), **15**, 481–485; *rev.: Jahresb. d. gesammt. Med.*, (1877), **12**, 602; *also: Ztschr. f. Thiermed.*, (1877), **3**, 334–340.—Ueber primäre Aktinomykose des Gehirns beim Menschen. *Münchener med. Wchnsch.*, (1887), **34**, 789–792.—* Ueber primäre Aktinomycose der Fusswurzelknochen. *Ibid.*, (1903), **50**, 2–4.

* Von Bonsdorff, H. Actinomycosis hominis. Ett bidrag till kännedomen om denna sjukdoms förekomst i Finland. (. . . Contribution to knowledge of the disease and its prevalence in Finland.) *Finska läk.-sällsk. handl.* Helsingfors, (1894), **36**, 253; 395.

*Bonvicini, A. Una nuova forma di micosi cutanea nei bovini determinata da un ifomicete appartenente al genere streptothrix-Cohn. *Nuovi Ercolani*, (1899), **4**.

* Borelius, J. Om aktinomykos i Blekinge. *Hygiea*, (1896), **58**, 156–164.—* Bauchaktinomykose in Form eines cirkumskripten, beweglichen Tumors. *Nord. med. Ark.*, (1903), III., **3**, 1–16, 1 pl.

* Borszéky, K. Idegen test által okozott aktinomykosis a nyakon. (Actinomycosis in the neck caused by foreign body). *Orvosi hetil.*, (1904), **48**, 708; *also transl.: Ungar med. Presse*, (1904), **9**, 603; *also: Pest. med. chir. Presse*, (1905), **41**, 334.

Boström. Demonstration mikroskopischer Präparate von Aktinomykose der Menschen und Thiere. *Berl. klin. Wchnch.*, (1885), **22**, 14; 274.—* Ueber Actinomycose; Demonstration von Culturen des Actinomyces. *Verhandl. d. Cong. f. innere Med.*, (1885), **4**, 94–99.—Untersuchungen über die Aktinomykose des Menschen. *Beitr. z. path. Anat. u. z. allg. Path.*, (1891), **9**, 1–240, 10 pl.—* and Herr, M. Ein seltener Fall von Aktinomykose bei einem sechsjährigen Kinde. *Prakt. Aerzt*, Wetzlar, (1891), **32**, 265–271.

* Bouffard. Pied de Madura observés a Djibouti. *Ann. hyg. et méd. colon.*, (1902), 636.—* Du mycétome a grains noirs en Afrique. *Ibid.*, (1905), **8**, 579–590, 1 pl.—*See also* Brumpt, Bouffard, and Chabaneix. *See also* Chabaneix and Bouffard.

* Bourquin, E., and de Quervain, F. Sur les complications cérébrales de l'actinomycose. *Rev. méd. Suisse*, Rome, (1897), **17**, 145–163.

Boyce, Rubert. Eine neue Streptothrixart, gefunden bei der weissen Varietät des Madurafusses. *Hyg. Rundschau*, (1894), **4**, 529–531; *also abstr.: Centralbl. f. Bakt. etc.*, (1894), Abt. 1., **16**, 918–919.—and Surveyor, F. Madura disease (Mycetoma carteri). *Lancet*, (1893), **1**, 846–847.—The fungus-foot disease of India. *Brit. Med. Journ.*, (1894), **2**, 638–639.—Upon the existence of more than one fungus in Madura disease (mycetoma). *Phil. Trans. Roy. Soc.*, London, (1894), **185 B**, 1–14, 4 pl.; *also abstr.: Proc. Roy. Soc.*, London, (1893), **53**, 110–112.—* *Trans. Path. Soc.*, London, (1894), April.

* Boyd, W. H. A case of melanoid mycetoma of the foot, in which the glands in the popliteal space and groin were enlarged; amputation; recovery. *Trans. Med. & Phys. Soc.* Bombay, (1883), III, **3**, 45–48.—* Notes on a case of ochroid mycetoma of the right foot, in which one gland in the popliteal space, and two beneath and one above Poupart's ligament were enlarged. *Ibid.*, 49–52.

† Braatz, E. Zur Aktinomykose; Zweigbacterien im Harn. *St. Petersb. med. Wchnsch.*, (1888), n. s., **5**, 119; 127; *ref.: Centralbl. f. Bakt. etc.*, (1888), Abt. 1., **4**, 208–209.

* Brabec, A. Sur l' actinomycome. *Cosop. lék. cesk.*, (1902), **41**, 839–843; 865–870.—* Ueber eine weniger bekannte Erscheinungsform der Aktinomykose beim Menschen. *Wien. klin. Rundschau*, (1902), **16**, 917; 937.

Bragadio, Alvise. Di un raro caso di localizzazione actinomicotica. *Giorn. Soc. ed Accad. vet. ital.*, (1908), **57**, 267–269.

Brault, J. Contribution à l'étude de l'actinomycose. Un cas d'actinomycosique constaté à Alger. Peritonite actinomycosique chez le lapin et le cobaye. *Arch. parasit*, (1899), **2**, 535–547.—Un cas d'actinomycose de la joue droite observé à Alger. *Compt. rend. Soc. biol.*, (1899), XI, 1, 17–18.—* Mycétome à forme néoplasique simulant un fibrosarcome de la face dorsale du pied. *Bull. et mém. Soc. chir.* Paris, (1906), n. s., **32**, 589–594, 2 pl.—* Étude anatomo-pathologique et bacteriologique d'une maladie dite de madura à forme néoplasique. *Arch. méd. exp.*, (1907), **19**, 145–151, 1 pl.

* Braun, H. Ueber Actinomykose beim Menschen. *Cor.-Bl. d. allg. ärztl. Ver. v. Thüringen*, (1887), **16**, 37–51.

* Brazzola, F. Sull' istogenesi delle lesioni anatomo-patologiche dell' actinomicosi. *Ercolani*, (1888), **1**, 65–78, 1 pl.

* Brett. *In:* Surgery of India. (1840).

* Briau, E. Actinomycose temporo-maxillaire gauche, à forme chronique, remontant à cinq ans, chez un homme de 25 ans . . . . *Lyon méd.*, (1897), **84**, 119–123.—* Travaux de la clinique du Professeur Poncet sur l'actinomycose. *Rev; Gaz. hebd. méd.*, (1897), n. s., **2**, 534–538.—* Actinomycose cutanée. *Lyon méd.*, (1898), **87**, 371.

* Brigidi, V. Actinomicosi dell' uomo. *Gazz. ôsp.*, (1892), **13**, 897–900.

* Bristowe. *Trans. Path. Soc.* London, (1871), **22**,—* The so-called fungus-foot of India. *Indian Med. Gaz.*, Calcutta, (1875), **10**.—* Fungus foot of India. *Trans. Path. Soc.* London, (1881), 320.—* and Acland. *Ibid.*, (1886).

Brocq-Rousseu and Pietttre. Sur les spores d'un streptothrix. *Compt. rend. Acad. sci.*, (1906), **142**, 1221–1223.

* Brown, E. L. A case of actinomycosis. *Chic. Med. Rec.*, (1894), **7**, 251.

* Bruas. Pied de Madura à grains noirs observé a Madàgascar. *Ann. hyg. et méd. colon.*, (1903), **6**, 602–605.

Brumpt, Émile. Notes et observations sur les maladies parasitaires (Mission de M. le Vte du Bourg de Bozas en Afrique Centrale), II.:—Mycétome à grains noirs. *Arch. parasit.*, (1902), **5**, 151–156; 460.—Notes et observations sur les maladies parasitaires (Mission de M. le Vte du Bourg de Bozas en Afrique Centrale), II.:—Mycétome à grains blanc. *Arch. parasit.*, (1902), **5**, 156–158.—Sur le mycétome à grains noirs, maladie produite par une mucédinée du genre Madurella n. g. *Compt. rend. Soc. biol.*, (1905), **58**, 997–999.—Les mycétomes. *Arch. parasit.*, (1906), **10**, 489–572, 10 pl.—* Les mycétomes. *Thése*, Paris, (1906), 94 pp., 10 pl.—Bouffard, and Chabaneix, J. A. Notes et observations sur les maladies parasitaires (Mission de M. le Vte du Bourg de Bozas an Afrique Centrale):—Note sur quelques cas de paludisme et sur un cas de mycétome observés à Djibouti. *Arch. parasit.*, (1901), **4**, 564–567.—*See also* Reynier and Brumpt.

* Brun. Encyclop. Léauté, (1893).—* Maladies des pays chauds, **2**, 177.

* Bruní, C. Streptothicose des voies urinaires; recherches experiméntales. *Ann. mal. des organes génito-urin.*, (1904), **22**, 1601–1622, 4 pl.—Ueber die Streptotricose der Harnwege. *Monatsb. f. Urol.*, (1905), **10**, 87–90.

* Brunner, F. Beiträge zur Kenntniss der Actinomycose in der Sœweiz. *Cor.-Bl. f. schweiz. Aerzte*, (1896), **26**, 369–380.

† Bruns, Hayo. Ein Beitrag zur Pleomorphie der Tuberkelbacillen. *Inaug.-Dissert.*, Strassburg, (1895); *also: Centralbl. f. Bakt. etc.*, (1895) Abt. 1., **17**, 817–826.—Zur Morphologie des Actinomyces. *Ibid.*, (1899), Abt. 1., **26**, 11–15.

von Bruns. Zentrale Aktinomykose des Unterkiefers. *Münchener med. Wchnsch.*, (1903), **50**, 236–237.

Brunswic-Le Bihan. *See* Blanchard. (Rap.).

Buchholtz, Hermann. Ueber menschenpathogene Streptothrix; ein Beitrag zur Aetiologie des acuten Lungenzerfalls. *Ztschr. f. Hyg. u. Infectionskrankh.*, (1897), **24**, 470–487.

* Buday, K. Aktinomykosis abdominalis egy esete. *Orvosi Hetilap.*, (1889).—* Néhány aktinomykosis-eseteröll. (Some cases of actinomycosis). *Gyógyászat*, (1904), **44**, 100; 120; 150.

* Buerger, L. Actinomycosis of the skin of the foot. *Am. Journ. Med. Sci.*, (1907), n. s., **134**, 702–709.

Büsgen, M. Culturversuche mit Cladothrix dichotoma. *Ber. deutschen bot. Ges.*, (1894), **12**, 147–152, 1 pl.

* Bütschli. Ueber den Bau der Bakterien und verwandter Organismen, Leipzig, (1890).

† Bujwid, Odo. Hodowle promienicy. (Die Kultur des Strahlenpilzes). *Gaz. Lekarsk.*, (1889), no. 52; *ref.: Centralbl. f. Bakt. etc.*, (1890), Abt. 1., **8**, 430.—Ueber die Reinkultur des Actinomyces. *Ibid.*, (1889), Abt. 1., **6**, 630–633.—* Sur la culture pure de l'actinomycose. *Cong. internant. hyg. et démog. Compt. rend. et mém.*, 1889, (1890), 607.

* Bukhman, A. Contemporary condition of the science of actinomycosis: (two cases of actinomycosis of the jaw). *Vratch*, (1898), **19**, 158–164.

* Bulhões, O., and Magalhães, P. S. Observação de um caso de actinomycose humana. *Brazil-med.*, (1888), **2**, 12–14; *also: Gaz. med. de Bahía*, (1887–8), III., **5**, 449–459.

* Bullitt, J. B. Report of a case of actinomycosis hominis of the lungs. *Ann. Surgery*, (1900), **31**, 600–608; *also abstr.: Louisville Month. Journ. Med. & Surg.*, (1900–1), **7**, 172.

* Burci, E. Contributo allo studio dell' actinomycosi umana. Roma, (1898), 14 pp.

Burckhardt, H. Kombination von Aktinomykose und Adenocarcinom des Dickdarms. *Centralbl. f. Bakt. etc.*, (1908), Abt. 1., Orig., **46**, 396–400.

* Burrell, H. L., and Bottomley, J. T. Three cases of actinomycosis. *Med. & Surg. Rep. Boston City Hosp.*, (1898), IX, 138–140.

* Busachi, T. L'attinomicosi dell'uomo con relative osservazioni personali. *Riv. clin.*, (1891), **30**, 441–464.

Buschke, A. Die Blastomykose. *Arch. f. Derm. u. Syph.*, (1903), **68**, 415–434.

* Butler, G. R. Pulmonary actinomycosis; recovery under the use of oil of eucalyptus. *Med. News*, (1898), **72**, 513–515.

* Butterfield, E. E. A case of pulmonary infection with an acid-fast actinomyces. *Journ. Infect. Dis.*, (1905), **2**, 421–430, 1 pl.

* Buzzi, F., and Galli-Valerio, B. Osservazioni sopra un caso di actinomicosi dell'uomo guarito col ioduro di potassio. *Riforma med.*, (1893), **9**, 351–353.

* Bychowski. Ueber Aktinomykose. *6. Kongress der Pirogoff'schen Ges. russischer. Aerzte.*

* Byron, J. M. A case of actinomycosis in man. *N. Y. Med. Journ.*, (1889), **50**, 716.

† Caglieri, G. E. Actinomycosis of the lung, pleura, and chest wall. *Pacific Rec. Med. & Surg.*, (1898–9), **13**, 149–153; *also: Journ. Am. Med., Assoc.*, (1898), **31**, 1173–1174.

* Calkins. *Ann. N. Y. Acad. Sci.*, (1898), **11**, 379; 400.

* Calmette, E. L'actinomycose. *Bull. méd.*, (1896), **10**, 543.

Calnek, Thomas N. *See* Desvernine, Albertini, Calnek, Debayle, and Moréno.

* Cameron, H. C. A grass (oat?) seed removed from an actinomycosis growth on the cheek. *Glasgow Med. Journ.*, (1905), **64**, 277.

† Caminiti, R. Sulla variabilità del potere cromogeno dei microbi, e di alcune condizioni che la determinano nella streptotrix da me isolata. *Giorn. int. sci. med.*, (1907), n. s., **29**, 357–359; *also abstr.: Centralbl. f. Bakt. etc.*, (1907), Abt. 1., Orig., **43**, 754–755.—Ueber eine neue Streptothrixspecies und die Streptothricheen im allgemeinen. Uebersetzt von Dr. Kurt Tautz, Berlin. *Ibid*, **44**, 193–208, 4 pl.

* Canali, L. La broncoactinomicosi nell'uomo. *Riv. clin. d. Bologna*, (1882).

* Capdepon, and Vitrac. Actinomycose cervico-faciale. *Soc. anat. et phys.* Bordeaux, (1896).

* de Capua, S. Un caso di actinomicosi. *Incurabili*, (1891), **6**, 417–426.

* Carl. Ein Fall von ausgebreiteter Aktinomykose beim Schweim. *Deutsche tierärzt. Wchnsch.*, (1898).

* Carless. A case of actinomycosis. *Brit. Journ. Derm.*, (1897), **9**, 474.

* Carpot. Quoted by Le Dantec.

Carr, Walter, Johnson, Raymond, and Power, D'Arcy. Two cases of actinomycosis of the skin in children. (With case notes by A. H. Curtis.). *Lancet*, (1904), **2**, 1216–1217.

* Carroll, F. F. A case of actinomycosis. *Journ. Roy. Army Med. Corps*, (1905), **4**, 655.

* Cart, Edouard. Contribution à le étude de l'actinomycose chez l'homme. Paris, (1890), 83 pp.—De l'actinomycose. *Arch. gén. méd.*, (1894), **1**, 342–353.—* Actinomycose primitive de la peau. *Lyon méd.*, (1898), **88**, 152–154.

* Cartais, Adrian. Contribution a l'étude de l'actinomycose des conduits lacrymaux. *Thèse*, Bordeaux, (1904), 45 pp.; *also abstr: Poitou méd.*, (1904), **8**, 109–113.

* Carter, Henry Vandyke. On a new and striking form of fungus disease, principally affecting the foot, and prevailing endemically in many parts of India. *Trans. Med. & Phys. Soc.*, Bombay, (1860), n. s., **6**, 104–141, 3 pl.—* The fungus disease of India. *Trans. Path. Soc.* London, (1861), **24**, 169.—* Notes on the possibility of the embryos of the guinea-worm and so-called fungus-disease of India, respectively, entering the human body through the sudorific ducts. *Trans. Med. & Phys. Soc.*, Bombay, (1862), n. s., **7** (app.), 3–7.—* On mycetoma, or the fungus-disease of India; including notes of recent cases and new observations on the structure, etc., of the entophytic growth. *Ibid.*, 206–221, 2 pl.—* Memoranda accompanying a specimen of mycetoma. *Trans. Path. Soc.* London, (1863–4), **15**, 251–253, 1 pl.—* Note on mycetoma, the fungus disease of India. *Trans. Med. & Phys. Soc.*, Bombay, (1863), n. s., **8**, (app.), 26–28.—* On mycetoma. *Brit. & For. Med.-Chir. Rev.*, (1863), **32**, 198–203.—* Brief notes of two cases of mycetoma, or the fungus disease of India, with sketches. *Trans. Med. & Phys. Soc.*, Bombay, (1869), n. s., **9**, (app.), 47–49.—* The parasitic fungus of mycetoma, or the fungus of India. *Trans. Path. Soc.* London, (1872–3), **24**, 260–263.—* The etiology of Madura-foot. *Indian Med. Gaz.*, Calcutta, (1874), **9**, 220.—† On mycetoma, or the fungus disease of India, London, (1874), 113 pp., 11 pl; rev.: *Lancet*, (1874), **2**, 591.—† On the nature of mycetoma, or the fungus disease of India. *Lancet*, (1874), **2**, 44; 113; *also* (*translation*): *Arch. méd. nav.*, (1875), **24**, 157–170.—* The so-called fungus-foot of India. *Indian Med. Gaz.*, (1875), **10**.—† Note on the apparent similarity between mycetoma and actinomyces. *Trans. Med. & Phys. Soc.*, Bombay, (1886), III, **9**, 86–90; *also abstr: Lancet*, (1886), **1**, 321.—* *Trans. Path. Soc.* London, (1886).

* Casagrandi. Sulle relazioni tra batteri proto, meta, paratrofi. *Questi Ann.*, (1901–3).

* Catterina, A. Contributo alla cura dell'actinomicosi con alte dosi di joduro di potassio. *Clin. chir.*, (1898), **6**, 49–56.

* Cazalis, C. A. Streptothrix Foersteri symbiosé avec un micrococcus recontré dans la secrétion de la conjonctivite granuleuse pseudotuberculose expérimentale. *Nouveau Montpel. méd.*, (1896), **5**, 309; 331; 351, 1 pl.

* Ceci, A. Mucormicosi in mano affeta da osteocondroma (mano di Madura). Contribuzione allo studio delle infezioni ifomicetiche, Genova, (1887).

* Chabaneix, J. A., and Bouffard. Pied de Madura, observés à Djibouti. *Ann. hyg. et méd. colon.*, (1901), 452.—*See also* Brumpt, Bouffard, and Chabaneix.

* Charmoy. Trois cas d'actinomycose. Réflexions à leur sujet. *Bull. Soc. méd.* Yonne, (1901), **42**, 63–74.

Chiari, H. Ueber primäre Darmactinomycose des Menschen. *Prager med. Wchnsch.*, (1884), **9**, 93–94.

* Chiari, O. De la "pharyngomycosis leptotricca." *Rev. mens. de laryngol. etc.*, (1887).

* Chiarolanza, R. Ricerche sperimentali sulla emolisina di una streptotrix. *Med. ital.*, (1907), **5**, 425.

Childe, F. L. A remarkable case of mycetoma. *Lancet*, (1894), **2**, 1271–1273.—*See* Hatch and Childe.

Chipman, W. W. The clinical aspect of actinomycosis. *Montreal Med. Journ.*, (1905), **34**, 93–98.

* Choux. Un cas d'actinomycose. *Arch. méd. et pharm. mil.*, (1891), **18**, 490–494.—Étude clinique et thérapeutique de l'actinomycose. *Arch. gen. méd.*, (1895), **1**, 401–421; 565–588; 664–697.

* Chrètien, E. De l'actinomycose humaine. *Sem. méd.*, (1895), **15**, 17–24; *also transl.: Med. Week*, Paris, (1895), **3**, 37; 49.—* Des manifestations buccofaciales de l'actinomycose chez l'homme. *Odontologie*, (1895), II., **2**, 65–74.

* Cienkowski, L. Zur Morphologie der Bacterien. *Mém. Acad. imp. sci.* St. Petersbourg, (1877), VII., **25**, 18 pp., 2 pl.

Ciechanowski, Stanislaw. Zur Aktinomycesfärbung in Schnitten; technische Notiz. *Centralbl. f. Bakt. etc.*, (1903), Abt. 1., Orig., **33**, 238–239.

* Claisse. De l'actinomycose linguale primitive. *Presse méd.*, (1897); *ref: Centralbl. f. Chir.*, (1898), no. 10.

Claus, Eduard. Ueber die Localisation und geographische Verbreitung der Aktinomykose beim Rind in Bayern. *Deutsche Ztschr. f. Thiermed.*, (1888), **13**, 290–300.

* Claussen. Aktinomykose der Zunge beim Rind. *Mitt. f. Tierärzte*, **3**.

* Cleghorn, J. Case of Madura foot. *Indian Med. Gaz.*, Calcutta, (1874), **9**, 260.

Clemow, Frank G. Mycetoma (Madura foot) in the Yemen. *Brit. Med. Journ.*, (1906), **1**, 918–919.

* Clerc. Actinomycose de l'amygdale. *Thèse*, Lyon, (1905).

* Cleveland, R. A. A case of mycetoma in Cyprus. *Journ. Trop. Med.*, (1907), **10**, 223.

* Codman, E. A. A case of actinomycosis. *Boston Med. & Surg. Journ.*, (1898), **139**, 134.

* Cohn, Ferd. Biologische Mitteilungen über Bakterien. *Jahresb. d. schles. Ges. f. vaterl. Kult.*, (1874), **51**, 116–119.—* Untersuchungen über Bacterien, II. *Beitr. z. Biol. d. Pflanz.*, (1875), **1**, 141–208, 2 pl.

* Goignet. Actinomycose du maxillaire supérieur. *Mém. et compt.-rend. Soc. sci. méd.* Lyon, (1893), **33**, 94–96.

* Colebrook. Indian Army Med. Rep., (1844).—* Med. Rep. by Med. Board, Madras, (1850).—*See also* Godfrey and Colebrook.

* Collas. Leçon sur la dégénération endémique des os du pied, Pondichery, (1861); *also: Arch. méd. nav.*, (1883), 81, 204.

Colle, J. *See* Arquembourg and Colle.

* Cominacini. Un cas di actinomicosi umana. *Gazz. osp.*, (1893), **14**, 922.

* Conti, P. L'actinomicosi bronco-polmonare primitiva nell'uomo. *Riv. ven. sci. med.*, (1885), **3**, 105; 231, 1 pl.

Cooper, Ludford. Actinomycosis of the tongue. *Trans. Path. Soc.* London, (1899), **50**, 61–62.

Coppen Jones, A. Ueber die Morphologie und systematische Stellung des Tuberkelpilzes und über die Kolbenbildung bei Aktinomykose und Tuberkulose. *Centralbl. f. Bakt. etc.*, (1895), Abt. 1., **17**, 1–16; 70–76, 1 pl.—Ueber die Nomenklatur des sogen. Tuberkelbacillus. *Ibid.*, (1896), Abt. 1., **20**, 393–395.

* Coppez, H., and Depage, A. Un cas d'actinomycose orbitaire. *Journ. méd.* Bruxelles, (1903), **8**, 753–756.

† Coquerel, Ch. Note sur l'examen microscopique des lésions que l'on observe dans l'affection connue sous les noms de pérical, pied de Madura. *Compt. rend. Soc. biol.*, (1866), IV, **2**, 191–196; *also: Gaz. méd.* Paris, (1896), **21**, 504.

* Corda. Prachtiflora enropäischer Schimmelbildungen, Leipzig u. Dresden, (1839), 23.—* Anleitung zum Studium der Mykologie, nebst kritischer Beschreibung aller bekannten Gattungen und einer kurzen Geschichte der Systematik. (1842), 43.

* Cornil. Actinomycose. *Journ. conn. méd. prat.*, (1886), III., **8**, 289; 397.—* and Babes. Les Bacteries. 3. ed., (1890).

Cornwall, J. W. Notes on the cultivation of Streptothrix madurae. *Indian Med. Gaz.*, Calcutta, (1904), **39**, 208–209.

* Corre, A. La maladie de Ballingal (pied du Maduré) d'après des notes inédites du docteur Collas, mises en ordre, completées et annotées par A. Corre. *Arch. méd. nav.*, (1883), **39**, 204–224, 1 pl.—* Traité clinique des maladies des pays chauds, (1887), 535.

* Cortes, Antonio de. Due casi di actinomicosi umana. *Gazz. osp.*, (1902), **23**, 1272–1275.

* Costantin, J. Les Mucédinées simples, Paris, (1888), 210 pp.

* Cotte, G. Actinomycose cervicale. *Lyon méd.*, (1905), **104**, 107.

* Cousin, Gabriel. Traitement de l'actinomycose par la levure de bière et l'arsenic associés (méthode du docteur Martel). *Thèse*, Lyon, (1904), 76 pp.

Cowie, D. Murray. A preliminary report on acid-resisting bacilli, with special reference to their occurrence in the lower animals. *Journ. Exp. Med.*, (1901), **5**, 205–214.

Cozzolino, Vincenzo. Ein neues Fadenbacterium, eine pseudo-aktinomykotische Erkrankung erzeugend. *Ztschr. f. Hyg. u. Infectionskrankh.*, (1900), **33**, 36–52, 1 pl.

* Craig, J. F. Disease in a heifer resembling actinomycosis. *Vet. Journ.*, (1906), **62**, 707.

* Cranwell, D. J. La actinomicosis humana en la República Argentina. *Rev. Soc. med. argent.*, (1903), **11**, 318–342; *also: Semana med.*, (1903), **10**, 454–456.—* Sobre dos nuevos casos de actinomycosis humana. *Rev. Soc. med. argent.*, (1903), **11**, 691–708; *also abstr.: Semana med.*, (1903), **10**, 1140.—* Sobre la frecuencia de la actinomicosis humana en la República Argentina. *Rev. Soc. med. argent.*, (1903), **11**, 789–792.—* Contribución al estudio de la actinomicosis humana; su frecuencia en la República Argentina. Buenos Aires, (1904), 193 pp., 10 pl.—* La actinomicosis en la República Argentina. *Rev. Soc. med. argent.*, (1904), **12**, 5–44; *also: Semana med.*, (1904), **11**, 725.—* Nueva contribución al estudio de la actinomicosis humana en la República Argentina. *Rev. Soc. med. argent.*, (1906), **14**, 316–321.

Crawford, D. G. Mycetoma, or fungus foot. *Indian Med. Gaz.*, (1903), **38**, 171–172.—† Two cases of actinomycosis. *Lancet*, (1905), **2**, 157–8; *also: Liverpool Med.-Chir. Journ.* (1905), **25**, 125–128, 2 pl.

* Crookshank, E. M. Actinomycosis hominis. *Med.-Chir. Trans.*, (1889), **72**.—* Actinomycosis. *Trans. 7 Int. Cong. Hyg. & Demog.*, (1891), London, **3**, 192–196.—* The morphology of actinomyces. *Ibid.*, **2**, 105.—Madura foot (Discussion). *Lancet*, (1893), **1**, 933.—Actinomycosis and Madura disease. *Ibid.*, (1897), **1**, 11–18, 1 pl.

* Csokor. Die Strahlenpilz-Erkrankung, Actinomykosis. *Allg. Wien. med. Ztg.*, (1881), **26**, 430.

Cuff, Archibald. A case of actinomycosis of the abdominal wall produced by an infected foreign body which had escaped from the intestine. *Brit. Med. Journ.*, (1906), **2**, 137–138.

Cunningham, D. D. Is mycetoma primarily owing to the action of the fungal elements ordinarily associated with its products? *Sci. Mem. Med. Off. India*, (1895), **9**, 67–74, 2 pl.—*See also* Lewis and Cunningham.

* Curtze, R. Die Aktinomykose und ihre Bekämfung. *Deutsche Med.-Ztg*, (1889), **10**, 569; 581; 593.

* Cusack, S. A. Fungus hematodes of the hand. *Proc. Path. Soc.* Dublin, (1852–8), 295–297.

* Cyze, J. Vysledky pri léceni aktinomykoosy. (Results of the treatment of actinomycosis). *Casop. lék. Cesk.*, (1904), **43**, 141; 171.

Czaplewski. Ueber einen aus einem Leprafalle gezüchteten alkohol- und säurefesten Bacillus aus der Tuberkelbacillengruppe. *Centralbl. f. Bakt. etc.*, (1898), Abt. 1., **23**, 97–107; 189–194, 2 pl.

Dalchow. Actinomykose. *Arch. f. wiss. u. prak. Thierhlknde.*, (1900), **26**, 362–363.

Damman, G. W. On the genera of streptothrix and cladothrix of Cohn. *Lancet*, London, (1893), **1**, 356–357.

* Dandois. Contribution au diagnostic et au traitement de l'actinomycose. *Ann. Soc. belge de chir.*, (1896–7), **4**, 297–302; also: *Rev. méd.*, Louvain, (1896–7), **14**, 289–294.

* Danlos. Un cas d'actinomycose faciale. *Bull et mem. Soc. méd. d. hôp. de Par.*, (1904), III, **21**, 466.

* Le Dantec. Étude bactériologique sur le pied de Madura de Sénégal (variété truffoïde). *Arch. d. méd. nav.*, (1894), **62**, 447–454.—* Précis de pathologie exotique, Paris. (1900), 662.

* Darier, J., and Gautier, G. Un cas d'actinomycose de la face. *Bull. Soc. franç. de dermat. et syph.*, (1891), **2**, 278–284; *Also: Ann. de dermat. et syph.*, (1891), III, **2**, 449–455, 1 pl.

* Darling, C. G. A case of actinomycosis. *Physician & Surg.* (1903), **25**, 315–318.

* Davids. Ueber die sog. Actinomycosis musculorum suis. *Inaug. diss.*, Giessen, (1898).

* Dawson, W. Actinomycosis of abdomen. *Vet. Rec.*, (1902–3), **15**, 595.

Dean, George. On a new pathogenic streptothrix. *Trans. Jenner Inst. Prev. Med.* (1899), **2**, 17–45, 3 pl. On a new pathogenic streptothrix. *Trans. Path. Soc.*, London, (1900), **51**, 26–47.

Debayle. *See* Desvernine, Albertini, Calnek, Debayle, and Moréno.

* Delbanco, Ernst. Ein amerikanischer Fall von Mycetoma pedis. *Deutsche Med-Zeitung*, (1897), **18**, 497–500; also: *Arb. a. Unna's Klin. f. Hautkr. in Hamb.*, (1898), 18–28; also *Festchrift f. Neumann* (1900), 117.—Eine neue Strahlenpilzart nebst Bemerkungen über Verfettung und hyaline Degeneration *München. med. Wchnsch.*, (1898), **45**, 48–50; 82–85.—*See also* Unna and Delbanco.

* Delearde. Contribution a l'étude de l'actinomycose. *Thèse.* Lille, (1895).—* Contribution à l'étude de l'actinomycose. *Rev. biol. du nord de la France*, (1894–5), **7**, 364; 401, 3 pl.—* Contribution à l'étude de l'actinomycose. *Bull. méd. du nord*, (1895), **34**, 409–434, 2 pl.

* Delépine, S. A case of actinomycosis hominis. *Trans. Path. Soc.*, London, (1888–9), **40**, 408–443, 3 pl.

* Delore, X. Actinomycose cerébrospinale. Méningite suppurée. *Gaz. hebd. de méd. et chir.*, (1896), **42**, I. 498.—* Actinomycose anale, périrectale et périvésicale. *Lyon méd.*, (1898), **68**, 372–374.

* Dematteis. Di alcuni microrganismi rivenuti nell' intestino. *Giorn. d. R. Accad. di Torino*, (1896).—* Il leptothrix nella enterite cronica e nell' anemia perniciosa progressiva. *Gazz. degli ospedali*, (1899), **37**, 392.

* Démias, Silvain. De l'actinomycose animale en particulier dans la région lyonnaise. Lyon, (1897), 45 pp.

* Dentler, M. Ueber Aktinomykose der Bauchdecken. *Monatschr. f. Geburtsh. u. Gynaek.*, (1903), **17**, 1236–1238.

Le Dentu. Remarques relatives au diagnostic de l'actinomycose. *Bull. Acad. méd.*, (1904), III, **51**, 111–115; also: 259–260.

* Depage. Un cas d' actinomycose. *Bull. Soc. roy. sc. méd. et nat. de Brux.*, (1903), **61**, 135–137.

† Desvernine, C. M., Albertini, Antonio Diaz, Calnek, Debayle, and Moreno. Madura foot in Cuba, Costa Rica and Nicaragua. *Pan-American Congr. Med. Record*, (1901), February 25. Also: (Note) *J. Trop. Med.*, (1901), **4**, 106.—*See also* Albertini and Desvernine.

* Didsbury, G. Sur un cas d'actinomycose pharyngée. *Rev. de laryngol.*, (1895),

* Dieulafoy. L'actinomycose thoracique. *Rev. gén. de clin. et de thérap.*, (1906), **20**, 499.

Doepke, Karl. Beitrag zur Kenntniss der Erregers d. mensch. Aktinomycose. *Münch. med. Wchnsch.*, (1902), **49**, I, 873–875.—Weitere Mitteilungen über den Erreger der menschlichen Aktinomykose. *München. med. Wchnsch.*, (1903), **50**, II, 2245–2250.

Domec, Th. Contribution à l'étude de la morphologie de l'actinomycose. *Arch. méd. exp. et d'anat. path.*, (1892), **4**, 104–113; Ref: *Baumgarten's Jahresb.*, (1892), **8**, 387.

* Donalies, Carl. Die Aktinomykose des Menschen. Halle a. S., (1894), 40 pp.

* Donius, René. Psoïtis actinomycosique. *Thèse.* Lyon, 1905, 50 pp.

Di Donna, A. Su di una streptothrix patogena con esperimenti sull'immunizzazione. *Ann. d'ig. sper.*, (1904), n. s., **14**, 449–459. 1 pl.

* Dor, L. Procédé simple pour rechercher les grains jaunes de l' actinomycose. *Lyon méd.*, (1904), **102**, 1231.—* Une nouvelle mycose à grains jaunes; ses rapports avec l'actinomycose; est-ce une variété de l'actinomycose ou une espèce différente? *Gaz. hebd. de méd. et chir.*, (1896) **43**, 553.— Une observation d'actinomycose de la joue et du maxillaire inférieur droit, avec propagation ou poumon droit. *Gaz. hebd. de méd. et de chir.*, (1893), II, **30**, 40–42; Ref: *Baumgarten's Jahresb.*, (1893), **9**, 433.—* Un cas d'actinomycose. *Mercredi méd.*, (1895), 586.—Actinomycose; recherche du champignon rayonné; ses caractères morphologiques et réactionnels dans les tissus. *Presse méd.* (1903), **11**, 654–658.

Dorn, C. Operation von Neubildungen actinomycotischer Natur. *Berl. tierärzt. Wchnsch.*, (1901), 492–493.—* Aktinomykotische Neubildung im Nasenrachen. *Wchnsch. f. Tierh. u. Viehzucht*, (1903), **47**, 247.—* Operative Behandlung von aktinomykotischen Neubildingen des Rindes. *Wchnsch. f. Tierh. u. Viehzucht*, (1907), **51**, 321; 341; 361.

* Downie, K. M. Case of Madura foot; mycetoma. *Indian Med. Gaz.*, (1873), **8**, 321.—* Madura foot disease; mycetoma of India. *Med. Press and Circ.*, (1874), 28.

* Doyen. *Ref.* Poncet and Bérard. Traité clinique de l'actinomycose humaine, pseudo-actinomycose et botryomycose. Paris, 1898.—* and Roussel. Trois cas d'actinomycose chez l'homme. *Trans. 7. Internat. Cong. Hyg. & Demog.*, (1891), London, **3**, 197–203.—* Deux cas d'actinomycose chez l'homme; opération; guérison. *Cong. franç. de chir. Proc.-Verb., etc.*, (1891), **5**, 463–465.

Dreyfus, Wilhelm. Ein Fall primärer Hautaktinomykose. *München. med. Wchnsch.*, (1903), **50**, II, 2291–2292.

Dubler, A. Ein Fall von Leptothrixmycose des pharynx, larynx und oesophagus. *Virchows Arch.*, (1891), **126**, 454–455.

* Dubreuilh. Sur un cas d'actinomycose. *Soc. d'anat. et de phys. de Bordeaux,* (1900).—* Actinomycose cervico-faciale. *Journ. méd. de Bordeaux,* (1906), **36,** 764; also: *Soc. d'anat. et de phys. de Bordeaux,* (1906).—* and Frèche, D. L'actinomycose dans le sud-ouest de la France. *Ann. de dermat. et syph.,* (1895), III, **6,** 788–803.—* and Sabrazès. Observation d'actinomycose humaine. *Soc. d'anat. et de phys. de Bordeux.* (1893).

* Duckworth, Sir D. A case of acute actinomycosis. *Trans. Clin. Soc.,* London, (1901), **34,** 1.—* A case of actinomycosis of the thoracic wall. *St. Barth. Hosp. Rep.,* London, (1895), **31,** 23–29.

*Ducor. Un cas d'actinomycose de la face. *Méd. mod.,* (1896).—* Actinomycose circonscrite datant de neuf ans. *Gaz. hop.,* (1896).—* Contribution à létude de l'actinomycose en France; actinomycose du maxillaire à forme néplasique limitée datant de huit ans. *Gaz. hòp.,* (1896), **69,** 966; 1015; 1030, 1071.—* Contribution à l'étude de l'actinomycose en France: actinomycose circonscrit du maxillaire inférieur. *Journ. méd. Paris,* (1896), II, **8,** 573–576.—* Contribution à l'étude de l'actinomycose en France. *Bull. Acad. méd.* Paris, (1897), III, **38,** 494–496.

* Düms. Ueber Aktinomykose in der Armee. *Deutsche mil.-ärztl. Ztschr.,* (1894), **23,** 145–162.

* v. Düring. Madurafuss *In:* Eulenberg's Realencyklopädie d. ges. Heilkunde, 3. ed., **14,** 203.

* Dufils. Actinomycose primitive des centres nerveux. *Thèse,* Lyon. 1905.

* Duguet. Sur un cas d'actinomycose bucco-faciale guéri. *Bull. Acad. méd.,* (1896), III, **31,** 36–41.—* Sur un cas de guérison d'actinomycose. *Sem. méd.,* (1896).—*Un cas d'actinomycose bucco-faciale guérie. *Presse méd.,* (1897), 213.

* Duma. Un cas d'actinomycose. *Bull et mém. Soc. chir.* Bucarest, (1900), **3,** 111.

* Dumont, G. Actinomycose à marche rapide. *Bull. méd. du nord,* (1896), **35,** 785–792.

* Duncker. Ein neues Färbungsmittel für Aktinomykose bovis. *Ztschr. f. Fleisch-u. Milchhyg.,* (1891), **1,** 56.—* Strahlenpilze (Actinomyces) im Schweinfleisch. *Ztschr. f. Mikr. u. Fleischschau,* (1884), **3,** 17–19; *Also transl.: Vet. Journ. & Ann. Comp. Path.,* (1884), **19,** 1–5.

* Duval. Quoted by Le Dantec.

* Dymock, D. F. Fungus disease; recurrence after amputation. *Indian Med. Gaz.,* (1881), **16,** 56.

* Ebermann. Material upon the bacteriology of suppuration (Russian). *Inaug. Dissert.* St. Petersburg. (1893).

Eddowes, Alfred. A case of actinomycosis primarily and solely affecting the cheek. *Brit. Journ. Dermat.,* (1906), **18,** 151–152.

* Eddowes, W. On the fungus foot disease and its treatment. *Indian Med. Gaz.,* (1867), **2,** 218.

* Ehrhardt. Zungenaktinomykose beim Pferde. *Schweiz-Arch. f. Tierhlk.,* (1896), **37.**—* Aktinomykose. *Schweiz-Arch. f. Tierhlk.,* (1897), **38.**

* Ekehorn, G. Om Aktinomykosi Blekinge. *Hygiea,* Stockholm, (1896), **58,** 422–424.

* Ekkert, N. Ob aktinomikoze u. krupnavo rogativo skota. (Aktinomycosis in horned cattle). *Vestnik obsh. vet.,* St. Petersb., (1889), **1,** 38–41.

* Eliasson, A. Om aktinomykos hos menniskan. (. . . in man). *Hygiea,* Stockholm, (1896), **58,** 337–358.—* Två fall af actinomykos, framgangsrikt behandlade med jodkalium. (Two cases of actinomycosis treated succesfully by iodide of potassium.) *Eira,* Stockholm, (1897), **21,** 262–264.

* Elschnig. Actinomyces im Thränenröhrchen. *Klin. Monatsbl. f. Augenh.*, **33.**

† Engel, A. Die Jodkaliumbehandlung der Actinomykose. *Veterinarius*, (1893), *Ref: Jahresb. d. gesammt. Med.*, (1893), **28**, I, 651.

Engelhardt, G., and Löhlein, M. Zur Kenntniss der Streptothrixpyämie. *Deutsches Arch. f. klin Med.*, (1903), **75**, 112–131.

* Engelmann, G. Ueber Actinomycose beim Menschen. *St. Petersb. me. Wchsch.*, (1893), n. s., **10**, 445–449.

Eppinger, Hans. Die durch Cladothricheen (Streptothricheen, etc.) hervorgerufenen Erkrankungen. *Ergeb. d. all. Path.*, (1896), **1**, I. 872–891; also: (1896), **3**, I. 328–364.—Ueber eine neue pathogene Cladothrix und eine durch sie hervorgerufene Pseudotuberculosis (cladothrichica). *Beitr. z. path. Anat. u. z. all Path.*, (1891), **9**, 287–328; also: *Wien. klin. Wchnsch.*, (1890), **3**, 321–323.

* Epstein, A. Ueber Angina chronica leptothricca bei Kindern. *Prag. med. Wchnsch.*, (1900).

† Ercolani (1865). Cited by Rivolta in his "Sul cosi detto mal del rospo, etc., *Clinica vet.; rev: Jahresb. d. gesammt. Med.*, (1878), **13**, I, 619.

* Ernst, Wilhelm. Ueber Nekrosen und den Nekrosebacillus (Streptothrix necrophora). *Monatsh. f. prak. Tierhlk.*, (1902), **14.**

Esmarch. Ref: *München. med. Wchnsch.*, (1886), **33**, 269.

† Esser. Die Uebertragung der menschlichen Aktinomykose auf das Rind. *Preuss. Mitteil.*, (1892), 30; ref: *Jahresb. d. gesammt. Med.*, (1882), **1**, 528.—Ueber Aktinomykose. *Berl. tierärzt. Wchnsch.*, (1889), **5**, 307–309.

* Etiévant, R. Un nouveau cas d'actinomycose temporo-maxillaire; réflexions sur certaines particularités. *Prov. méd.*, (1897), **11**, 7.

* Etiology (The) of Madura Foot. *Indian Med. Gaz.*, (1874), **9**, 100; (1875), **10**, 44–46.

Evans, Willmott. A case of actinomycosis (demonstration). *Brit. Journ. Derm.*, (1903), **15**, 250.—A case of actinomycosis (demonstration). *Brit. Journ. Derm.*, (1906), **18**, 216–217.

† Eve, Frederic S. Case of actinomycosis of the liver. *Trans. Path. Soc.* London, (1888–9), **40**, 405–408; also (Abstr.): *Brit. Med. Journ.* (1889), **1**, 584–585.—† Actinomycosis. *Practitioner*, (1888), **40**, 321–331; also (Abstr.) *Trans. Odont. Soc. Gr. Brit.*, (1887–8), n. s., **20**, 232–256, 1 pl.; ref.: *Lancet*, (1889), **1**, 328.

† Ewetzky, Th. L'actinomycose des conduits lacrymaux. *Arch. d'ophtalmol.*, (1896), 209; ref: *Baumgarten's Jahresb.*, (1896), **12**, 615.

* Eyre. Madura foot. *Indian Ann. Med. Sc.*, (1860), No. 12 (app.), 513.

Fadyeau, J. M. Zur Actinomykose beim Pferde. Ref: *Deutsche Ztschr. f. Tiermed.* (1889), **15**, 444–445.

Fairweather, D. The progress and treatment of a case of actinomycosis commencing in the vermiform appendix. *Brit. Med. Journ.*, (1896), **1**, 1555–1556.—and Dickson, R. S. Case of actinomycosis of the lower jaw. *Brit. Med. Journ.*, (1903), **2**, 1462–1463.

* Fantino, G. Contributo allo studio dell' actinomicosi umana. *Riforma med.*, (1898), **14**, 170; 180; 193.

Farquhar, T. *See* Fox and Farquhar.

* Faulkner, A. Clinical notes on mycetoma (Chionyphe carteri). *Indian Med. Gaz.*, (1886), **21**, 259.

* Fayrer, J. A case of so-called Madura foot. *In his:* Clinical and pathological observations in India, London, (1873), 637–644.

* Fechter, A. Beitrag zur Lehre von der Actinomykose. Halle a. S., 1883.

* Feldbausch. Ein Fall von Aktinomykose des Unterkiefers und der Halsgegend. *Vereinsbl. d. pfäz. Aerzte.*, (1899), **15**, 25–28.

* Félegházi, E. Actinomycosis colli. *Orvosi hetil.*, (1907), **51**, 830.

* Fernández Osuna, G. F. De algunas micosis patólogicas, especialmente de la actinomicosis. *Gac. méd. de Granada*, (1892), **11**, 33; 65; 129; 193; 225; 253, 1 pl.

* Ferré and Faguet. Sur un abcès du cerveau à streptothrix. Association française pour l'avancement des sciences, Session de Bordeaux, (1895), *Sem. méd.*, (1895), 359; also: *Mercredi méd.*, (1895), 441.

Fessler, Julius. Ueber Actinomykose beim Menschen. *München. med. Wchnsch.*, (1889), **36**, 532–534.

* Fibich, S. Actinomycosis miliaris primaria musculorum bovis. *Przegl. wet. Lwów*, (1903), **18**, 41–51.

* Firket, C. L'actinomycose de l'homme et des animaux. *Rev. méd.* (1884), **4**, 273–320.—* Sur un cas d'actinomycose humaine observé à l'hopital militaire d'Ostende, par M. le Dr. Lejeune. *Ann. Soc. méd.-chir.* Liége, (1889), **28**, 17–22.

* Firth, R. H. Note on a case of actinomycosis. *Indian Med. Journ.*, (1886), **5**, 322–325.—* On the nature of the actinomyces or ray-fungus. *Indian Med., Journ.* (1886), **5**, 505–509.

Fischel, Friedrich. Zur Morphologie und Biologie des Tuberkelbacillus. *Berl. klin. Wchnsch.*, (1893), **30**, 989–993.

* Fischer, A. Vorlesung über Bakterien. 1903.

† Fischer, Erwin. Beitrag zur Kenntniss der aktinomykotischen Granulationen und der Histologie aktinomykotischer Herde im Gehirne und seinen Häuten. *Inaug. Dissert.* Tübingen, 1887, 24 pp.; also (Abstr.): *Beitr. z. path. Anat. u Physiol.*, (1883), **2**, 484–492.—* Zur Aetiologie der Aktinomykose. *Centralbl. f. Chir.*, (1890).

* De Fischer, O. Sopra un caso di actinomycosi umana. *Riv. veneta di sc. med.*, (1890), **12**, 558–562; also: *Boll. d. clin.*, (1890), **7**, 341–344.

* Fleming, G. Actinomykosis; a new infectious disease of animals and mankind. *Vet. Journ. & Ann. Comp. Path.*, (1883), **16**, 1; 73; 153; 236; 318.

* Flexner, S. Actinomycosis of the human lung. *Proc. Path. Soc.* Phila., (1899–1900), n. s., **3**, 141–145.—† Pseudo-tuberculosis hominis streptotricha. *Trans. Ass. Am. Physicians*, (1898), **13**, 31–43, 1 pl.; also: *Journ. Exp. Med.*, (1898), **3**, 435–450, 1 pl.

* Floriani, L. Actinomicosi. *Boll. med. trentino*, (1908), **28**, 17; 43.

* Florkiewicz, W. Dwa przypadki promienicy (Aktinomykosis). *Gaz. lek.*, Warsawa, (1885), II, **5**, 911; 938; 955; 984.—* Kilka uwag nad objawami promienicy u ludzi (actinomycosis hominis). *Medycyna*, Warsawa, (1888), **16**, 861–868.

Florman, Arwid. Ueber die Tinction des Actinomyces bovis. *Ztschr. f. wiss. Mikr.*, (1889), **6**, 190–191.

* Folet. De l'actinomycose. *Nord méd.*, (1894–5), **1**, 113–115.—* Actinomycose cutanée de la région fessière. *Écho méd. du nord*, (1903), **7**, 115.

Le Fort. Un cas d'actinomycose. *Écho méd. du nord*, (1906), **10**, 405–406.—Actinomycose cervico-faciale. *Écho méd. du nord*, (1907), **11**, 246.

* Fortineau, L., and Soubrane, L. Note sur un streptothrix chromogène acido-résistant. *Gaz. méd. de Nantes*, (1906), II, **24**, 174.

Foulerton, A. G. R. On streptothrix infections. *Lancet*, (1899), **2**, 779–780.—The pathology of streptothrix infections. *In:* Allbutt's System of Medicine, (1906), **2**, part 1, 302–324.—Pathology of streptothrix infections *Lancet*, (1906), **1**, 970–971.—and Jones, C. Price. Streptothrix infections in the lower animals. *Journ. Comp. Path. & Therap.*, (1901), **14**, 45–59, 1 pl.—On the general characteristic and pathogenic action of the genus Streptothrix. *Trans. Path. Soc.* London, (1902), **53**, 56–127, 1 pl.

* Fox, Tilbury. Fungus foot of India. *Trans. Path. Soc.* London, (1869–70), **21**, 411; (1870–71), **22**, 320.—The so-called "fungus-foot" of India. *Lancet*, (1876), **1**, 190–191.—* and Farquhar, T. On the fungus foot of India; or mycetoma. *In their:* On certain endemic skin and other diseases of India, etc., London, (1876).

* Frarier. Actinomycose pulmonaire. *Prov. méd.*, (1897), **11**, 32; also: *Gaz. hôp. de Toulouse*, (1897), **11**, 105.

* Frèche. Actinomycose de la joue. *Soc. méd. et chir. de Bordeaux*, (1894).—* Actinomycose cervico-faciale. *Soc. méd. et chir. de Bordeaux*, (1894).—See Dubreuilh, W. and Frèche, D.

* Frey, L. Fungus articulat. talo-calcaneae; injection von iodoformäter; Heilung und volkomme Wiederherstellung der Functionsfähigkeit. *Wien. med. Presse*, (1888), **29**, 1325.

* Frey, O. Klinische Beiträge zur Aktinomykose. *Beitr. z. klin. Chir.*, (1897), **19**, 577–608.—* Beiträge zur Aktinomykose. *Inaug. Dissert.*, Tübingen, 1897.

Freymuth. Streptothrixerkrankung der Lunge. *Deutsche med. Wchnsch.*, (1907), **33**, 780–781.

* Freytag, G. W. Beiträge z. Aetiol. d. Aktinomykose. *Dissert.*, München, 1901.

† Frick, K. A. Die Behandlung der Actinomykose und ihre Erfolge. *Deutsche tierärzt. Wchnsch.*, (1896), **4**, 407; ref: *Baumgarten's Jahresb.*, (1896), **12**, 618.

Friedberger, Franz, and Fröhner, Eugen. Lehrbuch der speziellen Pathologie und Therapie der Haustiere. 4. ed, **2**, 506.

* Friedrich. Tuberkulin und Aktinomykose. *Deutsche Ztschr. f. Chir.*, (1896), **43**, 579.—Über Strahlenpilzähnliche Wuchsformen des Tuberkelbacillus im Thierkörper. *Deutsche med. Wchsch.*, (1897), **23**, 653–654.—Ueber die Häufigkeit und operative Prognose der Bauchaktinomykose des Menschen. *Deutsche med. Wchnsch.*, (1906), **32**, II, 1258–1261.—and Nösske, H. Studien über die Localisirung des tuberkelbacillus bei deirecter Einbringung desselben in den arteriellen Kreislauf (linken Ventrikel) und über aktinomycesähnliche Wuchsformen der Bacillenherde im Thierkörper. *Beitr. z. path. Anat.*, (1899), **26**, 470–510, 1 pl.

* Friis, I. Lidt om aktinomykose. *Hosp.-Tid.*, Kjbenh., (1898), IV, **6**, 1153–1158.

* Fröhlich, A. Die Veränderungen des Hufbeins bei Strahlkrebs. *Monatsh. f. prakt, Tierh.*, (1905), **16**, 416–460.

Fröhner, Eugen. *See* Friedberger, Franz and Fröhner, Eugen.

* Fromme, Ant. Ueber die Strahlenpilzähnlichen Bildungen des Tuberkelbacillus, Giessen, 1903.

Fuchs, Ernst. Ueber Färbbarkeit der Streptotricheen nach Methoden der Tuberkelbacillenfärbung. *Centralbl. f. Bakt. etc.*, (1903), Abt. 1, Orig., **33**, 649–656.

Fürbringer, Paul. Beobachtungen über Lungenmycose beim Menschen. *Virch. Arch.*, (1876), **66**, 330–365, 1 pl.

Fütterer, Gustav. A case of actinomycosis hominis, with remarks concerning the differential macroscopic diagnosis between actinomycotic and tuberculous peribronchitis. *N. Y. Med. Journ.*, (1901), **74**, 337–341.—Ein Fall von Aktinomykose der Lunge, der Leber und des Herzens beim Menschen. *Virch. Arch.*, (1903), **171**, 278–284.

* Gabourd. Actinomycose cervico-faciale gauche; actinomycose et grossesse. *Lyon méd.*, (1906), **106**, 62–64.

* Gabriélidès. Diagnostic de l'actinomycose. *Gaz. méd. d'Orient*, (1906), **51**, 72–75.

* Galli-Valerio, B. Richerche sulla probabile esistenza allo state indipendente dell' achorion Schönleinii e dell' actinomyses bovis. *Bull. sc. med. di Bologna*, (1894), VII, **5**, 296–302.—† Actinomycosi e pseudo-actinomicosi, a proposito di un caso osservato nell' uomo. *Gazz. degli osped. e delle clin.*, (1896); ref: *Centralbl. f. Bakt. etc.*, (1897), Abt. 1, **21**, 546.—Contribution à l'étude de la morphologie du Bacillus mallei. *Centralbl. f. Bakt. etc.*, (1899), Abt. 1, **26**, 177–180.

* Gamgee, L. A case of abdominal actinomycosis. *Birmingh. Med. Rev.*, (1904), **55**, 41.

* Gappisch. Zur Kenntnis der aktinomycoseähnlichen Körner in den Tonsillen. *Verhandl. d. deutsch. path. Gesellsch.*, Jena, (1905), 130–139, 1 pl.

Garcin. Ein Fall von Aktinomykose. *Berl. klin. Wchnsch.*, (1895), **32**, 47.

* Garré, K. Behandlung der Aktinomykose etc. Penzoldt-Stintzing, Handbuch der spez. Ther. innerer Krankheiten, **1**.

* Garten. Ueber einen beim Menschen chronische Eiterung erregenden pleomorphen Mikroben. *Deutsche Ztschr. f. Chir.*, (1895), **41**, 257.

* Gascoin, H. L'actinomycose en Touraine. *Gaz. méd. du centre*, (1907), **12**, 7–10.

* Gasperini, G. Recherches morphologiques et biologiques sur un microorganisme de l'atmosphère le Streptothrix foersteri Cohn. *Ann. de micrographie*, (1890), 449–474, 4 pl.—* Sopra una nouva specie appartenante al genere Streptothrix Cohn. *Soc. Toscana de Sc. Nat.*, Pisa, (1891), 267–277.—* Richerce morphologique a biologiche sul genere Actinomyces Harz. *Ann. dell' Istit. d'Igiene speriment. d. r. Univ. di Roma*, (1892), **2**, 167.—* Ulteriori ricerche sul genere Streptothrix come contributo allo studio dell' actinomyces. *Riv. gen. ital. di clin. med.*, (1892), **4**, 210–218.—Versuche über das Genus Actinomyces. *Atti dell'XI congresso med. internaz.* Roma, (1894), **6**, igiene, 80–82. Ref: *Centralbl. f. Bakt. etc.*, (1894), Abt. 1, **15**, 684–686.—* *Proc.-Verb. soc. Tosc. di sc. nat.*, (1895), **9**, 64–292.—† Nove ricerche sull' actinomicosi sperimentale. *Ann. d'ig. sper.*, (1896), n. s., **6**, 475–482; ref: *Baumgarten's Jahresb.*, (1896), **12**, 612–613.

† Gassner. Behandlung der Zugenaktinomykose beim Rind. *Tierärzt. Mitt.*, (1890); ref: *Berl. tierärzt. Wchsnch.*, (1890), **6**, 261.

* Gaucher, Combe and Gastou. Actinomycose du maxillaire inférieur gauche (ostéo-périostite actinomycosique). *Bull. Soc. franc. dermat. et syph.*, (1903), **14**, 61–64; also: *Ann. dermat. et syph.*, (1903), IV, **4**, 152–155.—* Actinomycose de la joue simulant une fluxion dentaire et guérie par le traitement iodoioduré. *Ann. dermat. et syph.*, (1903), IV, **4**, 837.—* Gastou and Chompret. Nouveau cas d'actinomycose du maxillaire inférieur simulant une fluxion d'origine dentaire. *Bull Soc. franc. dermat. et syph.*, (1903), **14**, 94–96; also: *Ann. dermat. et syph.*, (1903), IV, **4**, 228–230.

* Gaudier. À propos d'un cas d'actinomycose du maxillaire supérieur droit chez un enfant, simulant un ostésarcome. Rap. d' A. Broca. *Bull. et mem. Soc. chir.* Paris, (1907), **33**, 125–127.

* Gautier, G. Actinomycose de la face guéri par less ponctures electro-chimiques. *Rev. internat. d'electrother.*, (1891), **2**, 364–366.—See Dauer, J. and Gaûtier, G.

Gedoelst, L. Les champignons parasites de l'homme et des animaux domestiques. Bruxelles, 1902, 199 pp.

* Geissler. Casuistische Beiträge zur Aktinomykose des Menschen. *Breslau. aerzt. Ztschr.*, (1889), **11**, 58–61.—* Ein fall von Actinomykosis des Bauches. Berlin, 1898, 36 pp.

* Gémy and Vincent, H. Affection parasitaire du pied, analogue, si non identique à la maladie dite de Madura. *Congrès dermat. et syph.*, (1892) April 25; also: *Ann. dermat. et syph.*, (1892), May.—* Sur une affection parasitaire du pied non encore décrite (variété du pied de Madura?). *Ann dermat. et syph.*, (1892), III, **3**, 577–584, 1 pl.; also: *Bull. Soc. franç. dermat. et syph.*, (1892), **3**, 258–292, 1 pl.—* Sur un nouveau cas de pied de Madura. *Ann. dermat, et syph.*, (1896), III, **7**, 1253–1262.

* Generali, G. Actinomycosis in un bue. *Atti d. Soc. d. naturalisti di Modena*, (1887), III, **3**.

* Genersich, A. A sugárgomba-betegségröl (actinomycosis). *Orvosi hetil.*, Budapest, (1885), **29**, 925; 953; also, transl. (Abstr.); *Pest. med.-chir. Presse*, Budapest, (1885), **21**, 824; 843.

Gerlach. Cited by Rivolta in his "Sul cosi detto mal del rospo, etc." *Clinica vet.;* rev: *Jahresb. d. gesammt. Med.*, (1878), **1**, 619.

* Gibbes, H. Actinomycosis bovis. *N. Am. Pract.*, (1889), **1**, 337–339.—* Miescher's or Rainey's corpuscles in actinomycosis. *Ann. Surg.*, (1890), **11**, 96–101.

Gilbert. Ueber Actinomyces thermophilus und andere Aktinomyceten. *Ztschr. f. Hyg. u. Infektionskrank.*, (1904), **47**, 383–406, 1 pl.—Noch einmal die Actinomycetenfrage. *Ztschr. f. Hyg. u. Infektionskrank.*, (1905), **49**, 196–198.

Gildersleeve, N. *See* Abbott and Gildersleeve.

* Gill. *Indian Army Medical Reports* (1842).

* Ginsberg, Siegmund. Casuistische Beiträge zur Kenntniss der Actinomykose des Menschan. Berlin, 1890, 30 pp.

* Glaser, Johannes P. A. Ein Beitrag zur Casuistik und klinischen Beurteilung der menschlichen Aktinomykose. *Inaug. Dissert.*, Halle a. S., 1888, 33 pp.

* Glikman, T. S. Primary actinomycosis of the thorax. *Vratch. Gaz.*, (1905), **12**, 501; 566.

Globig. Ueber Bacterien-wachsthum bei 50 bis 70°. *Zeitschr. f. Hyg.*, (1888), **3**, 294–321.

† Gluck. Geschwulstartige Aktinomykose des Unterkiefers. *Berl. med. Gesellschaft*, Sitzung am 4. Nov., 1896. ref: *Deutsche med. Wchnsch.*, (1896), **5**, 210.

* Godfrey. Tubercular disease of the foot. *Lancet* (1843), June 10. Quoted by Manson.—* *Indian Army Medical Reports*, (1884).—* and Colebrook. *Military Medical Reports from the Presidency of Madras* (1850).

Godlee, Rickman J. A series of cases of actinomycosis. *Lancet.* (1901), **1**, 3–9.

* Gohn, H. W. Actinomycosis in a dog. *Journ. Comp. Med. & Veter. Arch.* (1902), **23**, 241.

* Goluhinin, L. E. Dva sluchaya aktinomikoza lyokhkikh. (Two cases of actinomycosis of the lungs.) *Med. obozr.*, Mosk., (1888), **29**, 733–747.

* Golyás, K. Altalános sugárombakór és sugárgomba által okozott elephantiasis egyegy esete szarvasmarhánál. (A case of each of generalized actinomycosis and elephantiasis caused by actinomycosis in horned cattle). Hüsszemle, Budapest, (1906), **1**, 17.

* Gomont, V. Recherches sur les enveloppes cellulaires des nostocacées filamenteuses. *Bull. Soc. bot. France*, (1888), **35**, 204–236, 2 pl.

Gooch, F. L. Is actinomycosis transmissible by cutaneous inoculation. *Journ. Comp. Path. & Ther.*, (1894), **7**, 59–61.

Gottschlich, E. In: Kolle-Wassermann's Handbuch d. path. Mikroorg. (1903), **1**, 136.

* Grancher and Ledoux. Recherches sur la tuberculose zoogléique. *Arch. de méd. exp. et d'anat. path.*, (1889), **1**, 2.

GRANDE ROSSI, F. See Acosta and Grande Rossi, F.

* GRAVES, W. R. A case of actinomycosis. *Intercolon. Med. Journ. Australas.*, (1906), **11**, 38–40.

GRAWITH, PAUL. Ueber Schimmelvegetationen im thierischen Organismus. *Virch. Arch.*, (1880), **81**, 355–376.

* GRECO, N. V. Streptothrix madurae; estudio sobre el parásito del pie de madura ó micetoma. *Anal. Circ. méd. argent.*, Buenos Aires, (1905), **28**, 375–410.

* GREVEN, PAUL CLEMENS. Beiträge zur Kasuistik der Aktinomykose. *Inaug. Dissert.*, Bonn, 1901, 37 pp.

* GRIGORYEFF, V. (Actinomycosis in horses.) *Vet. Obozr.*, Mosk., (1904), **6**, 281–283.

* GRILL, A. Ueber Aktinomycose des Magens und Darms beim Menschen. Tübingen, 1895; also: *Beitr. z. klin. Chir.*, (1895), **13**, 551.

* GRILLO. Contributo allo studio dell' actinomycosi umana. *Riforma med.*, (1898), **14**, 301; 315; 325.

* GRUBER. Über eine neue Eiterung erregende Microbenart: Mycromyces Hofmanni. *Trans. 7th Internat. Congress of Hyg. & Demog.*, London, (1891), **2**, 65.—Mycromyces Hoffmani (Demonstration). *Münch. med. Wchnsch.*, (1891), **38**, 653; also: *Centralbl. f. Bakt. etc.*, (1891), Abt. 1, **10**, 648.—Mikromyces Hofmanni eine neue Hyphomycetenart. *Arch. f. Hyg.*, (1893), **16**, 35–52, 1 pl.—* Zungenaktinomykose bei einem Pferde. *Deutsche tierärzt. Wchnsch.* (1895), 107.

† GRÜN. Lokale Behandlung der Aktinomykose. *München. tierärzt. Wchsnch.*, (1895), **1**; ref: *Jahresb. d. gesammt. Med.*, (1895), **1**, 552.

* GUDER, E. Étude sur l'actinomycose chez l'homme, en Suisse. *Rev. méd. de la Suisse Rom.*, (1891), **11**, 741–758.

* GUÉGUEN, F. Les champignons parasites de l'homme et des animaux. Paris, 1904, 292 pp.

* GUELLIOT, O. Quelques cas d'actinomycose. *Union méd. du nord-est*, (1902), **26**, 1–5.

*GUERMONPREZ and AUGIER. L'actinomycose en Flandre. Lille, 1892; also: *Journ. sc. méd. de Lille*, (1892), **2**, 25; 53; also: *Gaz. hôp.*, (1892), **65**, 162–165.—* and BÉCUE. Actinomycose. Paris, (1894), 271 pp.

* GUILLEMIN. Actinomycose des conduits lacrymaux. *Thèse*, Lyon, 1905.

* GUILLEMOT. Actinomycose cutanée de la joue gauche. *Lyon méd.*, (1896), **81**, 118–126.

* GUINARD. Actinomycose temporo-faciale. *Bull. et mém. Soc. chir. de Paris*, (1900), **26**, 339.—* Actinomycose cervico-faciale chez un sellier. *Bull. et mém. Soc. chir. de Paris*, (1906), n. s., **32**, 994–996.

* DE GUMBLETON, DAUNT, R. Medical letters from Brazil. *Dubl. Med. Press*, (1861), 4.

* GUNTHER. *Indian Army Medical Reports* (1844).

* GUTMANN. Actinomyces bovis. *Arch. f. Veterinärmed.*, (1882); also: *Jahresb. d. gesammt. Med.*, (1882), **17**, I, 528.

HAASS, E. Beitrag zur Kenntnis der Aktinomyceten. *Centralbl. f. Bakt., etc.*, (1905–6), Orig. **40**, 180–186.

HABEL, A. Ueber Aktinomykose. *Virch. Arch.*, (1896), **146**, 1–35.

* VON HACKER. Actinomykose der Zunge. *Anz. d. k. k. Gesellsch. d. Aerzte in Wien*, (1884–5), 185.—Ein Fall von Aktinomykose. *Wien. med. Wchnsch.*, (1885), **35**, 530.—Ueber die seit circa einem Jahre an der chirurgischen Klinik in Innsbruck beobachteten Fälle von Actinomykose. *Wien. klin. Wchnsch.*, (1897), **10**, 302–303.

* Hadden, W. B. Actinomycosis. *St. Thomas's Hosp. Rep. London*, (1884), n. s., **14**, 292.

† Hallander, W. Aktinomykos hos häst. (Actinomykose bei einem Pferd). *Svensk. Veterinärtdskr.*, (1896), **1**, 144; ref: *Jahresb. d. path. Mik.* (*Baumgarten*), (1896), **12**, 618.

* Hallauer, O. Klinischer Beitrag zur Lehre von der Aktinomykose. Zürich, 1892.

* Halter. Ein Fall von Zungenbeinaktinomycose. *Inaug. Dissert.*, Greifswald, 1898.

Hamburger, H. J. Actinomyces im Knochensysteme eines Pferdes. *Virch Arch.*, (1889), **117**, 423–427.

Hamilton, D. J. A text-book of pathology . . . (1894), **2**, pt. 2: Actinomycosis. 1021–1027, 1 pl.; Mycetoma (Madura foot), 1027–1028.

Hamilton, W. F. Actinomycosis from the medical point of view. *Montreal Med. Journ.*, (1905), **34**, 98–100.

* Hamoir. Aktinomykose des Gesäuges beim Schwein. *Ann. méd. vét.*, (1898), ref: *Ztschr. f. Fleisch- u. Milchhyg.*, (1899), **9**, 150.

* Hanau, A. Zwei Fälle von Actinomycose. *Cor.-Bl. f. schweiz. Aerzte*, (1889), **19**, 165–173.

* Hanken, J. H. Een geval van actinomycosis hominis. *Nederl. Tijdschr. v. Geneesk.*, (1887), II, **23**, 494–496.—* Nog twee gevallen van actinomycosis hominis. *Nederl. Tijdschr. v. Geneesk.*, (1890), II, **26**, 282–284.

† Harbitz, F. Bidrag til laeren om actinomyces hominis. (Beiträge zur Lehre von dem Actinomyces hominis). *Norsk Mag. f. Laegevidensk.*, (1898), IV, **13**, 1–44, 1 pl.; ref: *Jahresb. d. path. Mik.* (*Baumgarten*), (1898), **14**, 627–628.

* Hardoüin, P. L'actinomycose à Rennes; 2 nouvelles observations. *Bull. Soc. sc. et méd. de l'ouest*, (1904), **13**, 495–502.

* Harger, S. J. J. Actinobacillosis. *Journ. Comp. Med. & Vet. Arch.*, (1903), **24**, 300–303.

Harley, John. A case of acute tubercle of the liver agreeing completely with the so-called actinomycosis. *Brit. Med. Journ.*, (1885), **2**, 1018–1019; also (Abstr.), *Lancet*, (1885), **2**, 993–994.—* A case of so-called actinomycosis of the liver. *Med.-Chir. Trans.*, London, (1886), **69**, 135–156, 3 pl.; also (Abstr.): *Proc. Roy. Med. & Chir. Soc.* Lond., (1885–6), n. s., **2**, 18–23; also: *St. Thomas's Hosp. Rep. 1885*, Lond., (1886), **15**, 235–237, 1 pl.

Harris, Vincent Dormer. A case of actinomycosis hominis affecting the pleura and lung of the left side; the spleen, the liver, and peritoneoum, and probably the left intestine. *Journ. Path. & Bact.*, (1898), **5**, 182–188, 2 pl.

* Harsha, W. H. Actinomycosis of jaw. *Ill. Med. Journ.*, (1903–4), n. s., **5**, 598.

Hartl, R. Casuistische Beiträge zur Actinomycose bei Thieren. *Berl. tierärzt. Wchnscr.*, (1901), 1–6.

* Hartmann, R. Ein Beitrag zur Actinomycosis hominis. *Allg. med Centr.-Ztg.*, (1888), **57**, 869–871.

* Hartmann, W. M. A case of actinomycosis hominis. *Trans. Chicago, Path. Soc.*, (1903–4), **6**, 157–161.

* Harz, C. O. Actinomyces bovis, ein neuer Schimmel in den Geweben des Rindes. *Jahresb. d. k. Centr.-Thierarzneischule in München* (1877–8) 125–140; also: *Deutsche Ztschr. f. Tiermed.*, supp. (1878); ref: *Jahresb. d. gesammt. Med.* (1879), **14**, I, 626–627.

* Hatch, Wm. Keith. Statistics of cases of mycetoma in the Jamsetjee-Jejeebhoy Hospital during the years 1878–1897. *The Med. Annual* (1900), **18.**—and Childe, F. L. A remarkable case of mycetoma. *Lancet,* (1894), **2,** 1271–1273.

Haussmann. De l'actinomycose. *Arch. gén. méd.,* (1889), **2,** 449–464.

† Havas, J. Behandlung der Actinomycose mit Jodkalium. *Veterinarius,* (1893) (Ungarish); ref: *Jahresb. d. gesammt. Med.,* (1893), **1,** 652.

* Hay, J. Case of actinomycosis. *Liverpool Med.-Chir. Journ.,* (1903), **23,** 405.

Hayes, M. H. Actinomycosis in a horse. *Journ. Comp. Path. & Ther.,* (1896), **9,** 42–44.

* Haynes, H. Is actinobacillosis in Michigan cattle? *Am. Vet. Rev.,* (1903–4) **27,** 337.

† Hebb, R. G. A case of actinomycosis hominis. *Westminst. Hosp. Rep.,* Lond., (1888), **3,** 150–153, 1 pl.; also (Abstr.): *Proc. Roy. Med.-Chir. Soc.* Lond., (1885–7), n. s., **2,** 197–200; also (Abstr.) *Lancet,* (1887), **1,** 313; also (Abstr.): *Brit. Med. Journ.,* (1887), **1,** 331.

* Hegner, J. Ein Fall von Aktinomykosis. *Schweiz. Vrtljschr. f. Zahnh.,* (1907), **17,** 338–340.

Heider, Adolf. Ueber das Verhalten der Ascosporen von Aspergillus nidulans (Eidam) im Thierkörper. *Centralbl. f. Bakt. etc.,* (1890), Abt. 1, **7,** 553–556.

* Heidingsfeld, M. L. Actinomycosis and X-ray therapy. *Am. Journ. Dermat.,* (1903), **7,** 47–49; also: *Lancet-Clinic,* (1903), n. s., **50,** 326.—* Actinomycosis (?) case for diagnosis. *Lancet-Clinic,* (1905), n. s., **55,** 523.

Heine. Behandlung der Actinomycesknoten mit Arsen. *Berl. tierärzt. Wchnsch.,* (1897), 459–460.

Heinick, Emil. Beitrag zur Kenntniss der Bakterienflora des Schweinedarmes. *Arch. f. Thierhlk.,* (1903), **29,** 476–525.

* Heinrichs. Ein weiterer Fall von Aktinomykose des Kehlkopfes. *Arch. f. Laryngol. u. Rhinol.,* (1904), **16,** 350.

* Heinzelmann, G. Die Endresultate der Behandlung der Aktinomykose in der von Bruns'schen Klinik. *Beitr. z. klin. Chir.,* (1903), **39,** 526–557, 2 pl.

* Heller, Arnold. Actinomycose beim Menschen. *Amtl. Ber. ü. d. Versamml. deutsch. Naturf. u. Aerzte,* 1883, (1884), **56,** 147.—Ein Fall von Aktinomykose unter dem Bilde einer acuten Infectionskrankheit verlaufend. *Deutsches Arch. f. klin. Med.,* (1885), **37,** 372–375.

Henry, G. M'W. Actinomycosis. *Lancet,* (1904), **2,** 1449.

† Hering. Cited by Rivolta in his "Sul cosi detto mal del rospo, etc." *Clinica vet.;* rev: *Jahresb. d. gesammt.* Med., (1878), **1,** 619.

* Herlofsen, T. Et tilfaelde af actinomycosis hominis. (Sur un cas d'actinomycose chez l'homme.). *Norsk Mag. f. Laegevidensk,* Kristiania, (1897), IV, **12,** 1141–1143.

Herr, M. *See* Boström and Herr.

Hertwig. Ueber den Actinomyces musculorum der Schweine. *Arch. f. wiss. u. prak. Tierhlknde.* (1886), **12,** 365–372, 1 pl.—Ueber das Vorkommen des Strahlenpilzes (Actinomyces) und der Strahlenkrankkheit (Actinomycosis), nebst Literatur der Actinomycose bei Thieren und Menschen. *Cor.-Bl. d. allg. ärztl. Ver. v. Thüringen,* (1888), **17,** 323–337.

Herz, H. Ueber Aktinomykose des Verdauungsapparates. *Centralbl. f. d. Grenzgeb. d. Med. und Chir.,* (1900), **3,** 561–580; 621–638; 665–668.

* Hesse, G. Ueber Aktinomykose. *Deutsche Ztschr. f. Chir.,* (1892), 274–307, 2. pl.

* Hessert, W. Actinomycosis of the jaw. *Ill. Med. Journ.,* (1905), **7,** 289.

HEUCK, H. Ein Fall von Aktinomykose der Wirbelsäule und Brustwandungen mit Propagation auf die Lunge. *München. med. Wchnsch.*, (1892), **39**, 419–422; 437–440.

HEUSSER, TH. Ein Fall von primärer Actinomycose der Lungen. *Berl. klin. Wchnsch.*, (1895), **32**, 1029–1031.

HEWLETT, R. T. On actinomycosis of the foot, commonly known as Madura foot. *Lancet*, (1892), **2**, 18–19; 506–507.—* On Madura disease (mycetoma) of the foot. *Trans. Path. Soc.* Lond., (1892–3), **44**, 172–177, 1 pl.—Madura foot. (Demonstration of three cases.) *Lancet*, (1893), **1**, 933.

* HEYNE, BENJAMIN. *In:* Historical and statistical news of India (1806).

HICHENS, P. S. Some remarks on a case of actinomycosis of the lungs. *Brit. Med. Journ.*, (1905), **2**, 1168–1171.

HINK, AUG. Lungen-aktinomykosis einer Kuh. *Centralbl. f. d. med. Wiss.*, (1882), **20**, 817–819.

HIRSCH, AUGUST. Der Madura-Fuss. *Arch. f. path. Anat. etc.*, (1863), **27**, 98–115.—Handbuch der historisch-geographischen Pathologie: Madurafuss (fungus disease of India, Mycetoma), 2. ed. (1886), **3**, 483–491.

HOCHE, CL. L. Histogénèse du nodule actinomycosique et propagation des lésions *Arch. méd. exp. et d'anat. path.*, (1899), **11**, 599–613, 2 pl.

* HOCHENEGG, J. Ein Fall von abdominaler Actinomykose. *Wien. med. Presse*, (1886), **27**, 1436.—* Bauchwandactinomycose. *Anz. d. k. k. Gesellsch. d. Aerzte in Wien*, (1886), 161; (1887), 1; also transl. (Abstr.): *N. Y. Med. Journ.* (1887), **45**, 297.—* Actinomycosis abdominis. *Jahresb. d. erst. chir. Klin. d. Prof. Albert in Wien* (1887), 79–84; also: *Wien. med. Presse*, (1887), **28**, 614–618.—* Zur Casuistik der actinomycose des Menschen. *Wien. med. Presse.* (1887), **28**, 537; 582; 613.

* HODENPYL, E. Actinomycosis of the lung. *Med. Rec.*, (1890), **38**, 653–657; also: *Stud. Path. Lab. Coll. Phys. & Surg. Columbia Coll.*, N. Y., (1890–1), **1**.

* HOFFMANN. Ein Fall von Actinomycose der Linken Halsseite. *Monatschr. f. Ohrenh.*, (1905), **39**, 26–31.—* Ueber Aktionomykose des Kehlkopfes und des Kopfnickers. *München. med. Wchsnch.*, (1906), **53**, 449–453.

* HOFMEISTER, F. Ueber eine ungewöhnliche Erscheinungsform d. Blinddarmaktinomykose. *71 Versammlung deutscher Naturforscher und Aerzte in München.* Ref: *Centralbl. f. Chir.*, (1899).

* HOGG, J. The Madura fungus foot of India. *Med. Times and Gaz.*, (1871), **2**, 93.—* Fungus-foot disease of India. *Trans. Path. Soc. London*, (1871–2), **23**, 294–297.—* Mycetoma; the fungus-foot disease of India. *Month. Micr. Journ.*, (1872), **7**, 98–100.—*See also* Moxon and Hogg.

HOLLANDT, RICHARD. Die Zungenaktinomykose des Schweines; neue, krenothrixähnliche Fruktifikationsformen des Aktinomyces in der Zunge und in den Tonsillen. *Arch. f. wiss. u. prak. Tierhlkde.*, (1905), **31**, 417–465.

* HOLMSTEAD, T. Notes on mycetoma, with cases. *Trans. Med. & Phys. Soc.* Bombay, (1870), n. s., **10**, 304–310.

* HOLST, A. Et Tilfaelde af actinomycosis humana. *Norsk. Mag. f. Laegevidensk.*, (1888), IV, **3**, 326–328.

* HOLST, P. F. Lungeaktinomykose. *Forh. med. Selsk. i Kristiania*, (1897), 194–198.

* HORST, A. Ein Fall von Streptothrixpyämie beim Menschen. *Ztschr. f. Heilk.*, (1903), **24**, Abt. f. path. Anat. etc., 157–176, 2 pl.

HOWARD, WILLIAM TRAVIS, *jr.* Actinomycosis of the central nervous system, with the report of a case due to an unidentified member of the actinomyces group. *Journ. Med. Research*, (1903), **9**, 301–323.—A study of actinomyces cultivated from commercial vaccine virus. *Journ. Med. Research*, (1903–4), **10**, 493–512.

* De La Hoz, E. Champignons pathogénes et mycoses du continent américain. Paris, 1905, 125 pp.

* Hudson, W. H. A case of actinomycosis hominis, involving the tissues of the back and lungs. *Ann. Surg.*, (1897), **26**, 626–633.

* Hue, F. Actinomycose cervico-faciale? *Normandie méd.*, (1907), **22**, 9–12.

* Hummel, E. Zur Entstehung der Aktinomykose durch eingedrungene Fremdkörper. *Beitr. z. klin. Chir.*, (1895), **13**, 534–544.

* Huntly, W. Case of Madura in its initial stage. *Glasgow Med. Journ.*, (1889), **32**, 344; (1890), **33**, 339.

* Huth. Ein Fall von Actinomykose des Auges. *Centralbl. f. prak. Augenh.*, (1894), **18**, 106–109.

* Hutyra. Jodkalium bei Aktinomykose. *Ungar. Veterinärber.*, (1896), 38.

* Hyde, James N. A contribution to the study of mycetoma of the foot as it occurs in America. *Trans. Am. Dermat. Ass.*, N. Y., (1896), **19**, 74–90, 1 pl.
* Senn, Nicholas, and Bishop, D. D. A contribution to the study of mycetoma of the foot. *Journ. Cut. & Gen.-Urin. Dis.*, (1896), **14**, 1–15, 1 pl.

† Ignatiew. Ueber Heilung der Actinomykose. *St. Petersb. Journ. f. allg. Veterinärmed.*, (1893), 4; ref: *Jahresb. d. gesammt. Med.*, (1893), **1**, 651.

* Illich, Alberto. Beitrag zur Klinik der Aktinomykose. Wien, 1892, 205, pp. 2 pl.

* Imminger. Zur therapeutischen Behandlung der Actinomycose des Rindes. *Wchnsch. f. Theirh. u. Viehzucht*, (1889), **33**, 149–154.—Die operative Behandlung der Actinomycome beim Rind. Ref: *Berl. thierärzt. Wchnsch.*, (1899), 528.

* Iokheles, L. E. K kazuistikie pechonochnikh abstsessov; sluchai netipichnavo aktinomikosa pecheni. (Liver abscesses; a typical actinomycosis of the liver). *Med. Obozr.*, Mosk., (1902), **57**, 939–950.

* Isemer, Fr. Zwei Fälle von Blinddarmentzündung, hervogerufen durch Actinomyces. *Inaug.-Dissert.*, Greifswald, 1898.

Israel, James. Neue Beobachtungen auf dem Gebiete der Mikosen des Menschen. *Virch. Arch.*, (1878), **74**, 15–53, 4 pl.—Neue Beiträge zur mykotischen Erkrankungen des Menschen. *Virch. Arch.*, (1879), **78**, 421–437.—Ein Schlusswort zur Geschichte der Actinomykose. *Virch. Arch.*, (1882), **88**, 191–194.—Einige Bemerkungen zu Herrn Ponfick's Buch: "Die Actinomykose des Menschen." *Virch. Arch.*, (1882), **87**, 364–380.—† Erfolgreiche Uebertragung der Aktinomykose des Menschen auf das Kaninchen. *Centralbl. f. d. med. Wiss.*, (1883), **21**, 481–482; also: *Verhandl. d. Berl. med. Gesellsch.*, (1882–3), **14**, 194–196.—Demonstration einer erfolgreichen Uebertragung der Actinomycose von Mensch auf Kaninchen. *Berl. klin. Wchnsch.* (1883), **20**, 636.—* Klinische Beiträge zur Kenntniss der Aktinomykose des Menschen, Berlin (1885), 152.—* Ein Beitrag zur Pathogenese der Lungenaktinomykose. *Verhandl. d. deutsch. Gesellsch. f. Chir.* (1886), **15**, pt. 2, 36 to 40, 1 pl.—Ein Beitrag zur Pathogenese der Lungenaktinomykose. *Arch. f. klin. Chirurgie* (1887), **34**, 160 to 164, 1 pl.—* Präparate mit Actinomycosis von der Leiche einer 44 jähr. Frau. *Verhandl. d. Berl. méd. Gesellsch.* (1888), **19**, 3 to 6.—Ein Fall von Bauchactinomykose. *Deutsche med. Wchnsch.* (1889), **15**, 179.—Actinomykose der Coecalgegend. Ref.: *Deutsche med. Wchnsch.* (1895), **21**, Ver.-Beil., 54.—and Wolff, Max. Gelungege Züchtung des Strahlenpilzes ausserhalb des thierischen Organismus und Uebertragung seiner Reincultur auf Thiere *Deutsche med. Wchnsch.*, (1890), **16**, 413 to 414.

Israel, Oscar. Ueber Actinomyces. *Arch. f. path. Anat., etc.* (1884), **96**, 175 to 176.—Ueber die Cultivirbarkeit des Actinomyces. *Arch. f. path. Anat., etc.* (1884), **95**, 140 to 145.—Ueber Doppelfärbung mit Orcëin. *Virch. Arch.* (1886), **105**, 169 to 172.

* Iszlai. Welche Rolle können cariöse Zahne bei actinomykotischen Infectionen haben? *Pest. med.-chir. Presse* (1892).

* Van Iterson. Bijdrage tot de therapie der aktinomykose. *Nederl. Tijschr. v. Geneesk.*, (1892), II., **28**, 838.—* and Siegenbeek van Heukelom. Actinomycose bij den Mensch. *Nederl. Tijdschr. v. Geneesk.* (1889), II., **25**, 329 to 336.

* Iwetzky. Actinomykose der Thränenröhrchen. *Med. Rundschau* (1895).

* Jaboulay. Actinomycose de la paroi abdominale. *Lyon méd.* (1908), **110**, 428.

* Jackson, R. H. Actinomycosis hominis; report of two cases. *Wisconsin Med. Journ.* (1904–5), **3**, 11 to 15.

* Jakimovitch, N. N. (Actinomycosis of lung and pleura.) *Vratch* (1888), **9**, 125; 170; 211; 228; 267.

* Jakowski, M. Promienica pluc i oplucnej. (Actinomycosis of the lungs and pleura.) *Gaz. Lek., Warszawa*, (1897), II., **17**, 10 to 14. *Also, transl.* (Abstr.): *Vestnik med., Kharkov* (1897), **2**, 94.

Jameson, H. *See* Kemper and Jameson.

* Jamieson, S. Streptothrix infections, with special reference to the pulmonary form. *Australas. Med. Gaz.* (1907), **26**, 16 to 23.

Jamieson, W. A., and Edington, Alex. Observations on a method of prophylaxis, and an investigation into the nature of the contagium of scarlet fever. *Brit. Med. Journ.* (1887), **1**, 1262 to 1267.

* Janicke. Ueber Actinomykose des Menschen mit Kranken-Demonstration. *Jahresb. d. schles. Gesellsch. f. vaterl. Kult.* (1888), **66**, 241 to 249.

* Jans, F. W. Ueber einen Fall von Lungenactinomykose. Kiel (1897).

* Jascoin, Henri-Augustin-Jules. Contribution à l'étude de l'actinomycose en Touraine. Bordeaux (1906), 95.

* Jeandin, Joseph. Étude sur l'actinomycose de l'homme et des animaux. Genéve, Bâle, Lyon (1886), 143.

Jeanselme, E. Le tokelau dans l'Indo-Chine française. *C. R. Soc. Biol.* (1901), **53**, 122 to 123.—* Cours de dermatologie exotique, Par. (1904).

† Jelenewski, S. F. Pathologische Histologie und Bacteriologie der Lippenactinomykose beim Rinde (*Actinomyces labii*) (Russisch.). *Abstr.: Arch. f. Veterinärwiss.* (1901), 793; 873; *also Arch. sci. vet.*, **2**, 148; *ref.: Baumgarten's Jahresb.* (1901), **17**, 495 to 497.

* Jensen, C. O. Zur. Kenntnis der Actinomykose. *Monatsch. f. prakt. Thierh.* (1892–3), **4**, 166 to 177.

* Jervell, K. Fire Tilfaelde af Aktinomykose, udgaaende fra Mundhulen. (Four cases of actinomycosis proceeding from the cavity of the mouth.) Festskr.... Prof. Heibergs (etc.), Christiania (1895), 100 to 112.

* Jirou, J. Contribution à l'étude de l'actinomycose en France et en particulier dans la région lyonnaise. Lyon (1893).

* Johne. Aktinomykosis. Bericht über das Veterinärwesen im Königreich Sachsen (1879), 70, 98.—Die Actinomykose ist eine durch Impfung übertragbare Infectionskrankheit. *Centralbl. f. d. med. Wissensch.* (1880), **18**, 881 to 882.—* Aktinomykose der Zunge des Rindes. *Sächs. Vet.-Ber.* (1881), 76.—Weiteres zur Kenntniss des Strahlenpilzes (*Actinomyces bovis*). *Centralbl. f. d. med. Wissensch.* (1881), **19**, 273 to 274.—Aktinomykose. *Ibid* (1882), **20**, 625.—Die Aktinomykose oder Strahlenpilzerkrankung, eine neue Infectionskrankheit. *Deutsche Ztschr. f. Thiermed.* (1882), **7**, 141 to 192, 3 pl.—* Zur Aetiologie der Samenstrangfistel. Bericht über, das Veterinärwesen im Königreich Sachsen (1884), 40.—Aktinomykosis *in* Koch's Encyklopädie d. Thierheilk. (1885), **1**, 51 to 64, 1 pl.—Zur Frage der Actinomykose beim Schwein. *Deut. Zeit. f. Thiermedizin*, Leipzig (1888), **13**, 140 to 145.

JOHNSON, RAYMOND. *See* Carr, Johnson, and Power.

* JONES, C. PRICE. Preliminary note on cultures of streptotrichiae from various sources. *Tr. Path. Soc.* (1900), **51**, 217 to 220.—*See* Foulerton and Jones.

JONES, EDWARD T., and MACKAY, JOHN C. A case of actinomycosis. *Lancet* (1900), **2**, 255 to 256.

JONG, D. A. DE. Actinomykom in dem Schlund eines Rindes. *Deutsche Ztschr. f. Thiermed.* (1886), **12**, 308 to 309.—*Ibid* (1888).

JUDET, E. Les streptothricées et les cladothrix. *Ann. méd.-chir. du centre* (1904), **4**, 32 to 35.

JURINKA. Ein Beitrag zur Zungenaktinomykose. *Beit. z. klin. Chirurgie* (1895).—Zur konservativen Behandlung der menschlichen Aktinomykose. *Mitteil. a. d. Grenzged. d. Med. u. Chir.* (1896), **1**, 139 to 151.

* JUSTO, J. B. Un caso de actinomicosis abdominal. *Rev. Soc. méd. argent.*, Buenos Aires (1904), **12**, 660 to 664.

* KAARSBERG, F. Aktinomykosis. *Nord. med. Ark.*, Stockholm (1887), **19**, 7 to 11.

* KADYAN, A. A. Sluchaĭ khirurgicheskavo lĭecheniya aktinomikoza lyokhkavo. (Case of surgical treatment of actinomycosis of the lungs.) *Trudi Obsh. Russk. Vratch.* (1891–2), **53**, 28 to 37.

* KAEMPFER, E. Perical, i.e., hypersarcosis ulcerosa pedum; sive pedarthrocases, morbus Malabaris vernaculus. *In his:* Amænitatum exoticarum, Lemgoviæ (1712). **3**, 561 to 564.

v. KAHLDEN. (Demonstration eines Präparates von Lungenaktinomykose.) *Münchener med. Wchnsch.* (1898), **45**, 1040.

* KÁLMÁN, B. Aktinomykosis abdominalis egy esete. (Case of:) *Orvosi hetil.*, Budapest (1889), **33**, 550; 564. *Also, transl.* (Abstr.): *Pest. med.-chir. Presse*, Budapest (1889), **25**, 1249 to 1253.

KAMEN, LUDWIG. Ein Fall von primärer Hautactinomycoses. *Wien med. Wchnsch.* (1898), **48**, 834 to 838.

* KAMPELMANN, G. Ein Fall von Aktinomykose der Lunge und Leber. Kiel (1901).

* KANTHACK, A. A. Actinomycosis bovis; the morbid changes in the inferior maxila. *Illust. M. News* (1889), **5**, 29 to 32.—Madura disease (mycetoma) and actinomycosis. *J. Path. and Bact.* (1892), **1**, 140 to 162, 3 pl.—Madura disease of hand and foot. *Lancet*, Lond. (1892), **1**, 195.—On actinomycosis of the foot, commonly known as Madura foot. *Lancet* (1892), **2**, 169 to 170.—* *Tr. Path. Soc.*, Lond, (1892), July 19.—* A pyaemic form of actinomycosis. *Ibid* (1893–94), **45**, 233 to 238.—* Madura Diesase and Actinomycosis. *Ibid* (1894), **45**, 159.—* A pyæmic form of actinomycosis. *St. Barth. Hosp. Journ.* (1896), 51.

* KAPOSI. Actinomycosis. *Jahresb. d. Wien. k. k. Krankenanst.* (1892), **1**, 497.

* KAPPER, F. Ein Fall von acuter Actinomycose. *Wien med. Presse* (1887), **28**, 94 to 96.

KAREWSKI. Aktinomykose der Lunge und der Thoraxwand. *Arch. f. klin. Chir.* (1898), **57**, 576 to 590.—Beitrag zur Lehre von der Aktinomykose der Lunge und des Thorax. *Berl. klin. Wchnsch.* (1898), **35**, 328 to 330; 350 to 353; 373 to 376.—Ein Beitrag zur Lehre von der Aktinomykose des Thorax und der Lunge. *Deutsche med. Wchnsch.* (1898), **24**, Ver.-Beil., 53 to 54.—* and KÜMMELL. Aktinomykose der Lungen. Handbuch d. pract. Chir. **2**, 707.

KARSTEN, H. Der Strahlenpilz; Actinomyces Harz und seine Entwicklungsformen. *Deutsche med. Wchnsch.* (1884), **10**, 264 to 266.

* KASCHENSKI, P. (On the possibility of cultivating actinomyces.) *Med. Obozr.*, Mosk. (1888), **29**, 989 to 996.

KASHIWAMURA, S. Vier Fälle von primärer Lungenaktinomykose. *Arch. f. path. Anat. (etc.)* (1903), **171**, 257 to 278.

* KASPAREK. Die Wurmkrankheit der Rinder (Farcin du boeuf). *Thierärztl. Centralbl.* (1895), 168.

† KATZ, OTTO. Aktinomykose des Unterkiefers (Demonstration). *Berl. med. Gesellsch.* Nov. 6 (1895); *Berl. klin. Wchnsch.* (1895), **32**, 1013.

KEDZIOR. Ueber eine thermophile Cladothrix. *Arch. f. Hyg.* (1896), **27**, 328 to 338.

* KELLOCK, T. H. Actinomycosis of the mouth and face, with the report of ten cases. *Tr. Odont. Soc. Gr. Brit.* (1905), n. s., **37**, 29 to 61.

† VAN KEMPEN, L. Aktinomykom in het strottenhoofd einer Koe. *Holl. (tierärztl.) Ztschr.* (1894), **20**, 332; *Ref.: Jahresb. f. ges. Med.* (1894), **1**, 661.

* KEMPER, G. W. H. Pied de Madura. *In:* Encyclop. intern. chirurgie, **3**, 19.—A case of podelcoma; with microscopic examination of the diseased structure. *Am Pract.* (1876), **14**, 129 to 135.—* and JAMESON, H. A case of podelcoma. *Ibid* (1876), **14**, 577.

* KERN, R. Ein Fall von Actinomycose des Bauchfelles ausegehend vom Blinddarm; operative Behandlung; Heilung. *Cor.-Bl. f. schweiz. Aerzte* (1891), **21**, 570 to 572.

KIEFFER, C. F. A new streptothrix pathogenic for cattle (streptothrix of bovine pneumonia). *Med. Rec.* (1907), **71**, 346 to 349.

KIESERITZKY, GERH., and BORNHAUPT, LEO. Ueber einige unter dem Bilde der Aktinomykose verlaufende Affectionen. *Arch. f. klin. Chir.* (1905), **76**, 835 to 882.

* KIJEWSKI, F. O promienicy u czlowieka. (Actinomycosis in man.) *Gaz lek.*, Warszawa (1889), II., **9**, 50 to 58.

* KINNELL, G. Actinomykosis in Cornwall. *Vet. J. & Ann. Comp. Path.* (1886), **22**, 8 to 10.

KIRKPATRICK, R. C. Mycetoma pedis. *Brit. Med. Journ.* (1900), **1**, 1545.—*See also* Adami and Kirkpatrick.

KISCHENSKY. Ueber Actinomycesreinculturen. *Arch. f. exper. Path. u. Pharmakol.* (1890), **26**, 79 to 87, 1 pl.—* Opiti razvodok chistoi kulturi actinomyces. (On the possibility of cultivating actinomyces.) *Med. Obozr.*, Mosk. (1888), **29**, 989 to 996.

* KITT. Aktinomykose der Haut und des Enters beim Schweine. *Jahresb. ü. d. Leistung. a. d. Geb. d. Vet.-Med.* (Ellenberger-Schütz.) (1906).—* Bakterienkunde, Wien, 4 ed. (1903), 466.

* KLAMANN. Ein Fall von Actinomycose mit Berücksichtigung der in den Abscessen enthaltenen Bacterien. *Allg. med. Centr.-Ztg.* (1897), **66**, 821.

KLEMM. Zur Behandlung der Aktinomykose. *Berl. thierärztl. Wchnsch.* (1889), **5**, 389 to 390.

KNOLL. Allegemeine Actinomycosis des Schweines. *Ibid.* (1891), **7**, 213 to 214.

* KNOX, D. N. Actinomycosis of the cheek and neck. *Glasgow Med. Journ.* (1896), **45**, 382 to 388.

* KNOX, ROBERT. Actinomycosis of the lung with secondary skin infiltration and multiple abscesses of the chest walls. *Med. Press & Circ.* (1902), n. s., **74**, 52 to 54.—A clinical study of actinomycosis. *Lancet* (1904), **2**, 1204–1207.—A case of actinomycosis of the cheek cured by iodide of potassium; with suggestions as to the possible means of infection and spread of the disease. *Ibid.* (1906), **2**, 1213 to 1214.—Another case of actinomycosis. *Ibid* (1906), **2**, 1307 to 1308.

† Koch, Carl. Drei Fälle von Actinomycosis hominis. *München. med. Wchnsch.* (1891), **38**, 216 to 219; 236 to 239. *Also: Sitz.-Protok. d. ärztl. Lokalver. zu Nürnb.* (1890), 66 to 85.—* Ueber das Vorkommen der Aktinomykose in Nürnberg und Umgebung. *Verhandl. d. chir. Section d. 65 Versammlung deutscher Naturforscher und Aerzte.*, Nürnberg (1893).—† Weitere sechs Fälle von Actinomycosis hominis aus Nürnberg und Umgebung. *Münchener med. Wchnsch.* (1894), **41**, 141 to 143; 167 to 169. *Also: Sitz.-Protok. d. ärztl. Lokalver, zu Nürnb.* (1893), München (1894), 42 to 57.—Aktinomykosis der Parotis. *Münchener med. Wchnsch.* (1895), **42**, 177.

Kochanoff, A. V. Three cases of actinomycosis of the abdominal wall. *Izvïest. Imp. Voyenno-Med.* (1905), **11**, 35 to 50.

Köbner. Demonstration eines Präparates von Madurafuss aus Italien. *Monatsh. f. prakt. Dermat.* (1890), **10**, 361.—Pilzpräparat von Madurafuss (Mycetoma pedis) aus Italian (Demonstration). *Berl. klin. Wchnsch.* (1891), **28**, 132. *Also Arch. f. Dermat. u. Syph.* (1891), **23**, 843.

* Koehler, A. Ein Fall von allgemeiner und ein Fall von lokaler Actinomycosis. *Charité-Ann.* (1885–6), **12**, 598.

* Koelsch, Franz, Haver. Klinische Beiträge zur Lehre von der Aktinomykose des Menschen. *Inaug.-Dissert.*, Erlangen (1900).

* Koenig. Ein Fall von Actinomycosis hominis. Berlin (1884).—Aktinomykose aus der Zunge (Demonstration). *Münchener med. Wchnsch.* (1901), **48**, 647.

* Köster, H. Primär actinomycosis pulmonum. *Göteborgs läk.-sällsk. Förh.* (1893), 181 to 186; (1894), 1 to 9.

* Koettnitz, A. Ein Fall von Actinomycose. *Allg. med. Centr. Ztg.* (1888), **57**, 725 to 728.—Zur Behandlung der Actinomykose. *Deutsche med. Wchnsch.* (1891), **17**, 1047 to 1048.

* Koguchi, T. A case of pseudo-actinomycosis. *Chiugai Iji Shinpo*, (1903), 884 to 893.

*Kono. Actinomycosis of the abdonimal walls. *Chiugai Iji Shinpo* (1904), **25**, 1379 to 1390.

* von Korányi, F. Die Strahlenpilzkrankheit, Actinomycosis. Spec. Path. u. Therap., . . . Nothnagel, Wien (1897), **5**, 5. Theil, 1. Abth., 80 to 136, 2 pl.

* Kosark, D. Kaktinomikozu rogat. skota. (Actinomycosis of cattle.) *Vestnik obsh. vet.*, St. Petersb. (1889), **1**, 291; 307.—† Zur Frage über Actinomyces an Getreidegrannen. *Arch. f. Veterinärwissenschaften* (1890), *Ref.: Banumgarten's Jahresb.* (1892), **8**, 395.

* Koshima, U. Actinomycosis following local suppuration. *Chiugai Iji Shinpo* (1904), **25**, 145 to 149.

* Kosinski, J. Ueber Aktinomykose der Bauchhöle. *Medycyna* (1894). *Ref: Centrbl. f. Chir.* (1894).

Kothe, Carl. Tannoform bei der Behandlung von Strahlkrebs. *Berl. thierärztl. Wchnsch.* (1903), 198.

* Kovács, S. The treatment of actinomycosis. *Allatorvosi lapok*, Budapest (1907), **30**, 351.

* Koványi, M. Extirpation of actinomycoma by arsenic. *Allatorvosi lapok.*, Budapest (1907), **30**, 411.

* Kowalewsky, Jean. Die Lungenaktinomykose des Rindes und die Analogie derselben beim Menschen. *Bote f. Öffentl. Veterinärwesen* (1902.)—* (Pulmonary actinomycosis in cattle and its analogy with that of man.) *Vestnik obsh. vet.*, St. Petersb. (1902), **14**, 997; 1044.—Actinomycose mammaire des vaches (mastitis actinomycotica purulenta fibrosa). *Journ. d. méd. vét. et d. zootech.* (1903), V. s., **7**, 513 to 529.—* (Actinomycosis of the testicle of

a bull.) *Arch. vet. nauk.*, St. Petersb. (1904), **34**, 254 to 257.—and SWIATOSLAWKY. Sur la forme miliare de l'actinomycose des ganglions chez les bovidés (Actinomycosis glandi lymphaticae forma miliaris). *Journ. d. més. vét. et d. zootech.* (1900), V., **4**, 331 to 334.

† KOZERSKI, A. Ein Fall von Actinomycose der Haut, behandelt mit grossen Dosen Jodkali. *Medycyna., ref.: Jahresb. d. gesammt. med.*, (1896), **2**, 223.

* KRASNOBAYEFF, T. P. Intestinal actinomycosis in a boy treated by potassium iodide. *Dietsk. med.*, Mosk., (1896) **1**, 233.

* KRATZ, H. Ueber die Aktinomykose um Unterkiefer, zugleich ein Beitrag zur Pathogenese der Aktinomykose. Giessen, 1888.

KRAUSE, PAUL. Beitrag zur Kenntniss des Actinomyces. *Centralbl. f. Bakt. etc.*, (1899), Abt. 1., **26**, 209–12.—Culturen von Streptothrix Aktinomyces (Demonstration). *Münch. med. Woch.*, (1899), **46**, 749.

* KREINBERG, A. Ein interessanter Fall von Aktinomykose beim Rinde. *Ztschr. f. Fleisch.-u. Milchhyg.*, (1905), **15**, 311.

KRUSE, W. Systematik der Streptothricheen *In:* Flügge's Mikroorganismen, (1896), 3. ed., **2**, 48–66.

* KUBACKI, T. Beiträge zur Casuistik und Pathogenese der Actinomycose des Menschen. *Inaug.-Dissert*, Berlin, (1899), 22 pp.

KÜLBS. Ein Fall von Aktinomykose der grossen Zehe. *Wien. klin. Wchnsch.*, (1907), **20**, 49–50.

* KUETZ. *Linnaea*, (1833), **8**, 385.

* KUIPERS, K. R. Actinomycosis bij rundvee. *Tijdschr. v. Veeartsenijk. Maandbl.*, (1903–4), **31**, 565–573.

KUNDRAT. Ein Präparat von Actinomycosis beim Menschen (Demonstration). *Wien. med. Woch.*, (1883), **33**, 482–83.

* KUSCHEFF, N. E. Actinomycosis of lungs and pleura revealed during life in the sputa. *Vratch*, St. Petersb., (1888), **9**, 359.

* LAACHE, S. Actinomycosis hominis intestinalis. *Norsk Mag. f. Laegevidensk.*, (1892), IV, **7**, 1435–1450.

* LABBE, R. Sur un cas d'actinomycose gingivo-jugale. *Bull. med.*, (1906), **20**, 962.

* LACHNER-SANDOVAL, V. Über Strahlenpilze. *Inaugural Diss*, Strassburg, (1898).

* LAKEB, C. Beitrag zur Charakteristik der primären Lungenactinomycose des Menschen. *Wien. med. Presse*, (1889), **30**, 1065; 1108; 1150. Also (Abstr.): *Mitth. d. Ver. d. Aerzte in Steiermark* (1888), **25**, 109–111.

LAMB, D. S. *See* Arwine and Lamb. *See also* Pope and Lamb.

* LANGE. Aktinomykosis der rechten inguinal Gegend. *N. Yorker med. Monatschr.*, (1894), **6**, 175–179.—* Cases of actinomycosis. *Ann. Surg.*, (1896), **24**, 371.

LANGE, B., and MANASSE, P. Actinomyces. (Work referred to by Levy). *Centralbl. f. Bakt. etc.*, (1899), Abt. 1, **26**, 5.

LANGENBECK, VON. Beobachtung in einem Fälle von Wirbelcaries mit Senkungsabscessen. *See:* Israel, *in: Virchows Archiv*, (1878), **74**, 50–51, plate 3, fig. 9, a, b, c, and d. *Ref. in:* Allbutt, System of Med., (1906), 2. pt. 1, 325.

† LANGHANS, TH. Drei Fälle von Aktinomykose. *Correspondenzb. f. schweiz. Aerzte*, (1888), **18**; *ref.; Centralbl. f. Bakt. etc.*, (1888), Abt. 1., **4**, 288–290.

* LASSERRE, JACQUES. Contribution à l'étude du genre Nocardia. (genre Streptothrix Cohn). Description d'une espèce nouvelle, Toulouse, (1904), 142 pp., 1 pl.

* LATIMER, T. S., and WELCH, W. H. A case of intestinal and hepatic actinomycosis in man, associated with leukaemia. *Trans. Assoc. Am. Phys.*, (1896), **11**, 328–339; 1 pl.; *also: Int. Clin.*, (1896), VI., **3**, 152–164, 1 pl.

LAVERAN, A. Au sujet d'un cas de mycétome à grains noirs. *Bull. Acad. méd.*, (1902), III., **47**, 773–776.

* LAW, J. Actinomycosis in animals. *Med. News*, (1883), **43**, 695.

* LAYET, A. Mycétome. Dict. encycl. sci. méd., Paris, (1876), **11**, 172–177.

* LEBEDEFF, I. F. Sluchaĭ aktinomykosa lyokhkikh, raspoznannavo po mokrotïe. (Actinomycosis of lungs diagnosed by sputa). *Med. Obozr.*, (1889), **32**, 417–420.

* LEBERT. Traité d'anatomie pathologique, **1**, 54; and atlas, **1**, pl. 2, fig. 16, Paris, (1857).

* LEBOUROUX, AYMÉ MARIE JEAN. Contribution a l'étude de la maladie dite, pied de Madure, considérée comme une trophonévrose. *Thèse*, Rochefort, (1886), 95 pp., 1 pl., 1 tab.—* Mycétome. *Thèse*, Bordeaux, (1887).

* LEBRAM, F. Ueber Miliar-Actinomykose der Pleura. *Arb. a. d. Geb. d. path. Anat. Inst. zu Tübing.*, (1904), **4**, 297–309.

* LEBRETON, LUCIEN. De l'actinomycose dans le département de la Seine, Paris, (1902), 102 pp., 2 fig.

* LECLAINCHE, E. Actinomycose et botryomycose. *Méd. mod.* (1894), **5**, 131–133.

LEDDERHOSE. Actinomykose der Wangenhaut. *Deutsche med. Wchnsch.*, (1895), **21**, Ver.-Beil., 8.

LEDOUX. *See* Grancher and Ledoux.

† LEGRAIN, E. Sur un cas d'actinomycose de la face. *Ann. derm. et syph.*, (1891), III., **2**, 772–776; *ref.; Jahresb. d. path. Mik.* (Baumgarten), (1891), **7**, 355.—* Actinomycose du menton et du maxillaire inférieur. *Ann. derm. et syph.*, (1895), III., **6**, 682–684; *also: Bull. Soc. franç. derm. et syph.*, (1895), **6**, 338–340.—Sur quelques affections parasitaires observées en Algérie; Actinomycose. *Arch. parasit.*, (1898), **1**, 155–158; Le mycetome, *Ibid.*, 158–162.—*See* d'Audibert and Legrain. *See also* Blanchard.

* LEGRAND, H. Pied de Madura. *Caducée*, (1903), **3**, 262.

LEHMANN, K. B., and NEUMANN, R. O. Actinomycetes. Atlas u. Grund. d. Bakt., (1904), pt. 2, 132–134; 430–505; Cladothrix dichotoma, 509–511.

* LEITH, R. F. C. Actinomycosis of the colon, liver, and lung. *Edinb. Hosp. Rep.*, (1894), **2**, 121–191, 5 pl.

* LÉJEUNE. Actinomycose. *Arch. méd. belges*, (1889), III., **35**, 157; 303.

* LEMIÈRE, G., and BÉCUE, G. Méthode de coloration de l'actinomyces. *Bull. Soc. anat. clin.* Lille, (1892), **7**, 84.

LENEVEU. Sur l'actinomycose de la face. *Rec. méd. vet.*, (1906), **83**, 305–307.

* LEPAGE and CLÉJAT. Actinomycose du maxillaire inférieur. *Journ. méd.* Bordeaux, (1904), **34**, 750.

* LESER. Drei Fälle von Aktinomykose. *Wien. med. Bl.*, (1889), **12**, 345.—† Ueber eine seltene Form der Aktinomykose beim Menschen. *Verhandl. d. deutschen Ges. f. Chir.*, (1889), **18**, pt. 2, 278–287; *also: Arch. f. klin. Chir.*, (1889), **39**, 722–731.

† LESIN. Aktinomykotische Pharyngitis, hompliziert mit einer aktinomykotischen Basilarmeningitis. *Vratch*, (1894); *ref.: Centralbl. f. Bakt. etc.*, (1895), Abt. 1., **18**, 180.

* LEUNIS. Synopsis der Pflanzenkunde, 3. ed., Hannover, (1886).

LEVY, E. Ein neues aus einem Fall von Lepra gezüchtetes Bacterium aus der Klasse der Tuberkelbacillen. Studium über diese Klasse. *Arch. f. Hyg.*, (1897), **30**, 168–183.—Ueber die Actinomycesgruppe (Aktinomyceten) und die ihr verwandten Bakterien. *Centralbl. f. Bakt. etc.*, (1899), Abt. 1., **26**, 1–11.—Die Wachstums-und Dauerformen der Strahlenpilze (Aktinomyceten) und ihre Beziehungen zu den Bakterien. *Centralbl. f. Bakt. etc.*, (1903), Abt. 1., Orig., **33**, 18–23.—and KLEMPERPER, F. Grund. d. klin. Bakt., (1898), 2, ed.; Actinomycose, 361–369; Pathogene Streptothricheen, 369–370.

* Lewis, T. R. The fungus disease of India. *In his:* Phys. and path. researches, London, (1888), 337–390, 2 pl.—* and Cunningham, D. D. The fungus disease of India. Appendix to 11th Rep. San. Com., India, (1875); *also: Indian Ann. Med. Sci.*, (1876), **18**, 354–433.—* Fungus disease of India. Quain's Dict. Med., (1894), **1**, 710.

* Liebman, V. L'attinomice dell'uomo; studi batteriologici. *Resoc. san. Sped. civ.* Trieste, (1889), 17, 138–163; *also: Arch. per sci. med.*, (1890), **14**, 361–402, 1 pl.—* L'actinomicosi con speciale riguardo alla forma polmonare. *Tratt. ital. pat. e terap. med.* (1892), **1**, pt. 2, 413–460, 2 pl.

* Lightfoot. Fungoid disease of the foot, and scrofulous caries of the wrist. *Newcastle and Gateshead Path. Soc.*, London, (1851–2), 26–28.

* Lignieres, J., and Spitz, G. Contribucion al estudio de las afecciones conocidas bajo el nombre de actinomicosis. *Semana med.*, Buenos Aires, (1902), **9**, 223–232, 2 fig.; *also: Rev. Soc. méd. Argentina*, (1902), *also: Bull. Soc. centr. méd. vét.* Paris, (1902), **56**, 487.—Contribution à l'étude des affections connues sous le nom d'actinomycose. *Arch. parasit.*, (1903), **7**, 428–479, 1 pl; *also: Bull. Acad. méd.* Paris, (1903), III, **49**, 125–126.—Contribution à l'étude, à la classification et à la nomenclature des affections connues sous le nom d'actinomycose. *Centralbl. f. Bakt. etc.*, (1904), Abt. 1., Orig., **35**, 294–308; *also: Bull. Soc. centr. méd. vét.*, Paris, (1905), **59**, 64–93.

* Linden, Peter. Aktinomykose, Bonn, (1892), 80 pp.

* Lindt, W., *jr.* Ein Fall von primärer Lungenspitzenactinomycose. *Cor.-Bl. f. schweiz. Aerzte*, (1889), **19**, 262–272.

* Lingard, A. A series of specimens of actinomycosis of the jaws and tongue of oxen, presented to the museum of the Royal College of Surgeons in 1884 and 1885. *Trans. Path. Soc.* London, (1886), **37**, 590–592.

Liphardt. Zur Behandlung von Actinomycesgeschwülsten beim Rinde. *Berl. thierärzt. Wchnsch.*, (1895), 134–135.

* Van Lissa. Bijdrage tot de Jodkaliumbehandeling van Aktinomykose. *Ned. Tijdschr. v. Geneesk.* Amst., (1895).

Litten. Ein Fall von aktinomykotischen Leberabscess. *Deutsche med. Wchnsch.*, (1900), **26**, Ver.-Beil., 17–18.—and Levy, Fritz. Ueber atypische Aktinomykose. *Ibid.*, (1906), **32**, 1772–1775.

Littledale, H. E. Case of human actinomycosis ending fatally, with pathological report. *Brit. Med. Journ.*, (1900), **1**, 1286–1287.—Clinical and pathological notes on a case of human actinomycosis; with conclusions drawn from it as to the nature of the disease. *Ibid.*, (1901), **1**, 12–14.

* Ljunggren, C. A. Om Actinomycosis hominis och dess förekomst inom Jödra Skäne. *Nord. med. Ark.*, (1895), n. s., **5**, 1–26.

Löhlein. Zur Kenntniss der Streptothrixpyämie. *Münchener med. Wchnsch.*, (1902), **49**, 1161–1162.—Streptothrixpyämie bei primärer Bronchopneumonie (mit Demonstration). *Centralbl. f. allg. Path.*, (1908), **19**, Ergänzungsheft: *Verhandl. d. deutsche path. Ges.* 209–211. *See also* Engelhardt and Lohlein.

* Löwe, Reinhold. Statistisches und Klinisches zur Kenntniss der Actinomycose des Wurmfortsatzes und des Coecums, Greifswald, (1904), 64 pp.

van Loghem, J. J. Zur Kasuistik der Streptothrixpyämie. *Centralbl. f. Bakt. etc.*, (1905–6), Orig., **40**, 298–305.

* Lombardo-Pollegrino, P. Di una streptothrix isolata dal sottosuolo. *Riforma med.*, (1903), **19**, 1065–1068.

* Lothrop, H. A. Two cases of actinomycosis hominis, with remarks from a clinical standpoint. *Boston Med. & Sur. Journ.*, (1895), **132**, 300–303.

Lozton, Arthur. Actinomycosis. *Brit. Med. Journ.*, (1906), **2**, 1128.

Lubarsch, O. Zur Kenntniss der Strahlenpilze. *Ztschr. f. Hyg. u. Infectionskrankh*, (1899), **31**, 187–220, 1 pl.

* Lührs, Carl. Beiträge sur Kenntniss der Aktinomykosis des Menschen. Göttingen, (1889), 36 pp.

* Lüning and Hanau. Zur Casuistik der Actinomycosis des Menschen. *Cor.-Bl. f. schweiz. Aerzte,* (1889), **19,** 481–501.

* Lüpke. Zwei neue Fälle von Aktinomykose beim Rinde. *Deutsche tierärzt. Wchnsch.,* (1897), **6,** 223.

* Luginger, J. Streptothricheen als Ursache von Endocarditis beim Rind. *Monatsh. f. prak. Thiermed.,* (1904), **15,** 289–336.

* Lunow. Beitrag zur Diagnose und Therapie der Aktinomykose. *Inaug.-Dissert.,* Königsberg, (1889).

* Maass, F. The method of infection of actinomycosis in man. *Ann. Surgery,* (1903), **38,** 292–299.

* Macaigne and Raingeard. Actinomycose pulmonaire, cutanée et vertébrale. *Gaz. hebd. méd. et chir.,* (1896), 442.—* Actinomycose thoracique, cutanée, vertébrale, pulmonaire. *Presse méd.,* (1898), **1,** 331–335.

* McArthur, L. L., and Hollister, J. C. Actinomycosis of caecum, liver, and lungs. *Med. & Surg. Rep. St. Louis Hosp.,* (1903–4), 66–79.

MacCallum, W. G. On the life history of actinomyces asteroides. *Centralbl. f. Bakt. etc.,* (1902), Abt. 1., Orig., **31,** 529–547.

M'Donald, Stuart. A case of general streptothrix infection. *Trans. Med.-Chir. Soc. Edinb.,* (1904), n. s., **23,** 131–146.

McDowall. *See* Miller and McDowall.

Macé, E. Sur les caractères des cultures du Cladothrix dichotoma Cohn. *Compt. rend. Acad. sci.,* (1888), **106,** 1622–1623.—* Traité pratique de bacteriologie, Paris, (1889), 2. ed., (1892), 3. ed., (1897).—De la décomposition des albuminoides par les Cladothrix (actinomyces). *Compt. rend. Acad. sci.,* (1905), **141,** 147–148.

M'Fadyean, J. Actinomycosis and tuberculosis in the horse. *Journ. Comp. Path. & Therap.,* (1888), **1,** 49–53.—Actinomycosis. *Ibid,* (1889), **2,** 1–21, 1 pl.—The morphology of the actinomyces. *Brit. Med. Journ.,* (1889), **1,** 1339–1344; ref.: *Centralbl. f. Bakt. etc.,* (1889), Abt. 1., **6,** 576–578.—The treatment of actinomycosis. *Journ. Comp. Path. & Therap.,* (1892), **5,** 173; 348–349; (1893), **6,** 163–164.

* McGovern, W. P. Actinomycosis. *Med. News.* (1892), **60,** 99–101.

* Machinski, N. Sluchai aktinomikosa lyokhkikh, raspoznanniy pri zhizni po mokrotie. (Actinomycosis of the lungs diagnosed during life by the sputa). *Ejened. klin. gaz.,* (1888), **8,** 497–507.

* McIntire, H. B. Notes on actinomycosis. *Boston Med. & Surg. Journ.,* (1897), **136,** 84–86.

* McJunkin, F. A. A case of infection with actinomyces bovis. *Physician & Surg.,* (1907), **29,** 213–216.

Macleod, J. M. H. Madura foot. *In:* Allbutt's System of Medicine, 2. ed., (1907), **2,** pt. 2., 754–759.

McMorrow, F. A case of actinomycosis of the pelvis. *Journ. Am. Med. Assoc.,* (1907), **48,** 2029.

* M'Questin, C. Fungus foot of India. *Pacific Med. & Surg. Journ.,* (1873–4), **7,** 552–555.

* McReddie, G. D. Cases of mycetoma. *Indian Med. Gaz.,* (1880), **15,** 156.

Madden, Frank Cole. Two cases of the pink variety of mycetoma. *Journ. Trop. Med.,* (1902), **5,** 243–244; *also: Rec. Egyptian Gov. School Med.,* (1901), 209–212, 1 pl.

Magalhães, P. S. *See* Bulhœs and Magalhães.

* Magli, E. Su di un caso di actinomicosi. *Med. ital.,* (1903), **1,** 1; 22.

* Magnan. L'actinomycose de la face. *Ann, méd.-chir. centr.*, (1904), **4**, 386.
* Magnussen, Lorenzo. Beiträge zur Diagnostik und Kasuistik der Actinomycose, Kiel, (1885), 19 pp.
* Mailland. Tumeur actinomycosique de la mâchoire sur un boeuf. *Lyon méd.*, (1904), **102**, 940.
* Maitland, J. Case of mycétoma of the abdominal wall. *Indian Med Gaz.*, (1898), **33**, 57.
† Majocchi, D. Attinomicosi della pelle umana e di alcuni animali (dermo-actinomicosis). *Atti 12. Cong. Assoc. med. ital.*, (1887), **2**, 367–369; *also: Monatsch. f. prak. Derm.*, (1887), **6**, 1050–1051.—* Appunti clinici e microscopici sopra un raro caso di attinomicosi cutanea dell'uomo. *Ateneo med. parmense*, (1887), **1**, 67–70.—* Sull'uso delle carni alimentari affette dall'-actinomyces. *Ibid*, (1889).—* Contribuzione allo studio della dermo-actinomicosi primitiva dell'uomo. *Atti Cong. gen. Assoc. med. ital.*, (1891), **14**, 455–461; *also: Bull. sci. med. Bologna*, (1892), VII, **3**, 517–539, 1 pl.—* Attinomicosi della bocca e della laringe. *Atti Cong. Soc. ital. laryngol.*, (1892), **1**, 206–210.
* Mallory, F. B. A case of actinomycosis. *Boston Med. & Surg. Journ.*, (1895), **132**, 296–300; *also: Med. & Surg. Rep. Boston City Hosp.*, (1895), VI, 179–189, 1 pl.—and Wright, J. H. Pathological Technique, (1904), 3. ed., Actinomyces, 226–232; Madura Foot, 232–235.
Manson, Patrick. Mycetoma, or the fungus disease of India, *In his: Tropical* diseases, (1899), 568–578; (1906), 3. ed. rev., 680–891; (1907), 4, ed. rev., 759–769.
* Manninger, V. Aktinomycosis-esetek. (Cases). *Orvosi hetil.*, (1906), **50**, 299.
* Mañe, Algorta, and Scaltritti. Un caso de actinomycosis. *Rev. méd. Uruguay*, (1907), **10**, 274–278.
* Maratuech, Léon. Actinomycomes (forme neoplasique) des parois abdominales, Lyon, (1904), 52 pp.
* Marchand. Botanique cryptogamique, Paris, (1883).—Ein Präparate eines Falles von Aktinomykose. (Demonstration). *Berl. klin. Wchnsch.*, (1896), **33**, 334.—* Un cas d'actinomycose. *Journ. chir. et ann. Soc. belge chir.*, (1905), **5**, 419–421.
* Marcus. Beitr. z. Kasuistik und Pathol. der Aktinomykose d. Menschen. *Dissert.* München, 1899.
* Mari, N. (Contributions to the study of actinomycosis). *Uchen. zapiski Kazan. Vet. Inst.*, (1890), **7**, 157; 255; 294; 371.—Häufigkeit der Aktinomykose bei geschlachteten Rindern in Moskau. *Berl. tierärzt. Wchnsch.*, (1890), **6**, 406.—Ueber die Lippenaktinomykose. *Centralbl. f. Bakt., etc.*, (1892), Abt. 1, **12**, 854–855.
* Marotta, R. A. Sobre un caso de actinomicosis torácico. *Anal. Circ. méd. argent.*, (1905), **23**, 341–346.—* and De Luccia Lettieri, E. Un caso de actinomicosis torácica. *Rev. Soc. méd. argent.*, (1906), **14**, 61–70.
* Marten, R. H., and Teichelmann, E. A case of actinomycosis hominis. *Intercolon. Quart. Journ. Med. & Surg.*, (1894), **1**, 209–217.
Martens. Zur Kenntniss der Lungen und Wirbelsäulenaktinomykose. *Arch. f. klin. Chir.*, (1902), **66**, 698–709, 1 pl.
* Martin. Ein Fall von Aktinomykose der Lunge und Bronchien. *Ztschr. f. Fleisch- u Milchhyg.*, **10**, 152.
Martin, C. H. A report of two cases of actinomycosis of the brain. *Journ. Path. & Bact.*, (1896), **3**, 78–86.

* Maseěk, D., Vitez, and Joanovic'. Slucaj aktinomykoma pazuha s klinickog i bakteriolozkog gledista. (A case of actinomycosis of the armpit from a clinical and bacteriological standpoint). *Liec. viestnik u. Zagrebu*, (1904), **26**, 57–60.

* Masoutin, N. G. (Differential distinction of actinomycosis). *Vratch*, St. Petersb (1889), **10**, 433–436.

Massaglio, Aldo. L'actinomycose du doigt. *Presse méd.*, (1904), **12**, 557.

* Mathis. Zur Behandlung der Aktinomykose mit Jodkalium. *Lyon méd.* (1893), 518.

* Matlack, J. A. Actinomycosis. *Interstate Med. Journ.*, (1905), **12**, 453–457.

* Matlakowski, W. Pnypadek wyleczonej promienicy (actinomynosis hominis). *Gaz. lek.*, Warsawa, (1889), II, **9**, 46–50; also, transl.: *St. Petersb. med. Wchnsch.*, (1890), **15**, 232–234.

Matthiesen. Rückenmarkslähmung durch Druck eines Actinomykoms auf das verlangerte Mark. *Berl. tierärzt. Wchnsch.*, (1894), 353.

* Mauri, F. Contribution à l'étude de l'actinomycose. *Rev. med. Toulouse*, (1884), **18**, 617–629.

* Maxutoff, A. (On actinomycosis and morphology of actinomyces). *Meditsina*, St. Petersb., (1893), **5**, 146; 163; 180.

* Maydl, K. Ueber Aktinomykose der Zunge. *Internat. klin. Rundschau*, (1889), **3**, 1713; 1756.

Mayer. Beiträge zur Aktinomykose des Menschen. *Prag. med. Wchnsch.*, (1887), **12**, 161–163.—* Euteraktinomykose beim Schwein, Uebertragung auf die Ferkel. *Wchnsch. f. Tierh. u. Vierzucht*, (1904), **48**, 634.

† Maylard. Ein Fall von Aktinomykose der Appendix. *Glasgow Journ.*, (1896); ref: *Jahresb. d. gesammt. Med.*, (1896), **2**, 222.

† Mayo. Actinomycosis bovis or "lump jaw." *Journ. Comp. Med. & Vet. Arch.*, (1893), **14**, 163; ref: *Jahresb. d. gesammt. Med.*, (1894), **1**, 661.

* Mazzini, G. Fegato con grande ascesso da actinomicosi. *Giorn. d. R. Soc. ed Accad. vet. ital.*, (1899), **48**, 5.

* Mazzoni. Osteomielite fungosa dei metatarsi, dei cuneiformi e del calcagno del piede destro; disarticolazione sotto astragalica col processo Nélaton; guarigione. *In his:* Clin. chir., Roma, (1884), **8–9**, 225.

Mertens, Victor E. Beiträge zur Aktinomykoseforschung. *Centralbl. f. Bakt.*, (1901), Abt. 1, **29**, 649–654.—Beiträge zur Aktinomykoseforschung. *Ztschr. f. Hyg. u. Infekt.* (1903), **42**, 45–89, 2 pl.

* Meunier. Trois nouveaux cas d'actinomycose. *Bull. méd.*, (1895), 675; also: *Gaz. méd. de Paris*, (1895). * Vier Fälle von Gesichts-und Halsaktinomykose. *Acad. de méd.*, Sitzung am 16. Juli 1895. Ref: *Wien. med Presse*, (1895), 1246.

* Meyen. *Linnaea*, (1827), **2**, 433.

* Meyer, P. Sur un cas d'actinomycose chez l'homme. *Mém. Soc. méd. de Strasb.*, (1883–4), **21**, 157–163; also: *Gaz. méd. de Stras.*, (1884), **43**, 99.

* Middeldorpf, K. Ein Fall von Actinomycose. *Amtl. Ber. ü. d. Versamml. deutsch. Naturf. u. Aerzte*, (1883), **56**, 172.—Ein Beitrag zur Kenntniss der Actinomykose des Menschen. *Deutsche med. Wchnsch.*, (1884). **10**, 225–227; 244–245.

† Miklaszewski, W. Ein Beitrag zur Casuistik der Actinomycose des Thorax. *Kronika lekarska* (1896); ref: *Jahresb. d. gesammt. Med.*, (1896), **2**, 321–322.

Mikulicz. Vorstellung eines geheilten Falles von "abdominaller" Actonimycose. *Berl. klin. Wehnsch.*, (1889), **26**, 122–123.

* Milian. Actinomycose de la plante du pied; de la biopsie dans l'actinomycose. *Bull. Soc. franç. dermat. et syph.*, (1905), **16**, 2–4; also; *Ann. dermat. et syph.*, (1905), IV, **6**, 63–65.

* Miller and McDowall. On a somewhat unusual proceeding in surgery, after amputation for mycetoma. *Trans. Med. & Phys. Soc.*, Bombay, (1869), n. s., **9**, (app), 32–34.

* Mills, W. A. Actinomycosis in man. *Dental Cosmos*, (1897), **39**, 949–955.

Milton. Actinomycosis of the cheek. *Lancet*, (1893), **2**, 1222.

* Minas, P. A. Observations in "Keereenagrah" (tuberculosis disease) of the foot. *Indian Ann. Med. Sc.*, (1860–1), **7**, 316.—* Chionyphe carterii, or fungoid foot, with cases, and results of operations. *Indian Ann. Med. Sc.*, (1868), **12**, 281–293.—* Four cases of Chionyphe carterii. *Indian Med. Gaz.*, (1868), **3**, 107.—* Case of Chionyphe carterii. *Indian Med. Gaz.*, (1869), **4**, 140.

* Minges, G. Spontaneous recovery of a case of human actinomycosis inoculated into a rat. *Tri-State Med. Journ.*, (1895), **2**, 263–266.

*Mitteldorf. Ueber die geographische Verbreitung der Aktinomykose beim Rind in Bayern. *Dissert.* Donnauwörth, 1901.

* Mixter, S. J. Actinomycosis of the abdominal wall. *Boston Med. & Surg. Journ.*, (1895), **132**, 303.

Miyamoto. *See* Aoyama and Miyamoto.

Moeller, Alfred. Ein neuer säure-und alkoholfester Bacillus aus der Tuberkelbacillengruppe, welcher echte Verzeeigungsformen bildet. *Centralbl. f. Bakt.*, (1899), Abt. 1, **25**, 369–373, 1 pl.—* On the relations of tubercle bacilli to other bacteria resistant to acids and to actinomyces. *Trans. Brit. Congress on Tuberculosis* (1901).

* Monestié, F. De l'actinomycose cutanée. *Thèse*, Paris, 1895.—* Symptomatologie de l'actinomycose cutanée. *Journ. sc. méd. Lille*, (1895), **2**, 193–206.

* Monti, A. Ueber gewerbliche Infektionen. II. Aktinomykose. *Deutsche klin. ther. Wchnsch.*, (1906), **3**, 810–814; also: *Wien. klin. ther. Wchnsch.*, (1906), **13**, 810–814.

* Montoya y Flores. Micetoma ó pie de Madura. *Anales Acad. med. de Medellin*, (1904), **12**, 379–381.

Moodie, E. L. Large actinomycoma of the liver, secondary to a circumscribed actinomycosis of the upper jaw. *Journ. Path. & Bact.*, (1903), **8**, 239–242.

Moore, Veranus A. Actinomycosis mistaken for tuberculosis at post-mortem following the tuberculin test. *Am. Vet. Rev.*, (1906–7), **30**, 181–184.

* Moore, W. J. On ulcus grave; morbus tuberculosis pedis; podelkoma; mycetoma; Madura foot; or fungus foot disease. *Indian Med. Gaz.*, (1867), **2**, 269.—* Notes on some points connected with fungus foot disease. *Indian Med. Gaz.*, (1873), **8**, 282–284.

* Moosbrugger. Ueber Aktinomykose des Menschen. Tübingen, (1886), 57 pp, 1 pl. also: *Beitr. z. klin. Chir.*, (1886), **2**, 339–395, 1 pl.; also ref: *Centralbl. f. Chir.*, (1886), **13**, 897–902.—* Fünf Fälle von Actinomycosis hominis. *Med. Cor.-Bl. d. würtemb. ärztl. Ver.*, (1892), **62**, 121–124.

Moréno. *See* Desvernine, Albertini, Calnek, Debayle, and Moréno.

* Morgen. Zur Kasuistik der Kehlkopf- und Luftröhren-Aktinomykose. *Ztschr f. Fleisch- und Milchhyg.*, **11**, 366.

* Moroni, A. Di una streptothrix dell'arie. *Arch. sc. med. Torino*, (1898), **22**, 253–257, 1 pl.

Morris, Malcolm. Actinomycosis involving the skin and its treatment by iodide of potassium. *Lancet*, (1896), **1**, 1553–1557, 1 pl.

* Moser, W. Actinomycosis of the liver. *N. Y. Med. Journ.*, (1894), **60**, 176.

* von Mosetig-Moorhof, R. Ein Fall von Pseudo-Aktinomykose seltener Ausbreitung. *Wien. med. Presse*, (1895), **36**, 1–4.

* Mossdorf and Birch-Hirschfeld. Ein Fall von tödlich verlaufener Actinomykose beim Menschen. *Jahresb. d. Gesellsch. f. Nat.- u. Heilk. in Dresden*, (1881–2), 80–88.

† Mosselman and Liénaux. L'actinomycose et son agent infectieux. *Ann. méd. vét.*, Brux., (1890), **39**, 409–426; ref: *Jahresb. d. path. Mikr.* (Baumgarten), (1890), **6**, 407–408.

* Mota, J. E. Actinomycosis. *Bol. Inst. path. Mex.*, (1902), **2**, 1–9.

* Moty. Pseudo-actinomycose du bras-droit. *Bull. Soc. centr. méd. dépt. du Nord*, (1901), II, **5**, 110–113.

† Moussu, G. Progn. und Behandlung der Aktinomykose des Unterkiefers beim Rind. *Rec. méd. vét.*, (1896), 465; ref: *Jahresb. d. gesammt. Med.*, (1896), **1**, 517–518.—* Actinomycose diffuse de la face (vache). *Rev. méd. vét.*, (1905), **82**, 473–480.

* Moxon and Hogg. *Trans. Path. Soc. London*, (1870), **21**, 411.

* Müller, E. Ein Fall von geheilter Aktinomykose. *Mitth. a. d. chir. Klin. zu Tübing.*, (1883–4), **1**, 489–493; also: *Med. Cor.-Bl. d. württemb. ärzt. Ver.*, (1884), **54**, 188.—* Ueber Infektion mit Aktinomykose durch einen Holzsplitter. *Beitr. z. klin. Chir.*, (1887–8), **3**, 355–363, 1 pl.

* Müller, H. Aktinomycose der Lungen und des Thorax. *Cor.-Bl. f. schweiz. Aerzte*, (1901), **31**, 187.

* Müller, K. A. J. Zwei Fälle von Aktinomykose und Carcinom. *Inaug. Dissert.*, Berlin, 1888.

Müller, W. Über Aktinomykose der Brustdrüse. *Münch. med. Wchsnch.*, (1894), **41**, 1027–1030.—Ueber Aktinomykose der Speicheldrüsen. *In* Joh. Orth. Path.-anat. Arb., Berlin, (1903), 278–298.

* Münch, A. W. Ein Fall von Actinomycosis hominis. *Cor.-Bl. f. schweiz. Aerzte*, (1887), **17**, 97; 129.—* Actinomycosis hominis. *Cor.-Bl. f. schweiz. Aerzte*, (1888), **18**, 234–241.

* Muller, L. Ueber Veränderungen im Augenhintergrunde bei miliarer Aktinomykose. *Klin. Monatsbl. f. Augenh.*, (1903), **41**, 236–247.

* Multhaupt, K. B. Beitrag zur Lehre von der Aktinomykose. Kiel, 1888.

* Mundler. Drei Falle der Aktinomykose des Kehlkopfs. *Beitr. z. klin. Chir.*, (1892), **8**, 615.

* Munro, J. C. Four cases of actinomycosis. *Bost. Med. & Surg. Journ.*, (1900), **143**, 255–256.

* Murphy, J. B. Actinomycosis in the human subject. *N. Y. Med. Journ.*, (1885), **41**, 17–19.—* Actinomycosis hominis with report of five cases. *N. Am. Pract.*, (1891), **3**, 593–607, 3 pl.; also: *Chicago Med. Rec.*, (1891–2), **2**, 485–499, 3 pl.

Musgrave, W. E., and Clegg, M. T. The etiology of mycetoma. Report of a case of the ochroid variety occurring in the Philippine Islands and caused by a new species of Streptothrix (Streptothrix freeri), with bibliography by Mary Polk. *Phil. Journ. Sc.* (1907), **2**, B, 477–511, 4 pl.

* Musser, Pearce, and Gwyn. *Trans. Ass. Am. Phys.*, (1901).

Mycetome and Actinomyces. (Report of meeting of the Medical and Physical Society of Bombay, Jan. 8, 1886). (Abstr.) *Lancet*, (1886), **1**, 321.

* Nacciarone. L'actinomicosi nell' uomo. *Riforma med.*, (1896), **12**, 805–808.

NAKAYAMA, H. Impfversuche mit Aktinomyces asteroides Eppinger an Meerschweinchen; zugleich ein Beitrag zur Frage der Ueberempfindlichkeit. *Arch. f. Hyg.*, (1906), **58**, 207–312, 4 pl.

NASSE. Actinomycosis der Schädelbasis. *Berl. klin. Wchsnch.*, (1889), **26**, 120–121.

* NAUSSAC, J. De l'actinomycose pulmonaire. Lyon, (1896).

* NEIRET. Notes médicales à Mayotte. *Arch. méd. nav.*, (1897), **67**, 453.

* NEKAM, L. A streptothrixekröl és némely altaluk élöidézett betegségröl. (The streptothrix and some diseases caused by it). *Orvosi hetil.*, Budapest, (1899), **43**, 12.

* NELHIEBEL. Ueber Aktinomykose. *Tierärzt. Centralbl.*, (1895), 221.

* NETTER. Observation d'actinomycose pulmonaire. *Soc. méd. des hôp.*, (1893).

NEUBER, E. Ueber Aktinomykosis. *Wien. med. Wchsnch.*, (1893), **43**, 1961–1966; 2004–2006; 2049–2051; 2087–2090.

* NEUKIRCH. Ueber Strahlenpilze (Aktinomyceten). Strassburg, 1902.—* À propos d'actinomycose. *Presse méd.*, (1903), **11**, 848.—Zur Actinomycetenfrage. *Ztschr. f. Hyg. u. Infektionskrk.* (1904), **48**, 463–470.

† NEUSCHMIED, J. Fall von Actinomycose der Zunge und der Lymphdrüsen beim Rinde. *Oester. Monatschr. f. Tierhlkd*, (1878), ref: *Jahresb. der gesammt. Med.* (1878), **13**, I, 618.

NEWJADOMSKY. Beitrag zur Lehre von der se überaus pleomorphen Gruppe der pathogenen Cladothricheen. Ref: *Ergeb. d. allg. Path.* (Lubarsch-Ostertag), (1898), **5**, 665–666.

* NEWMAN, J. H. Cases of Madura foot disease. *Indian Med. Gaz.*, (1874), **9**, 322.

NICHOLLS, A. G. The bacteriology of actinomycosis. *Montreal Med. Journ.*, (1905) **34**, 100–103.

NICOLLE, CHARLES. *See* Blanchard. (Rap.)—and PINOY, E. Sur un cas de mycétome d'origine aspergillaire observé en Tunisie. *Arch. parasit.*, (1906), **10**, 437–458, 1 pl.

* NIELLY. Elements de pathologie exotique, (1881).

VAN NIESSEN. Die Actinomyces-Reincultur. *Virch. Arch.*, (1897), **150**, 482–521, 4 pl.

NIKITIN, W. Ein Fall von ausgebreiteter Aktinomykose mit Lokalisation im Gehirn. *Deutsche med. Wchsnch.*, (1900), **26**, 612–613.

* NOCARD, E. Sur un cas d'actinomycose, le premier observé en France. *Arch. vet.*, (1884), **9**, 241–245.—Note sur la maladie des boeufs de la Guadeloupe connue sous le nom de Farcin. *Ann. Inst. Pasteur*, (1888), **2**, 293–302, 2 pl.—† Notes sur l'actinomycose des animaux. *Rec. méd. vét.*, (1892), VII, **9**, 167–182; also transl. (Abstr.): *Am. Vet. Rev.*, (1892–3), **16**, 190–192; and *Journ. Comp. Path. & Ther.*, (1893), **6**, 184–185.—and LECLAINCHE, E. Les maladies microbiennes des animaux. Paris. 3. éd., (1903), **2**, 339–362.

NÖSSKE, H. *See* Friedrich and Nösske.

* VON NOORDEN, W. Ueber fünf weitere Fälle von Aktinomykose, nebst Bericht über die Endresultate der früher an der Klinik operierten elf Fälle. *Beitr. z. klin. Chir.*, (1889–90), **5**, 213–226.

NORRIS, CHARLES, and LARKIN, JOHN H. Two cases of necrotic broncho-pneumonia with Streptothrix. *Journ. Exp. Med.* (1900–1) **5**, 155–193, 6 pl.

* NORTHORP, W. P. Actinomycosis in man. *Proc. N. Y. Path. Soc.*, (1888), 151.

NOSSAL, ALOIS. Die Aktinomykose der Brustorgane (Lunge, Pleura). *Centrbl. f. d. grenzgeb. d. Med. u. Chir.*, (1902), **5**, 465–474; 528–535.

* NOVOTNY. Aktinomykose d. Zunge b. Pferde. *Tierärzt. Centralbl.*, (1897), **20**, 325.

† Obraztsoff, E. S., and Petroff, N. V. (Case of simultaneous actinomycosis and mucorine mycosis in man). *Russk. Med.*, St. Petersb. (1889), **7**, 457–459; ref: *Centralbl. f. Bakt. etc.*, (1890), **7**, 514.

† Ochsner, A. J. Report of a case of actinomycosis. *Chicago Med. Journ. & Exam.* (1886), **53**; *also: Journ. Am. Med. Ass.*, (1886), **7**, 608–610.—* A case of actinomycosis. *Med. News*, (1891), **58**, 97.—* Demonstration of actinomyces. *Chicago Med. Rec.*, (1893), **5**, 154–156.

* Oechsler, Philipp Alfred. Beiträge zur Aktinomykosis hominis. Kiel, 1885, 22 pp.

* Ohage, J. Actinomycosis hominis. *Northwest. Lancet*, (1893), **13**, 1–3; 17.

Okuniewski, J. Aktinomykose nach einer Hautverletzung. *Wien. klin. Wchnsch.*, (1906), **19**, 377.

* O'Leary, J. Aktinomikosis. *Journ. Comp. Med. & Surg.*, (1883), **4**, 183–199.

* Oliver, John C. Report of a case of actinomycosis hominis. *Ann. Surg.*, (1900), **32**, 668–672.

O'Neill, William. A case of actinomycosis. *Lancet*, (1886), **2**, 342–343:

Oppenheim, Moriz. Vortrag über Mycetoma pedis (Madura-Fuss). *Arch. f. Schiffs-u. Trp.-Hyg.*, (1903), **7**, 346–350.—Die pathologische Anatomie des indischen Madurafusses (Mycetoma pedis). *Arch. f. Dermat. u. Syph.*, (1904); **71**, 209–242, 4 pl.

* Orb, M. Zur Kasuistik der Actinomycose. Berlin, (1892).

* Orloff, L. W. (Case of actinomycosis). *Vratch*, St. Petersb., (1889), **10**, 187.—† Aktinomyces in der Kinngegend. *Vratch*, (1889); ref: *Centralbl. f. Bakt.*, (1890), Abt. 1, **7**, 513–514.—Zur Frage der actinomycotischen Erkrankung des Gehirns und der Gehirnhäute. *Deutsche med Wochnsch.*, (1890), **16**, 328–332.

* Ostertag. Zur Jodtherapie der Actinomykose. *Monatsh. f. prakt. Thierhlk.* (1892–3), **4**, 208–214.—† Handbuch der Fleischbeschau für Thierärzte, Aerzte und Richter. Stuttgart, (1899); rev: by Johne in: *Ztschr. f. Thiermed.*, (1899), **3**, 339–340.

*Otto, Clemens. Zur Kenntniss der klinischen Verlaufes der Aktinomykose des Menschen. Berlin, (1892), 30 pp.

Ouervain, F. de. *See* Bourquin and Oervain.

* Palagi, A. Caso di rantolo in un vitello prodotto da actinomicosi esofagea. *Giorn. anat. fisiol. e patol. d. animali*, (1884), **16**, 312–314.

Paltauf, Rich. Präparate einer Aktinomykose des Herzens und des Herzbrutels. (Demonstration). *Wien. klin. Wchnsch.*, (1890), **3**, 487.—† Über Madurafuss. *Internat. klin. Rundschau*, (1894); also: *Centralbl. f. allg. Path.*, (1894), **5**, 865; also: *Wien. klin. Wchsnch.*, (1894), **7**, 466.—Madurafuss (Demonstration). *Arch. f. Dermat. u. Syph.*, (1903), **66**, 212–213.

* Paolo, Lombardo Pellegrino. Di una streptothrix isolata dal sotlosuolo. *Riforma med.*, (1903), **19**.—Sul comportamento delle streptotricee e di alcuni bacteri nei grassi. *Ann. ig. sper.*, (1904), n. s., **14**, 533–575.

* Park, R. Actinomycosis. *Buffalo Med. & Surg. Journ.*, (1892–3), **32**, 325–337.

* Parmenter, F. J. Infection with an organism of the actinomyces group; recovery. *Buffalo Med. Journ.*, (1907–8), **63**, 133–137.

* Parrascandolo, C. Actinomicosi del polmone del punto di vista chirrurgico. *Clin. med.*, (1900), **6**, 353–359.

* Partsch, K. Zwei Fälle von Actinomykosis. *Jahresb. d. schles Gesellsch. f. vaterl. Kult.*, (1881), **59**, 50–53; also: *Breslau. ärztl. Ztschr.*, (1881), **3**, 78.—* Einige neue Fälle von Aktinomykose des Menschan. *Deutsche Ztschr.*

f. Chir., (1885–6), **23**, 497–529.—* Die Actinomykose des Menschen vom klinischen Standpunkte besprochen. *Samml. klin. Vortr.*, (1888), No. 306–307.—* Aktinomykose der Augenlider. *Centralbl. f. prakt. Augenh.*, (1893), **17**, 161–165.—Die Eigangspforte des Aktinomyces. *Wien. klin. Wchsnch.*, (1893), **6**, 97–99.—* Die Aktinomykose der Mundorgane. Handbuch der prak. Chirurgie, (1900), **1**, 1022.

* DE PASO FERNÁNDEZ-CALVO, J. La actinomycosis. *Gac. méd. de Granada*, (1884), **2**, 267; 298; 328.

PATTON, W. S. Mycetoma (Madura foot) in the Yemen. *Brit. Med. Journ.*, (1906), **1**, 1401.

* PAULY and POLLOSSON, A. Nouveau cas d'actinomycose. *Mém et comp.-rend. Soc. sc. méd. de Lyon*, (1893), **33**, 67.

* PAWLOWSKY, A. (Remarks on actinomycosis). *Ejend. klin. gaz.*, St. Petersb., (1883), **3**, 356.—and MAKSUTOFF, MAG. Sur la phagocytose dans l'actinomycose. *Ann. Inst. Pasteur*, (1893), **7**, 544–549.

PEARCE. *See* Musser, Pearce and Gwyn.

* PEARSON, L., and RAVENEL, M. P. A case of pneumycosis due to the Aspergillus fumigatus. *Journ. Comp. Med. & Vet. Arch.*, (1900), **21**, 451–465.

* PEASE, H. T. Actinomycosis in the buffalo. *Vet. Journ. & Ann. Comp. Path.*, (1891), **33**, 14.

PELAGATI, MARIO. Blastomyceten und hyaline Degeneration. *Virch. Arch.*, (1897), **150**, 247–259.

* PELLEGRINI, L. Un caso di actinomicosi della faccia. *Boll. Soc. med. prov. Bergamo*, (1894), **5**, 55–59.

* PELLETIER. Mycétome à grains rouges observé à Saint Louis (Sénégal). *Ann. hyg. et med. colon.*, (1906), **9**, 578–580.

† PERINNI. Ueber die actinomicose der Rinder. *Koch's Monatschr.*, **18**, 168; *Jahresb. d. gesammt. Med.*, (1893), **1**, 652.

* PERRIN. Actinomycose cutanée de la face. *Ann. derm.*, (1899).

* PERRONCITO, E. Osteosarcoma della maxilla anteriore e posteriore net bovini. *Patologia Enciclopedia agraria italiano*, (1875), **3**, 599.—* L'aktinomyces bovis (Harz) ed i sarcomi nei bovini. *Ann. accad. agric.*, (1878).—* Ueber den Actinomyces bovis und die Sarkome der Rinder. *Deutsche Ztschr. f. Thiermed.*, (1879), **5**, 33–40, 1 pl.—Inoculation d'actinomyces accidentellement survenue à un cheval. *Arch. ital. biol.*, (1886), **7**, 143–144.—* Ancora sulla priorità dell' osservazione dell' actinomyces bovis per íl Prof. S. Rivolta. *Med. vet.*, (1887), **34**, 337–352; also: *Gior. d. R. Accad. med. Torino*, (1887), III, **35**, 352–366.

* PERTIK, O. Reklinghausen és De Bary intézeteiböl Strassburgban. Az actinomycosisról íy eset kapcsán az embernél. *Orvosi hetil., Budapest*, (1884), **28**, 1133; 1161; 1194, 1 pl.—Untersuchungen über Aktinomykose. *Wien. med. Wchnsch.*, (1885), **35**, 307.

PETGES, G. Notes sur l'actinomycose, en particulier dans la région bordelaise. *Gaz. hebd. sc. med. Bordeaux*, (1907), **28**, 145–151.

† PETIT, G. Actinomycose des bourses consécutives à la castration, chez le boeuf. *Bull. Soc. centr. méd. vét.*, (1903), n. s., **21**, 328–330; also: *Bull. et mém. Soc. anat.* Paris, (1903), **78**, 635.

* PETROFF, N. V. (Actinomycosis). *Russk. med.*, (1888), **6**, 186–189.—* (Actinomycosis in man). *Dnevnik Obsh. Vratch. g Kazani*, (1888), **12**, 41–58.—Ein Beitrag zur Lehre von der Aktinomykose. *Berl. klin. Wchnsch* (1888), **25**, 541–544.—*See also* Obraztsoff and Petroff.

† Petruschky. Demonstration von Präparaten und Kulturen von einem zweiten intra-vitam diagnostizierten Fälle von Streptotrichosis hominis. *Verhandl. d. Cong. f. inn. Med.*, Wiesbaden, (1898), **26**, 557–561; also: *Centralbl. f. allg. Path. u. path. Anat.*, (1898), **9**, 394–395; also: *Deutsche med. Wchsnch.*, (1898), **24**, Ver.-Beil. 78.—Die pathogenen Trichomyceten Streptothrix, Cladothrix, Leptothrix. *In:* Kolle und Wasserman's Mikroorganismen, (1903), **2**, 832–860.

Petterson, Alfred. Untersuchungen über säurefeste Bacterien. *Berl. klin. Wchnsch.*, (1899), **36**, 562–566.

* Peyre, Raymond. Symbiose actinomycosique; la symbiose morphologique et fonctionelle de l'actinomyces éclairée par quelques autres symbioses microbiennes. Lyon, (1903), 71 pp.

* Pflug, G. Lungenactinomycose in Form acuter Miliartuberculosis. *Oester. Vrtljschr. f. wiss. Veterinärk.*, (1882), **58**, 13–30, 1 pl.—Lungenaktinomykosis in Form acuter Miliartuberculosis bei einer Kuh. Vorläufige Mittheilung. *Centralbl. f. d. med. Wiss.*, (1882), **20**, 241–243.—† Ueber Actinomycosis. *Oester. Vrtljschr. f. wiss. Veterinärk.*, (1882), **58**, 1–12; *also abstr.:* Berl. *klin. Wchnsch.*, (1882), **19**, 46.—Die nicht actinomykotische Holzzunge des Rindes. *Deutsche Ztschr. f. Thiermed.*, (1891), **17**, 109–120.

* Phokas, G. Klinikón mathema peri aktinomukoseos. *Iatrike proodos, En Suro*, (1903), **8**, 1–5.

* Piana, G. P. Actinomicosi incipiente; actinomyces svilupparto alla superficie di frusti di tessuto vegetale fibro-vascolare incisti dati sotto la mucosa della lingua nei bovini. *Arch. sc. med.* (1886–7), **10**, 137–141, 1 pl.

* Pic, A. De l'actinomycose pulmonaire. *Prov. méd.*, (1896), **10**, 121; 135; 169.

Pieroni, N. Actinomycose der Méningen cérébrales (Una nouva forma actinica). *Journ. méd. vét.*, (1900), V, **4**, 245–246.

Piettre. *See* Brocq-Rousseu and Piettre.

* Pilliet, A. H. L'actinomycose; Nocardia bovis. *Pròqres méd.*, (1897), III. **6**, 257–260.

† Pilz. Actinomycose bei einem Pferde. *Ztschr. f. Veterinärk.*, **5**, 12; *Jahresb. d. gesammt. Med.*, (1893), **1**, 653.

Pinoy, E. Reproduction experimentale du mycétome à grains noirs. *Compt. rend. Acad. sc.*, (1906), **143**, 1175–1176.—*See also* Nicolle and Pinoy.

* Pir, Zahor. Amputation for treatment of mycetoma. *Indian Lancet*, (1901), **17**.

* Pitt. Primäre Lungenaktinomykose beim Rinde. *Ztchr. f. Fleisch-u. Milchhyg*, **10**, 134.

Plehn, Albert. Das Mycetoma. *In:* Mense's Handbuch der Tropenkrankheiten, (1905), **1**, 42–47.

* Plehn, F. Die Kamerkunküste (1896).

* Plicque, A. F. L'actinomycose chez l'homme et chez les animaux. *Gaz. hôp.*, (1890), **63**, 705–711.

* Plönnigs, J. F. A. Ein Fall von primärer Lungenaktinomykose hervogerufen durch Aspiration einer Gerstenähre. Greifswald, 1898.

* Poissonnier. Actinomycose cutanée de la face. *Lyon méd.*, (1905), **104**, 869.

* Poli, C. Asceso perilaringeo da actinomicosi. *Boll. R. Accad. med.* Genova, (1893), **8**, 42–44; *also: Gazz. d. osp.*, (1893), **14**, 322.

Pollosson, A. *See* Pauly and Pollosson.

Polverini, G. Richerche e oservazioni sul piede di Madura. *Sperimentale*, (1903), **57**, 659–684; also (Abstr.): *Journ. Trop. Med.*, (1905), **8**, 63–64; also (Review): *Monatsh. f. prak. Dermat.*, (1904), **38**, 575.

* Polyanski, N. S. (Co-existent actinomycosis and tuberculosis in man). *Prakt. Vratch*, St. Petersb., (1905), **4**, 275; 296.

* Poncet, A. Une observation d'actinomycose de la joue et du maxillaire inférieur droit, avec propagation au poumon drout. *Gaz. hebd. méd.*, (1893), II, **30**, 40–42; also (Abstr.): *Mém. et compt.-rend. Soc. sc. méd. de Lyon*, (1892), **32**, 166–168.—* Actinomycome de la région cervico-faciale droite. *Mercredi méd.*, (1895), **6**, 349–351.—* De l'actinomycose humaine à Lyon. *Gaz. hebd. méd.*, (1895), **42**, 182–190.—* Huit cas d'actinomycose humaine. *Prov. méd.*, (1895), **9**, 205–207. also: *Rev. gén. clin. et ther.*, (1895), **9**, 259.—* Note sur un nouveau cas d'actinomycose de la face, joue et région temporo-maxillaire gauches; médication iodurée; guérison. *Mercredi méd.*, (1895), **6**, 289–291.—* Note sur un nouveau cas d'actinomycose temporo-maxillaire; considération cliniques. *Lyon méd.*, (1895), **80**, 519–530.—* Pseudo-actinomycose cervico-faciale; phlegmons suppurés de la face, du cou, dus à une nouvelle mycose à gros grains jaunes. *Assoc. franc. chir. Proc.-verb. etc.*, (1896), **10**, 66–73.—* Sur un cas de fistule trachéo-oesophagienne d'origine actinomycosique. *Bull. Acad. méd.*, (1896).—* L'actinomycose en France pendant l'anne 1896. *Courrier méd.*, (1897), **47**, 142–144.—* De l'actinomicose anorectale. *Bull. Acad. méd.*, (1898), III, **40**, 168–176; also: *Gaz. hebd. méd.*, (1898), n. s., **3**, 889–891; also: *Gaz. hôp.*, (1898), **71**, 981–984.—* Un cas d'actinomycose. *Lyon méd.*, (1902), **99**, 873.—* Actinomycose angulo-et temporo-maxillaire droite. *Lyon méd.*, (1903), **100**, 607.—Actinomycose de la face. *Lyon méd.*, (1903), **101**, 48.—Actinomycose fessière d'origine rectale; vaste phlegmon actinomycosique de la fesse et de la partie postéroexterne de la cuisse gauche. *Gaz. hôp.*, (1903), **76**, 325–330.—* Otite actinomycosique; mastoïdite et phlegmon cervical de meme nature. *Lyon méd.*, (1903), **100**, 903–905.—* Actinomycose cervico-faciale aiguë, a forme-angulo-temporo-maxillaire; phlegmon lihneux caractéristique. *Lyon méd.*, (1904), **103**, 20–24.—Trismus au constriction actinomycosique des machoires. *Gaz. hôp.*, (1904), **77**, 237–239.—* Actinomycose cervico-faciale de forme très douloureuse; névrite actinomycosique. *Lyon méd.*, (1905), **104**, 1398–1402.—* Pseudo-actinomycose; mycose à graines jaunes. *Lyon méd.*, (1905), **105**, 261–263.—* Actinomycose temporomaxillaire gauche; actinomycose et grossesse. *Lyon méd.*, (1906), **106**, 226–229.—* Actinomycose à forme néoplasique. *Lyon méd.*, (1907), **109**, 533–535.—* Actinomycose cervicale. *Lyon méd.*, (1907), **109**, 259.—* Actinomycose de la joue droite. *Lyon méd.*, (1907), **109**, 949.—and Bérard, L. De l'actinomycose humaine, particulièrement en France. *Lyon méd.*, (1897), **85**, 467–477; also: *Odontologie*, (1897), II, **5**, 214–222; also: *Méd. mod.*, (1897), **8**, 499; also: *Bull. méd.*, (1897), **11**, 737; 925.—* Étiologie générale de l'actinomycose. *Lyon méd.*, (1898), **88**, 5–13; also: *Prov. méd.*, (1898), **12**, 202–204.—* Traité clinique de l'actinomycose humaine, pseudo-actinomycoses et botryomycoses. Paris, (1898), 410 pp, 4 pl.—† De l'actinomycose humaine pendant ces deux dernières années (1898–1900). *Bull. Acad. méd.*, (1900), III, **43**, 394–402; also: *Tribune méd.*, (1900), II, **33**, 265–268; also: *Gaz. hebd. méd. et chir.*, (1900), **47**, 289–292; *Gaz. hôp.*, (1900), **73**, 365–368; *Lyon méd.*, (1900) **93**, 433–441.—Die Aktinomykose. *Münch. med. Wchnsch.*, (1900), **47**, 815.—† De l'actinomycose humaine en France. Sa fréquence. Son pronostic éloigné. *Bull. Acad. méd.*, (1902), III, **47**, 424–432, 1 fig; also: *Gaz. hop.*, (1902), **75**, 377–380; *Tribune med.* (1902), II, **35**, 288–290; *Méd. mod.*, (1902), **13**, 125–126; *Presse méd.*, (1902), **10**, I, 437.—* Actinomycose angulo et temporo maxillaire droite. *Lyon méd.*, (1903), **100**, 607.—* Actinomycose de la face. *Lyon méd.*, (1903), **101**, 48.—* Actinomycose fessiére d'origine rectale, vaste phlegmon

actinomycosique de la fesse et de la partie posteroexterne de la cuisse gauche. *Gaz. hôp.*, (1903), **76**, 325.—* Otite actinomycosique; mastoïdite et phlegmon cervical de meme nature. *Lyon méd.*, (1903), **100**, 903.—† À propos du diagnostic clinique de l'actinomycose humaine. *Ann. méd.-chir. du centre*, Tours, (1904), **4**, 179–182; also: *Arch. gén. méd.*, (1904), **1**, 805–810; *Lyon méd.*, (1904), **102**, 609–619; *Bull. Acad. méd.*, (1904), III, **51**, 250–260; *Méd. mod.*, (1904), **15**, 89–91; *Arch. parasit.*, (1903–4), **8**, 548–557.—* Actinomycose cervico-faciale aiguë, à forme angulo temporo maxillaire; phlegmon ligneux caraceristique. *Lyon méd.*, (1904), **102**, 20.—* Trismus ou constriction actinomycosique des machoires. *Gaz. hop.*, (1904), **77**, 237.—* Actinomycose cervico-faciale de forme très douloureuse; névrite actinomycosique. *Lyon méd.*, (1905), **104**, 1398, with plate.—and Thévenot, L. † De l'actinomycose humaine en France et à l'étranger dans ces cinq dernières années. *Bull. Acad. méd.*, (1903), III, **49**, 722–731; also: *Bull. méd.*, (1903), **17**, 543–545; *Lyon. méd.*, (1903), **100**, 1055–1065; *Gaz. hop.* Lyon, (1903), **4**, 396–406.—and Lacomme and Thévenot, L. Recherches sur la toxicité des cultures d'actinomycose et la présence de leurs produits solubles. *Bull. Acad. méd.*, (1907), III, **57**, 449–452.

* Ponfick, E. Ueber das Vorkommen eigentümlicher gelblicher Körner u. s. w., *Bresl. ärzt. Ztschr.*, (1879).—† Ueber Actinomycose. *Berl. klin. Wchsnch.*, (1880), **17**, 660–661; also: *Verhandl. d. Berl. med. Gesellsch.*, (1880–1), **12**, 6–10.—* Ueber Actinomycose des Menschen und der Thiere. *Allg. Wien. med. Ztg.*, (1880), **25**, 431; also: (Abstr.): *Pest. med.-chir. Presse*, (1880), **16**, 936.—Die Aktinomykose des Menschen; eine neue Infektionskrankheit. Festchrift zum 25 jähringen Jubiläum Virchows. Berlin 1882, 138 pp. 6. pl.—Letztes Wort zur Aktinomykosen-Frage. *Arch. f. path. Anat. etc.* (1882), **88**, 195–196.—Zur Geschichte der Actinomykose. *Arch. f. path. Anat., etc.*, **87**, 541–561.—* Ueber Actinomycose ohne Actinomyces. *Jahresb. d. schles. Gesellsch., f. vaterl. Kult.*, (1884), **62**, 104–108; also: *Breslau. aerztl. Ztschr.*, (1885), **7**, 30.—Actinomycosis. *In* Twentieth Century Practice of Medicine, (1898), **15**, 475–493.

* Pont. À propos d'un cas d'actinomycose cervico-faciale suppurée à forme mentonnière en fer à cheval. *Odontologie*, (1904), II, **18**, 65–68.

* Pope, B. F., and Lamb, D. S. Mycetoma, the fungous foot of India. *N. Y. Med. Journ.*, (1896), **64**, 386–388.

* Popelin. Arthrite fonguese du pied. *Presse méd. belge*, (1888), **40**, 241.

* Porauer. Beitrag zur Zungenaktinomykese des Rindes. *Veterinarius*, (1898).

* Porter, C. A. Actinomycosis. *Boston Med. & Surg. Journ.*, (1900), **143**, 251–254.

* Porto, N. Um caso de actinomycose. *Journ. Soc. sc. med.* Lisb., (1894), **58**, 483–489.

* Pourpre. De l'actinomycose cutanée primitive de la face. *Thèse*, Lyon, (1901); also: *Lyon méd.*, (1902).

* Powell. Actinomycosis of the Lungs. *In* Diseases of the Lungs and Pleurae. London, (1893).

† Powell, R. D., Godlee, R. J., and Taylor, H. H. On a case of actinomycosis hominis; with an appended report on the morphology of the fungus by Edgar Crookshnak. *Med.-Chir. Trans.*, London, (1889), **72**, 175–208; 3 pl.; also, (Abstr.): *Proc. Roy. Med. & Chir. Soc.*, Lond., (1888–9), III, **1**, 67; 68–72; *Lancet*, (1889), **1**, 328; *Brit. Med. Journ.*, (1889), **1**, 357–358.

* Powell, S. Report on Mycetoma carterii, or fungus disease of India. *Army Med. Dep. Rep.*, (1896), **37**, 335–349.

POWER, D'ARCY. *See* Carr, Johnson and Power.

* PRALLE, G. Ueber Aktinomykose unter besonderer Berücksichtigung der Farbemethoden. Freiburg i. B., 1892.

* PRASAD, K. Notes on a case of fungus disease of India (mycetoma or Madura foot). *Indian Med. Gaz.*, (1906), **41**, 139.

* PREIS, K. Actinomycosis esete. (Case). *Orvosi hetil.*, Budapest, (1905), **49**, 505.

PREUSSE. Enzootisches Aufreten der Aktinomykosis in den Kreisen Elbing und Marienburg (Westpr.). *Berl. tierärzt. Wchsnch.*, (1890) **6**, 17–21.

* PRIETSCH. Ausgebreitete Aktinomykose bei einer Kuh. *Ber. d. Vetw. i. Kgr. Sachs.*, **39**, 96.

† PRINGLE, J. J. A case of actinomycosis involving the skin extensively. *Med.-Chir. Trans.*, (1894–5), **78**, 21–37, 1 pl.; also: (Abstr.): *Brit. Med. Journ.*, (1894), **2**, 1108; *Lancet*, (1894), **2**, 1154.—Actinomycosis of the cheek. *Brit. Journ. Dermat.*, (1896), **8**, 19–22.

*PROTOPOPOFF, N., and HARMER, H. Ein Beitrag zur Kenntniss der Aktinomycesculturen. *Ztschr. f. Heilk.*, (1890), **11**, 255–266, 1 pl.

* PROVERA, C. Sopra un metodo rapido di colorazione dell' actinomices bovis nelle sezioni. *Morgagni*, (1904), **46**, 270–272.

PRUTZ, W. Die Behandlung der Aktinomykose mit Jodkalium; experimentelle und klinische Untersuchungen. *Mitt. a. d. Grenzgeb. d. Med. u. Chir.*, (1899), **4**, 40–90, 2 pl.

PUSCH, GUSTAV. Beiträge zur Kenntniss der Lungenaktinomykose. *Arch. f. wiss. u. prak. Thierhlknde.*, (1883), **9**, 447–58.

* PUTZU, F. Un caso di actinomicosi temporo-mascellare guarito colle inie-zioni di jodiogelatina sclavo. *Riv. crit. di clin. med.*, (1906), **7**, 765–771.

* QUENET. De l'actinomycose du maxillaire superieur. *Thèse* Lyon, (1895).

DE QUERVAIN. Beitrag zur Aktinomykose des Schädelinnern. *Münchener med. Wchnsch.*, (1899), **46**, 709.

* RABAÏOYE, OCTAVE. Dégénérescence cancroïdale des vielles actinomycoses, Lyon, (1904), 44 pp.

* RABE. Über einen neuentdeckten pathogenen Mikroorganismus beim Hunde. *Berl. thierärzt. Wchnsch.*, (1888).

RADAIS, M. *See* Sauvageau and Radais.

* RAFFA, A. Attinomicosi addominale dell'uomo; saggio di clinica e microscopia. *Riv. ven. sci. med.*, Venezia, (1887), **6**, 577–584.—* Actinomicosi e sua cura. *Riforma med.*, (1892), **8**, 327–329.

* RAINGEARD. Des manifestations cutanée de l'actinomycose. *Thèse*, Paris, (1896).—*See also* Macaigne and Raingeard.

RAMOGNINI, PIETRO. Un caso di actinomicosi della pelle in un dito (patereccio actinomicotico). *Arch. per sci. med.*, (1904), **28**, 245–262.

* RANNEFT. Een geval van actinomycose bij den mensch. *Nederl. Tijdschr. v. Geneesk.*, (1889), II, **25**, 2. d., 633.

* RANSOM, W. B. A case of actinomycosis hominis of the urinary and alimentary tracts; recovery. *Med.-Chir. Trans.*, London, (1891–2), **75**, 85–102, 2 pl; *also abstr.: Proc. Rov. Med. & Chir. Soc.*, London, (1891–2), III, **4**, 14.—* A case of actinomycosis of the vermiform appendix causing perityphlits. *Med.-Chir. Trans.*, (1891–2), **75**, 63–84, 1 pl.; *also abstr.: Proc. Rov. Med. & Chir. Soc.*, London, (1891–2), III, **4**, 12.—The prognosis and treatment of actinomycosis. *Brit. Med. Journ.* (1894), **1**, 61–62.—A case of actinomycosis of the orbit, with a summary of seven cases of actinomycosis. *Ibid*, (1896), **1**, 1553–1555.

* RAPP, G. Trismus ou constriction actinomycosique des mâchoires, Lyon, (1904), 63 pp.

† Rasch, C. Actinomycosis cutis. *Hospitalstidende,* (1898), **6,** 713; *ref.: Jahresb. d. path. Mik.,* (Baumgarten), (1898), **14,** 628–629.

Rasmussen, P. B. Actinomykose. *Deutsche Ztschr. f. Thiermed.,* (1891), **17,** 455–458; (1894), **20,** 299–301.

Ravenal, M. P. *See* Pearson and Ravenal.

* Raybaud, A. Un cas d'actinomycose de la face du maxillaire supérieur. *Marseille méd.,* (1905), **42,** 161–165.—and Sépet, P. Sur un cas d'actinomycose humaine. *Compt. rend. Soc. biol.,* (1903), **55,** 1475–1477.

* Rayevski, P. I. (On the treatment of actinomycosis by iodin compound). *Med. Obozr.,* (1898), **49,** 147–150.

* Raynaud, R. Madura foot (Pied de Madura). *Prat derm.,* (1902), **3,** 448–454.

* Reboul. Un cas d'actinomycose thoracique. *Gaz. hôp.,* (1895), **68,** 1367.—* L'actinomycose et ses dangers pour les travailleurs agricoles. *France agr. et hortic.,* (1900), **18,** 11; *also: Naturaliste,* (1900), II, **22,** 29.

* Reclus, P. Sur une observation d'actinomycose cervico-faciale. *Bull. méd.,* (1906), **20,** 167.

* Redard. Ueber Aktinomykose. *Deutsche Monatschr. f. Zahnheilk.,* (1887).

Reeks, F. H. Successful treatment of actinomycosis. *Journ. Comp. Path. & Therap.,* (1893), **6,** 96.

* Rees. Morphologische und biologische Eigenschaften des Strahlenpilzes der Lungenaktinomykose. *Vereinsbl. d. pfälz. Aerzt.,* (1907), **23,** 3–5.

* Regnier, E. Beitrag zur Casuistik der Abdominal-Actinomycose. *Ztschr. f. Heilk.,* (1894), **15,** 255–264, 1 pl.

Reignier. Un cas d'actinomycose à la Rochelle. *Journ. méd. Paris,* (1905), II, **17,** 509–510.

* Reinsberg, V. Aktinomycosis cutis. *Casop lék. cesk.,* (1906), **45,** 1358.

Remedi, Vittorio. Contributo alle lesioni prodotte dal fungo raggiato. *Sperimentale,* (1903), **57,** 47–73.

* Remy. Trois cas d'actinomycose chez la vache. *Ann. Soc. méd. Gand.,* (1887), **66,** 67–70.—Aktinomykose. *Arch. f. prak. Thierheilk.,* (1898), **24,** 295.—* and Van Ongevalle. Diagnostic de l'actinomycose par l'examen des crachats. *Ann. Soc. méd. Gand.,* (1889), **68,** 272–282.

Report of the Committee of the Pathological Society of London appointed to consider the nomenclature of the conditions sometimes described as Pseudo-Tuberculosis. *Trans. Path. Soc.,* London, (1889), **50,** 361–364.

* Repp, J. J. Actinomycosis in a cow. *Am. Vet. Rev.,* (1903–4), **27,** 341.

† Reynier, Paul, and Brumpt, Émile. Observation parasienne de pied de Madura. *Bull. Acad. méd.* Paris, (1906), III, **55,** 709–723: *also: Bull. et mém. Soc. chir.* Paris, (1906), n. s., **32,** 618–623.

* Rho, F. Micetoma o piede di Madura. *Mal. predom. n. paesi caldi e temperati,* Roma, (1896), **2,** 595–603.

* Richter, Alexander. Zwie Fälle von Aktinomykose als Beitrag zur Kenntniss der Generalisation aktinomykotischer Erkrankungen. *Inaug.-Dissert.,* Kiel, (1901).

* Richter, H. E. Über das Madurabein, eine Pilz-Schmarotzerkrankheit. *Jahresb. d. Ges. f. Nat. u. Heilk.,* (1875–6), **8.**

* Rigal. Un cas d'actinomycose cervicale. *Lyon méd.,* (1903), **100,** 163–165.

* Rigler, Otto. Die Aktinomykose in Thüringen. *Inaug.-Dissert.,* Jena, (1901).

* Von Ritter. Ueber einen Fall von durch eine Streptothrix bedingter ulcerosa mit metastat. Gehirnabscessen. *Prager med. Wchnsch.,* (1900).

* Ritzhaupt, G. O. Ein Fall von Tumorartiger, secundärer Leberaktinomykose, Heidelberg, (1903).

* Rivière, A., and Thévenot, L. L'actinomycose de l'oreille. *Rev. chir.*, (1904), **29**, 42–50.

† Rivière, P. Étude d'un nouveau streptothrix parasite de l'homme. *Arch. clin.* Bordeaux, (1895?), 469; *ref.: Jahresb. d. path. Mik.* (Baumgarten) (1895), **11**, 462.

* Rivolta, S. Sarcoma fibrosa al bordo inferiore della brancha mascellare sinistra del bovo. *Medico vet.*, (1868), 125.—* Del cosi detto farcino o moccio dei bovini e della detta tubercolosi o mal del rospo (Trutta) della lingua dei medesimi animali. *Giorn. anat. fisiol. e. patol. d. animali*, (1875), **7**, 198–217.—† Sul cosi detto mal del rospo della Trutta, e dell'actinomices bovis di Harz, Milano, (1878), 14 pp.; *also: Clinica Vet.*, (1878); 149; *also: Deutsche Ztschr. f. Thiermed.*, (1879), **5**, 110; *ref.: Jahresb. d. gesammt. Med.*, (1878), **1**, 619.—* Sopra un nuovo micromicete del cavallo; nota preventiva. *Guglielmo da Saliceto*, (1879–80), **1**, 145.—Ueber die Priorität der Beschreibung der Formen der Actinomykose und ihrer eigenthümlichen Elemente bei den Rindern. *Arch. f. path. Anat.*, (1882), **88**, 389–390.—* Del micelio e delle varietà e specie di dyscomyceti patogeni. *Giorn. anat. fisiol. e patol.*, (1884), **16**.—* Ancora sulla priorità dell'osservazione dell'actinomyces bovis. *Giorn. anat. fisiol e patol d. animali*, (1887) **19**, 138–144.

* Robin. Pied de Madura. *Gaz. méd.*, (1863), 461.—Note sur le pied endémique de Madura. *Compt. rend. Soc. biol.*, (1864), III, **5**, 34.

* Rochefort. Du pied de Madura ou mycétome de Vandyke Carter. *Arch. méd. nav.*, (1876), **25**.

* Rochet. De l'actinomycose humaine. *Gaz. hebdom. méd.*, (1893), **40**, 149.—* and Martel. Trois cas d'actinomycose cervico-faciale; une forme nouvelle hyperostosante; une forme très atténuee; une forme commune. *Gaz. méd.*, Paris, (1898), XI, **1**, 1–6.

Roecki, G. Ueber Pneumonomykosen. *Ztschr. f. Thiermed.*, (1884), **10**, 122–132, 1 pl.

* Roger. Ueber Aktinomykose. *Aerztl. Int.-Bl.*, (1884), **31**, 583.

* Romanelli, R. Contributo allo studio dell' actinomicosi umana. *Cesalpino*, Arezzo, (1906), **2**, 220–226.

* Romognini, P. Sopra un caso di actinomicosi primitiva della pelle (patereccio actinomicotico). *Gior. d. R. Accad. di med. di Torino*, (1904), IV, **10**, 232–236. *Also: Gazz. d. osp.*, Milano, (1904), **25**, 720.

* Róna, S. Actinomycosis esete. *Gyógyászat*, Budapest, (1899), **39**, 105.

* Rosenbach. Zur Kenntniss der Strahlenpilzerkrankungen beim Menschen. *Centralbl. f. Chirurgie*, (1880).—Über das Erysipeloid. *Arch. f. klin. Chirurgie*, (1887), **36**, 346–350.

* Rosenfeld. Aktinomykose. Centralverband deutscher Aerzte in Böhmen. *Prager med. Wchnsch.*, (1896), *Ref: Hildebrand's Jahresb.*, **2**, 177.

Roser, Karl. Zwei Fälle von acuter Aktinomykose. *Deutsche med. Wchnsch.* (1886), **12**, 369–371.

Rossi, Carmelita. Contributo alla conoscenza dello stipite Actinomices albus varieta tossica. *Ann. ig. sper.*, (1905), n. s., **15**, 693–708; *Ref. Bull. Inst. Pasteur*, (1906), **4**, 121–22.

* Rossi-Doria, E. D. Su di alcune specie di "Streptothrix" trovate nell' aria, studiate in rapporto a quelle già note e specialmente all' Actinomyces, *Ann. Ist. ig. sper. Univ. Roma*, (1891), **1**.

* Rotter. Vorstellung mehrerer Patienten mit Actinomycose, *Verhandl. d. Deutsch. Gesellsch. f. Chir.*, (1886), **15**, 105–112.—Demonstration von Impfaktinomykose. *Ref.: Centralbl. f. Bakt. etc.*, (1888), Abt. 1, **3**, 446.

* Roussel, Georges. De l'actinomycose chez l'homme en France. Deux nouvelles observations, (1891), 73 pp.—*See* Doyen and Roussel.

* Roux. Traité pratique des maladies des pays chauds, (1888), **3**, 353.

* Rovsing, T. Et Tilfaelde af Aktinomykose. *Hosp.-Tid.*, Kjöbenh., (1887), III, **5**, 889–895.—* Om Aktinomykosen hos Mennesket. *Biblioth. f. Laeger*, Kjöbenh., (1889), **9**, 74–107.

Rowland, Penry W. Three cases of aktinomycosis. *Lancet*, (1902), **2**, 671–672.

* Rudneff, S. M. (Case of actinomycosis of neck.) *Chir. Laitop.*, Mosk., (1894), **4**, 585–594.—* On actinomycosis. *Khirurgia*, Mosk., (1897), **1**, 177–201. *Ref.: Centralbl. f. Chir.*, (1897).

* Ruelle, E. Contribution à l'étude du mycétome. *Thése*, Bordeaux, (1893).

Rütimeyer, L. Ein Fall von primärer Lungenactinomycose. *Berl. klin. Wchnsch.*, (1889), **26**, 45–48; 66–68.

Ruffer. Madura foot. (Discussion) *Lancet*, (1893), **1**, 933.

Ruge, Hans. Aktinomycesdrusen. *Deutsche. med. Woch., Ver.-Beil.*, (1896), **22**, 153.—Ueber Actinomyces-ähnliche Gebilde in den Tonsillen. *Zent. f. klin. Med.*, (1896), **30**, 529–37, 1 pl.

* Ruhräh, John. Actinomycosis in man, with especial reference to the cases which have been observed in America. *Ann. Surg.*, (1899), **30**, 417, 605, 722; *Also* (1900), **31**, 235–241.

Ruiz-Casabó, M. Descripcion de un cladothrix cromógeno. *Cron. méd. quirúr.* Habana, (1894), *Ref.: Centralbl. f. Bakt. etc.*, (1895), Abt. 1, **17**, 466.

† Rullmann, Wilhelm. Chemisch-bakteriologische Untersuchungen von Zwischendecken-füllungen mit besonderer Berücksichtigung von Cladothrix odorifera. *Inaug.-Diss.*, München, 46 pp., 1 pl., (1895); *Ref.: Centralbl. f. Bakt. etc.*, (1895), **17**, 884–85.—Ueber eine aus Sputum isolirte pathogene Streptothrix. *Munchen. med. Wchnsch.*, (1898), **45**, 919–922.—und Perutz, Fr. Ueber eine aus Sputum isolirte pathogene Streptothrix. *München. med. Wchnsch.*, (1899), **46**, 407–09; (*Abst.*) *in: Centralbl. f. Bakt. etc.*, (1899), **26**, 152–53.

* Rusloff, I. A. Actinomycosis of the lungs and pleura. *Med. Obozr.*, Mosk., (1899), **51**, 706–713.

Russell, H. L. Untersuchung über im Golf von Neapel lebende Bacterien. *Zeit. f. Hyg. etc.*, (1892), **11**, 165–204, 2 pl.

* Rustomjee, Bazunjee. Cases of diseased foot. *Tr. Med. and Phys. Soc.*, Bombay, (1860), n. s., **5**, 230–241.

Rydygier. Zur Therapie der Aktinomykosis. *Wien. klin. Wchsch.*, (1895), **8**, 649–650.

* Sabrazès, J. Les parasites du genre streptothrix dans la pathologie humaine. *Mercredi med.*, Paris, (1895), 458.—*and Cabannes, C. Actinomycose pulmonaire. *Rev. méd.*, (1899), **19**, 68–77.—Die Aktinomykose der Lungen. *München. med. Wchnsch.*, (1899), **46**, 425.—* and Rivière. Sur une streptothrix rencontré dans un cas d'abscès du cerveau et à un infarctus suppuré du rein. *Presse méd.*, (1894).—* Les parasites du genre streptothrix dans la pathologie humaine. *Deuxième Congrès francaise de médecine interne*, (1895). *Also: Sem. méd.*, (1895); *Mercredi méd.*, (1895), 485–486.—*See also* Dubreuilh et Sabrazès.

Saccardo, P. A. Streptothrix. *Sylloge Fungorum*, (1886), **4**, 282–283.

* Salinari, S. Contributo clinico sperimentale allo studio dell' actinomicosi umana. *Gior. med. esercito*, (1904), **52**, 641–656.

* Salmon, D. E. Report upon investigations relating to the treatment of lumpy-jaw, or actinomycosis in cattle. *Rep. U. S. Bureau Animal Ind.*, (1891–2), **8–9**, 109–176, 8 pl; (1893–4), **10–11**, 88–91.—Actinomycosis, or lumpy jaw. *Circular U. S. Bur. An. Ind.*, no. 7, (1896), 4 pp.—and Smith, Theobald.

Infectious diseases of cattle. *In:* Special report on diseases of cattle (U. S. Bur. An. Ind.), rev. ed. (1904). Actinomycosis. pp. 427–436, 3 pl. *Also Circular U. S. Bur. An. Ind.*, no. 96, (1906), 10 pp., 3 pl.

SAMES, THEODOR. Zur Kenntniss der bei höherer temperatur wachsenden Bakterien-und Streptothrixarten. *Ztschr. f. Hyg.*, (1900), **33**, 313–362, 1 pl.

* DE SAMPAIO BARROS, F., *jr.* Zur Pathologie und Therapie der Actinomycose beim Thiere und beim Menschen. Aachen, (1883), 27 pp.

SAMTER, E. O. Ein Beitrag zu der Lehre von der Aktinomykose. *Arch. f. klin. Chir.*, (1892), **43**, 257–351, 1 pl.

SANFELICE, FRANCESCO. Beiträge zur Kenntniss der Aktinomykose der Leber bei den Rindern. *Arch. f. wiss. u. prak. Thierhlknde.*, (1896), **22**, 153–170, 2 pl. —Delli' azione patogena di alcune streptothrix. *Centralbl. f. Bakt. etc.*, (1904).—* Tuberculosi e pseudotuberculosi. *Riforma med.*, (1904).—Ueber die pathogene Wirkung einiger Streptothrix-(Actinomyces) Arten. *Centralbl. f. Bakt.*, etc. (1904), Orig., **36**, 355–367.—Streptothrix-pseudotuberkulose. *Ibid*, (1905), Orig., **38**, 30–41, 1 pl.

* SANO, P. À propos d'actinomycose. *Journ. méd. de Brux.* (1905), **10**, 209.

SATA, A. Ueber die Fettbildung durch verschiedene Bakterien nebst einer neuen Färbung des Aktinomyces im Schnitte. *Centralbl. f. allg. Path. u. path. Anat.* (1900), **11**, 97–102.

* SATTERTHWAITE, T. E. Actinomycosis in man and animals. *Quart. Bull. Clin. Soc.* (1885–86), **1**, 160–163.

SAUVAGEAU, C., and RADAIS, M. Sur les genres cladothrix, streptothrix, actinomyces et description de deux streptothrix nouveaux. *Ann. d. l'Inst. Pasteur*, (1892), **6**, 242–273.

* SAWTSCHENKO, J. G. (Pseudo-actinomycosis produced by a bacillus; a new bacterium which produces suppuration clinically similar to actinomycosis). *Russk. Arch. patol. klin. Med. i Bakteriol.* (1896), **1**, 184–203, 1 pl. *Also transl. abstr.: Ibid.* 209–211. *Ref. Jahresb. pathogenen Microorganismen (Baumgarten)*, (1896), **12**, 613.

SAWYERS, J. L. Medical treatment of actinomycosis. *Journ. Am. Med. Ass.* (1901), **36**, 1314–19.

SCAFFIDI, V. Di un actinomyces che si trova nella bocca dell' uomo (actinomyces lingualis non liquefaciens) *Giorn. internaz. d. sci. med.* (1907), n. s. **29**, 409–415.

SCHABAD, J. A. Actinomycosis atypica pseudotuberculosa; streptothricosis hominis autorum. *Russk. Vratch*, (1903) **2**, 1242–45; 1282–85; 1317–21. *Also: Ztsch. f. Hyg.* etc., (1904), **47**, 41–80, 1 pl.

* SCHARTAU, H. F. Ein Beitrag zur Kenntniss der Aktinomykose, Kiel (1890), 43 pp.

† SCHEELE and PETRUSCHKY. Kulturen und Präparate einer menschpathogenen Streptothrixart (Diagnose in vivo). *Verhandl. des Kongresses f. inn. Med.* (1897), **15**, 550. *Also, ref.: Münch. med. Woch.* (1897), **44**, 686; *Deutsch. med. Woch.* (1897), **23**, *Ver.-Beil.*, 124.

† SCHEUBE, B. Madurafuss. *In: Eulenburg's encykl. Jahrb.* (1897), **7**, 243. Also: *In his:* Die Krankheiten der warmen Länder (1900), 2. ed., 618–631; (1903), 3. ed., 735–748; and Engl. translation by Pauline Falcke, edited by James Cantlie, London, (1903), 2. rev. ed., 552–563.

* SCHILLER, H. Ueber Aktinomykose des Menschen, ein Beitrag zur Pathogenese derselben, Giessen (1892).

* SCHILLING. Zungenaktinomykose beim Schweine. *Ztschr. f. Fleisch- u. Milchhyg.*, **10**, 134.

* SCHIRMER, A. A case of actinomycosis hominis. *Chicago Med. Journ. & Exam.* (1886), **53**, 354.

SCHLAGENHAUFER. Beiträge zur pathologischen Anatomie der Aktinomykose beim Menschen. *Virchow's Arch.* (1906), **184**, 491–515, 1 pl.

† SCHLANGE, H. Zur Prognose der Aktinomykose. *Arch. f. klin. Chir.*, (1892), **44**, 863–875; *also: Centralbl. f. Chir.*, (1892), 37; *also: Berl. klin. Wchnsch.*, (1892), **29**, 706–707; *also: Verhandl. d. deutschen Ges. f. Chir.*, (1892), **21**, pt. 2, 241–253; Discussion, pt. 1, 45.

SCHLEGEL, M. Aktinomykose des Menschen und der Thiere. *Ergeb. d. allg. Path.*, (Lubarsch-Ostertag), (1898), **5**, 404–446.—Zur Lungenaktinomykose. *Berl. tierärzt. Wchnsch.*, (1903), 409–411.

SCHMALTZ. Jodwirkung bei Actinomyces. *Ibid*, (1902), 401–402.

SCHMORL, GEORG. Über ein pathogenes Fadenbacterium (Streptothrix cuniculi). *Deutsche Ztschr. f. Thiermed.*, (1891), **17**, 375–408, 2 pl.; *also ref.: Hyg. Rundschau*, (1891), **1**, 882–883.

SCHNEIDEMÜHL, GEORG. Ueber Strahlenpilzerkrankungen bei Mensch und Tier. *Münchener med. Wchnsch*, (1890), **37**, 639–641.

* SCHNEIDER, I. G. (Actinomycose du boeuf; cas observé à l'abattoir municipal d'Ekathérinbourg.). *Zapiski Uralsk. med. Obsh.*, (1892), **2**, 89–94.

* SCHOU, J. (A case of abdominal actinomycosis cured by iodide of potassium.). *Ugesk. f. Laeger*, Kjöbenh., (1895), V, **2**, 1174–1178.

* SCHOUTEN, S. Een geval van actinomycosis. *Nederl. Tijdschr. v. Geneesk.*, (1902), II, **38**, 254–256.—* Bijrdräge tot de kennis van het voorhomen en de verspreiding der actinomycosis bij de huisdieren in Nederland. *Tijdschr. v. Veeartsenijk. Maandbl.*, (1903–4), **31**, 97–153.

* SCHREYER, HERMANN. Zwei Fälle von Actinomykose der Bauchdecken, Greifswald, (1890), 28 pp.

* SCHROEDER, F. Luchist. gribok (Aktinomykosis) v nizhem sleznom kanal. (Actinomycosis in inferior lachrymal canal). *Vratch*, (1894), **15**, 291; 351.

* SCHROETER, J. Pilze schles., 173.

* SCHRÖTTER, L. Ueber aktinomykose des Mediastinums und der Herzens. *Int. Beitr. z. innere Med.*, (1902), **1**, 535.

SCHTSCHEGLOW. Ein Fall von Lungen-Pseudoaktinomykose. *Ref.: Ergeb. d. allg. Path.* (Lubarsch-Ostertag), (1898), **5**, 665.

* SCHUCHARDT, B. Ueber das Vorkommen des Strahelnpilzes (Actinomyces) und der Strahlenkrankheit (Actinomycosis). *Cor.-Bl. d. allg. ärztl, Ver. v. Thüringen*, (1888), **17**, 323–337.—* Ueber das Vorkommen von Actinomycose beim Epephanten in Indian und über die Identität des Madura-fuses mit Actinomycose. *Cor.-Bl. d. allg. ärztl. Ver. v. Thüringen.* (1888), **17**, 376.

SCHÜRMAYER, BRUNO. Ueber Aktinomykose des Menschen und der Tiere. *Centralbl. f. Bakt. etc.*, (1900), Abt. I., **27**, 49–61; 101–107, 2 pl.

† SCHUKEWITSCH, J. J. Zur Lehre über die Actinomycose des Rindes. *Wiss. Abhandl. d. Kasan'schen Vet.-Inst.*, 193–230; 231–335; *ref.: Jahresb. d. gesammt. Med.*, (1902), **1**, 711–712.

SCHULZE. Behandlung der Actinomycome. *Berl. thierërzt. Wchnsch.*, (1899), 52–53.—Untersuchungen über Strahlenpilzformen des Tuberculoseerregers. *Ztschr. f. Hyg.*, **31**, 153–186, 1 pl.

* SCHWARTZ, E. Actinomycose de la face; traitement et guérison par l'iodure de potassium. *Bull. et mém. Soc. chir.* Paris, (1904), n. s., **30**, 948.

SCHWARZ. Actinomykose beim Pferd. *Berl. thierärzt. Wchnsch.*, (1901) 600.

* SCHWARZENBACK. Beiträge zur Pathologie und Therapie der Streptomykosen. *Cor.-Bl. f. schweiz. Aerzte*, (1905), **35**, 129; 172; 220.—* Die interne Behandlung der Streptomykosen mit Formaldehyd. *Ibid*, 766–777.

* SCOTT, J. Actinomycosis. *Am. Vet. Rev.*, (1890–1), **14**, 7–13.

SEMMER, E. Ueber die Morphologie des Tuberkel-und Rotzbacillus und der pathogenen Schizomyceten. *Ztschr. f. Thiermed.*, (1895), **21**, 212–216.

SENN, NICHOLAS. *See* Hyde, Senn, and Bishop.

SÉPET, P. *See* Raybaud and Sépet.

* SERGI TROMBETTA, D. Il primo caso di actinomicosi in Messina. *Morgagni*, (1894), **36**, 119.

* SHAH, T. M. Mycetoma; variaties; its clinical aspects; with cases. *Med. Rep.*, Calcutta, (1893), **2**, 225–227.

* SHATTOCK, S. G. Two cases of actinomycosis in man. *Trans. Path. Soc.* London, (1884–5), **36**, 254–261, 1 pl.—† Mycetoma papillomatosum. *Ibid*, (1897–8), **49**, 293–296, 1 pl.; *also: Brit. Med. Journ.*, (1898), **1**, 622.

* SHTSHEGLOFF, M. Pseudoactinomycosis of the lungs. *Med. Obozr.*, (1897), **48**, 649–660; *also transl.: Compt. rend. Cong. int. med.*, (1897), **2**, sec. 3, 284–296.

† Siedamgrotzky. Epulis vom Rinde mit Aktinomyces bovis. *Ber. ü. d. Veterinärw. im Kgr. Sachsen* (1877); ref: *Beitr. z. path. Anat.*, (1891), **9**, 180; *Jahresb. d. gesammt. Med.*, (1878), **1**, 618–619.

SILBERSCHMIDT, W. Sur un nouveau Streptothrix pathogène (Streptothrix caprae). *Ann. Inst. Pasteur*, (1899), **13**, 841–853.—Ueber zwei Fälle von Pilzmassen im unteren Thränenkanälchen. *Centralbl. f. Bakt. etc.*, (1900), **27**, 486–493, 1 pl.—Ueber Aktinomykose. *Ztschr. f. Hyg. u. Infekt.*, (1901), **37**, 345–380, 2 pl.—Zur bakteriologischen Diagnose der Aktinomykose. *Deutsche med. Wochnsch.*, (1901), **27**, 816–817.

SILBERSTERN, P. Hautemphysem und Lungenaktinomykose. *Wien. med. Woch.*, (1893) **43**, 1843–46; 1886–90; 1928–30.

SKERRITT, E. M. Actinomycosis hominis. *Am. Journ. Med. Sci.* (1887), n. s. **93**, 75–88.

SMITH, C. B. M. A case of actinomycosis treated by iodide of potassium. *Lancet* (1897), **1**, 734–35.

* SMITH, F. C. Case of mycetoma. *Trans. South Indian Branch Brit. Med. Ass.* Madras (1888), **2**, 287–289, 2 pl.

SMITH, G., HARSANT, W. H., and BUSH, J. P. Three cases of actinomycosis. *Lancet* (1897), **1**, 311.

* SMITH, H. B. W. Actinomycosis and notes on a case. *Transvaal Med. Journ.* (1907–08), **3**, 23–25.

† SMITH, T. (Note on a drawing made about 1855). Published by Kanthack in *St. Barth. Hosp. Journ.* (1896), 51. Ref.: Allbutt's System of Medicine (1906), **2**, pt. 1, 325.

* SMOLITSCH. Mastite streptococcique des vaches ou mastite contagieuse ou chronique des vaches laitieres de Nocard et Mollereau. *Travaux du 1. Congress vétérinaire russe*, St. Petersb. (1903), **1**, 478; 480.

* SMYTH, J. Notes on a case of mycetoma of the neck. *Indian Med. Gaz.* (1898), **33**, 56.

SNOW, HERBERT. Case of actinomycosis with tuberculosis. *Brit. Med. Journ.* (1891), **2**, 124–25.

* SÖDERBAUM, P. Fall af actinomycosis. *Upsala Läkaref. Forh.* (1893), **28**, 227–230.

* SOKOLOFF, K. V. (Actinomycosis of lungs recognized during life from sputa) *Voyenno-san. Dielo*, St. Petersb., (1889), **9**, 9.

* SOKOLOFF, N. A. (Prevalence of actinomycosis in Russia). *Vratch*, St. Petersb. (1893), **14**, 384; 420; 454.

* SOLTMANN. Zur Aetiologie d. Aktinomykose. *Bresl. Aerzteztschr.* (1885).—* Ueber Aetiologie und Ausbreitungsbezirk der Actinomycose. *Jahrb. f. Kinderheilk.* (1886), n. s., **24**, 129–139.

* von Sommer, G. Primo caso di actinomycosi osservato in Napoli. *Riv. internaz. di med. e. chir.* (1886), **3**, 9–14; (1887), **4**, 90–101. *Also: Boll. di clin.* (1887), **4**, 393–400; *also transl., abstr. Union med.* (1887), III, 44, 686–688.

* Sommer y Greco. Primer caso de mycetoma ó pie de Madura en la República Argentina. *Argent. Med.* (1904).

* Sonnenburg. Ein Fall von Lungenaktinomykose. *Berl. klin. Woch.* (1890).

Soubrane, L. *See* Fortineau and Soubrane.

† Soucail. Erfolgreiche Behandlung dreier Fälle von Aktinomyces Geschülsten des Kiefers durch grosse gaben von Jodkalium. *Rev. vet.*, (1893), 65, *Ref.: Jahresb. gesammt. Med.*, (1893), **1**, 651.

Spitz, G. *See* Lignières and Spitz.

* Spizharnîy, I. K. Actinomycosis of the lungs. *Med. Obozr.*, (1903), **59**, 85–93. *Also, transl.: Rev. balear de cien. med.*, Palma de Mallorca, (1903), **19**, 36–40. *Habana med.*, (1903), **6**, 57–59.

* Stanton, E. McD. Actinomycosis limited to the urinary tract. *Albany M. Ann.*, (1905), **26**, 738–742.

* Starkiewicz, W. Actinomycosis of the lungs and pleura. *Gaz. lek.*, Warszawa, (1902), 2. s., **22**, 949–954.

* Staub. Zur Therapie der Hautaktinomykose. *Therap. Monatsh.* (1894).

* Sterling, S. (Two cases of actinomycosis of the lungs.) *Czasopismo lek.* (1903), **5**, 327 to 334.

Sternberg, Carl. Zur Kenntniss des Aktinomycespilzes. *Wien. klin. Wchnsch.* (1900), **13**, 548 to 551.

* Stewart, G., and Muir, R. Notes on a case of actinomycosis of ovaries and liver. *Edinb. Hosp. Rep.* (1893), **1**, 96 to 113, 1 pl.

† Stokes, W. R. A study of the group actinomyces; with the report of a pathogenic species for man. *Am. J. M. Sc.*, Phila. & New York (1904), n. s., **128**, 861 to 875; *Also J. Alumni Ass. Coll. Phys. & Surg.*, Balt. (1907), **10**, 40 to 52.

* Stolpe. Ueber Aktinomykose der Lymphdrusen bei amerikanischen Rindern. *Ztschr. f. Fleisch. u. Milchhyg.* (1907), **17**, 339.

Stolz, Albert. Ueber einen Bacillus mit Verzweigungen: *Arch. f. Hyg.* (1897), **30**, 156 to 167.

* Stooss, M. Zur Aetiologie und Pathologie der Anginen. *Mitteilungen aus Kliniken u. med. Instituten der Schweiz* (1895).

* Strebel, M. Die Jodebehandlung der Aktinomykose beim Rinde, *Schweiz.-Arch. f. Tierh.* Zürich (1898), **40**, 49 to 59.

* Streiten, van der. Actinomycose du foie. *Bull. d. l'Acad. royale d. méd. d. Belgique* (1891), 549; *Ref.: in* Bécue, De l'actinomycose, *Thèse*, Paris (1892).

* Struve. Ein Fall von Aktinomykose beim Pferde. *Ztsch. f. Fleisch.-u. Milchhyg.* (1897), **3**, 29.

* Subbotic. Ein Beitrag zur Kenntniss der Verbreitung der Aktinomykosis. *Pester med. chir. Presse* (1886).

Surveyor, N. F. Madura foot of India. *Brit. Med. Journ.*, Lond. (1892), **2**, 575 to 576.—* Madura foot of India. *Rep. and Proc. Royal Soc.* (1893); *also* (Review) *Centrbl. f. Bacteriol. etc.* (1894), Abth. I, **16**, 798 to 799.—*See also* Boyce and Surveyor.

* Sykoff, V. (Actinomycosis of pleura and lungs.) *Russk. Med.* (1890), **16**, 767.

* Syms, P. Actinomycosis. *Ann. Surg.* (1897), **25**, 155 to 171.

* Szegedy, J. Hasi actinomycosis esete. (Case.) *Orvosi hetil.*, Budapest (1907), **51**, 502.

* Szénásy, S. Sugárgomba-betegség (aktinomycosis). *Gyógyászat*, Budapest (1886), **26**, 625 to 630; *also, transl. Pest. med.-chir. Presse*, Budapest (1886), **22**, 908; *Centrbl. f. Chir.* (1886).

* Taburet, J.- H.- E.- M. Contribution à l'étude clinique de l'actinomycose cutanée chez l'homme. Bordeaux, (1893).

* Takasu, K. Ein Fall der Lungenactinomycose. (Japanese text.) *Ztschr. d. med. Gesellsch. zu Tokyo* (1897), **11**, 911 to 916.

† Tansini. Behandlung der Aktinomykose. *Münchener med. Wchnsch.* (1900), **47**, 954.—Sopra un caso di actinomicose. *Clin mod.*, Pisa (1900), **6**, 90 to 91.

* Taufal. Aktinomykose des Mittelohres. *Prag. med. Wchnsch.* (1894).

* Van Tavel. Vergleichende Morphologie der Pilze (1892).

* Taylor, F. A case of actinomycosis of the liver and lungs. *Guy's Hosp. Rep.* (1891), III., **33**, 311 to 323.

Terni, C. Una nuova specie di actinomyces (actinomyces Gruberii). *Atti d. XI Cong. med. internaz.* (1894), **6**, igiene, 79 to 82; also *Centrbl. f. Bakt. etc.* (1894), **16**, 362 to 363.—Actinomycosi della lacertola, Actinomyces lacertae. *L'Ufficiale sanitario.* (1896), 160; Ref.; *Centrbl. Bakt.* (1896), **19**, 953.

* Terrier, F., and Dujarier, C. Un cas d'actinomycose cervico-crânienne. *Rev. d. chir.* (1906), **33**, 431 to 433.

* Thebault, Léon. Contribution à l'étude clinique de l'actinomycose; l'actinomycose à Rennes (1906), 63.

Thévenot, L. L'actinomycose du poumon et ses manifestations primordiales. *Arch. gén d. méd.* (1903), **2**, 3137 to 3145.—Panaris actinomycosique de l'annulaire droit. *Presse méd.*, Par. (1903), **11**, 659 to 660.—Actinomycose cervico-faciale de forme courante. *Gaz. d. hôp.* (1903), **76**, 769 to 771.—* Actinomycose de la fesse d'origine rectale. *Lyon méd.* (1903), **100**, 123 to 125.—* Actinomycose de la région de l'angle maxillaire. *Ibid.* (1903), **101**, 180.—* Actinomycose cervico-faciale; nouvelles observations. *Arch prov. d. chir.* (1904) **13**, 98 to 105.—* De l'actinomycose humaine dans le Dauphiné. *Dauphiné méd.*, Grenoble (1904), **28**, 213.—* Actinomycose de l'amygdale, phlegmon cervical consécutir. *Lyon méd.*, 12 juin (1904).—À propos de l'actinomycose humaine (1907), **108**, 942 to 945.

* Thiriar. Un cas d'actinomycose. *Clinique* (1891), **5**, 417 to 425; also: *Mercredi méd.* (1891), **2**, 349 to 352; *J. d. mal. cutan. et syph.* Par. (1891), **3**, 498 to 507; *Gaz. d. hôp. de Toulouse* (1891), **5**, 323; 331; transl.: *Gazz. d. osp.* Napoli (1891), **12**, 623 to 626.

* Thirion, G., and Augier, D. Actinomycose pleuropulmonaire propagée a là paroi abdominothoracique et au foie, prise pour un abcès du foie; cachexie; opération; mort neuf jours après. *J. d. sc. méd. de Lille* (1903), **2**, 121 to 132.

* Thiroux. Mycose de la jambe due au Streptothrix madurae (Vincent). *Ann. d'hyg. et d. méd. colon.* (1906), **9**, 453 to 455.

* Thiry, Georges. Contribution à l'étude du polychromisme bacterien; bacille et cladothrix polychromes cristaux colorés.—*Arch. de phys.*, (1897), **29**, 284–88. Un actinomyces mordoré et son pigment cristallisé. *Compt. rend. d. Cong. d. Soc. sav. d. Paris, Sect. d. Sc.* (1901), 64 to 66.

* Thollon. Du sarcome actinomycosique. *Thèse*, Lyon (1896).

* Tichow, P. J., Rasowoski, W. J., Stephanowski, T. K. Ueber Aktinomykose. *VII Kongr. d. russisch. Aerzte in Kasan* (1899); *Ref.: Centrbl. f. Chirurgie.* (1899).

* Van Tieghem. Traité de botanique (1891), 2. ed., 1069; 1191; 1207.

Tilanus, C. B. Ueber einen Fall von Actinomykosis cutis faciei. *Münchener med. Wchnsch.* (1889), **36**, 534 to 535.—* Twee gevallen von actinomycosis cutis faciei. *Nederl. Tijdschr. v. Geneesk.* (1889), 2. s. **25**, 517 to 520.

* Tizzoni, G. Ancora sull' adenomicosi septica; nuove ricerche. *Arch. per le sc. med.* (1881), **5**, 87 to 96.

† Toggia (1822). Cited by Rivolta *in his* Sul cosi dette mal del rospo. etc. *Clinica, vet., Rev.: Jahresb. d. ges. Med.* (1878), **1**, 619.

* Tommasoli, P. L. L'actinomicosi dal punto di vista dermatologico e nei suoi rapporti con altre dermatosi congeneri come il micetoma, la micosi fungoide, etc., *Gior. ital. d. mal. ven.* (1893), **28**, 409; 513.

De Toni, J. B., and Trevisan, V. Schizomycetaceae. *In:* Saccardo's Sylloge Fungorum (1899), **8**, 923 to 1087. (Sphaerotilus, 926; Cladothrix, 927; Nocardia, 927 to 928.)

Toussaint, M. Contribución al estudio de la actinomicosis. *Bol. d. Inst. patol.*, México (1907–8), II., **5**, 181; 241; 299; 371; 445.

† Treves, F. A case of supposed actinomycosis. *Tr. Path. Soc.* (1883–4), **35**, 356 to 362, 1 pl.; *also: Lancet* (1884), **1**, 70; 848.

Treves, W. Knight. On a case of actinomycosis. *Lancet* (1884), **1**, 107 to 108.

Trevithick, Edgar. A case of actinomycosis; primary abscess of the left lung exhibiting spontaneous cure; metastatic manifestation in the pelvis declaring itself after a year's latency with abscess formation and slow spontaneous cure. *Lancet* (1906), **2**, 158 to 159.

† Trinchera. *Clin. vet.* (1893), **16**, 485; *Jahresb. f. ges. Med.* (1893), **1**, 652.

Trolldenier. Ueber eine bei einem Hunde gefundene pathogene Streptothrix. *Ztschr. f. Thiermed.* (1903), n. s., **7**, 81 to 109. *Ref.: Centrbl. f. Bakt. etc.*, (1904), Abt. 1., *Ref.:* **34**, 124 to 125.

Truelsen, O. Zungenactinomycom des Pferdes. *Berl. thierärztl. Wchnsch.* (1893), 39 to 40; *Ref.: Jahresb. d. ges. Med.* (1894), **1**, 6 to 61.

† Trutta (1785). Cited by Rivolta *in his* Sul cosi detto mal del rospo, etc. *Clinica vet.: Rev.: Jahresb. d. ges. Med.* (1878), **1**, 619.

Tsilinsky, P. Sur les mucédinées thermophiles. *Ann. Inst. Pasteur* (1899), **13**, 500 to 505.

Tsiperkusa, T. V. Sluchai luchisto-gribkovoi lablizni. *Russk. Vratch.* S.-Petersb., (1906), n. s. **5**, 295 to 296.

† Tullberg. Lokale Behandlung der Actinomykose mit Arseniksalbe. *Svensk. Veterinartidskrift* (1896), **1**, 252; *Ref.: Jahresbr. ges. Med.* (1896), **1**, 517.

* Turner, G. I. (On the treatment of actinomycosis). *Vratch.* (1895), **16**, 1174–1177.

Tusini, Giuseppe. Über die Aktinomykose des Fusses. *Arch. f. klin. Chir.*, (1900), **62**, 249–287, 5 pl.

* Tuttle, G. A. A case of general streptothrix infection; with a study of the micro-organism. *Med. & Surg. Rep. Presbyterian Hosp.*, N. Y., (1904), **6**, 147–169.

* Ullmann, E. (Fall von Bauchactinomycosis). *Anz. d. k. k. Gesellsch. d. Aerzte in Wien*, (1887), 183.—* Beitrag zur Lehre von der Actinomycose. *Wien. med. Presse*, (1888), **29**, 1769; 1812; 1853.

Unna, P. G. Die Histopathologie der Hautkrankheiten Mycetoma, Madurafuss. *In:* Orth's Lehrbuch der speciellen pathologischen Anatomie Berl., (1894), Ergänzungsband, **2**, 469–472.—* Ueber Piedra nostras. *Deutsche med. Ztg.*, (1895), **16**, 255.—Ueber Aktinomykose und Madurafuss. *Münch. med. Wchsnch.*, (1897), **44**, 150.—* Aktinomykose und Madurafuss. *Deutsche med. Ztg.*, (1897), **18**, 49–51; *also: Arb. a. Unna's Klin. f. Hautkr. in Hamb.*, Berl., (1898), 1–9.—* and Delbanco, Ernst. Beiträge zur Anatomie des indischen Madurafuss (Mycetoma, Fungus disease of India). *Monatsh. f. prakt. Dermat.*, (1900), **31**, 545–567, 2 pl.

* UNTH. Ein Fall v. Aktinomykose d. Auges. *Centralbl. f. prakt. Augenheik.*, (1894).

URBAN. Aktinomykose. *Münch. med. Wchsnch.*, (1897), **44**, 124.

* URIZER, R. Dos nuevos streptotrix (*Streptotrix lignièri*). *Rev. Soc. méd. argent.* (1904), **12**, 570 to 578.

* VACHETTA. Studi e ricordi clin. Milano (1882). *Ref.: Oesterr. Vierteljahrsschr.*, Wien (1882), 37.

* VALERY, C. Sur un cas d'actinomycose du poignet. *Gaz. d. hôp.* (1906), **79**, 606.

* VALLE, L. Actinomicosis cervicofacial. *Rev. Soc. méd. argent.* (1903), **11**, 513 to 539, 1 pl.

VALLÉE, H. Sur un nouveau streptothrix (streptothrix polychromogène), *Ann. Inst. Pasteur* (1903), **17**, 288 to 292.

† VEER, A. VAN DER, and ELTING, A. W. A résumé of the subject of actinomycosis, with report of a case of actinomycosis abdominalis. *Med. News* (1902), **80**, 109 to 113. *Also: Albany Med. Ann.* (1902), **23**.

* VENNERHOLM. Aktinomykos hos vara husdjur. *Svensk Veterinärtidsskrift* (1896), **1**, 22; 33; 81.

* VERHAEGHE, É. L'actinomycose dans la région du Nord. *Écho méd. du nord* (1907), **11**, 445; 457.

* VERLIAC, HENRY. Recherches experimentales sur less toxins de l'actinomyces. Contribution à l'étude des poisons microbiens à l'action locale prédominante (1907), 83 pp.

† VEROCAY, J. Aktinomykose der Beckenorgane eines 14-jährigen Mädchens. *Verhandl. d. deutsch. path. Gesellsch.* (1905), 139 to 142; *Also abstr.: Centrbl. f. allg. path.* (1905); **16**, 808 to 809.

VIGNAL, W. Recherches sur les microörganismes de la bouches. *Arch. d. physiologie normale patholog*, (1886), **18**, II, 325 to 391, 8 pl.

* VINCENT, E., and SAULIÉ, H. Quelques cas de maladies à streptothrix chez l'homme en Algérie. *Bull. méd. d. l'Algérie*, (1905), **16**, 101 to 109.

VINCENT, H. Étude sur le parasite du "pied de Madura." *Ann. Inst. Pasteur*, (1894), **8**, 129 to 151, 3 fig.—* Sur l'unicité du parasite de la maladie de Madura (Streptothrix Madurae H. Vincent) et sur ses formes génératives. *Compt. rend. Soc. d. biol.* (1906), **61**, 153 to 155. *Also: Caducée* (1906), **6**. 245.—*See also* Gémy and Vincent.

* VINOGRADOV, K. N. (Actinomycosis in man.) *Russk. Méd.* (1885), **3**, 875 to 877, (1886), **4**, 3; 27.

VIRCHOW, RUD. Beiträge zur Lehre von den biem Menschen vorkommenden pflanzlichen Parasiten. *Virch. Arch.* (1856), **9**, 557 to 593, 1 pl.—Beiträge zur Kenntniss der Trichinosis und der Actinomycosis bei Schweinen. *Ibid* (1884), **95**, 534 to 547.

* VISCONTI, A. Primo caso dittinomicosi polmonale nell'uomo osservato in Milano. *Reudic. R. Ist. Lomb. di sc. e lett.* (1898), 2. s., **31**, 615 to 630. *Also, in his:* Microscopie, Milano (1898), 11 to 26.

* VITA, D. DE. Il primo caso di actinomicosi in militari italiani. *Ann. di med. nav.* (1901), **7**, 653.

VITRAC. *See* Capdepon and Vitrac.

* VLAYEFF, G. M. (Case of actinomycosis; on the morphology and biology of the exciting agent of this disease.) *Voyenno-méd. J.*, St. Petersb. (1897), **188**, 1158 to 1176.

* VOLKOVITCH, N. M. (Some remarks on actinomycosis in man, - its distinction, pathology, and propagation.) *Chir. Laitop.*, Mosk. (1893), **3**, 3 to 22, 1 pl.

* VORONOFF, S. L'actinomycose en Égypte. *Gaz. d. hôp.* (1904), **77**, 1301 to 1303.

Voss, J. Die Behandlung des Strahlkrebses mit grüner Seife. *Berl. thierärztl. Wchnsch.* (1904), 592 to 593.

* Walker, J. E. Actinomycosis. *Memphis Lancet* (1898), **1**, 279 to 287.

† Walley. The abortive treatment of actinomycosis. *Veterin.* (1886), **59**, 53; *Ref.: Jahr. d. gesam. Med.* (1886), **1**, 550.

Wallhauser, H. J. F. A case of actinomycosis. *Journ Cutan. Dis.* (1904), **22**, 77 to 79, 1 pl.

† Walther. Jodtherapie bei Actinomycose. *Bericht über das Veterinärwesen im Königreich Sachsen* (1894), 127; 128. *Ref.: Jahresb. f. ges. Med.* (1894), **1**, 661 to 662.

* Waring, H. J. A case of actinomycosis hominis. *St. Barth. Hosp. Rep.*, London (1891), **27**, 173 to 175.—* Actinomycosis of the caecum, vermiform appendix, and right iliac fossa, including an account of seven cases. *Ibid* (1905), **41**, 197 to 210.

* Warren, G. A. A case of actinomycosis. *Hot Spring, Med. Journ.* (1903), **12**, 100 to 102.

Warthin, A. S., and Olney, H. S. Pulmonary streptothricosis; notes of a case with clinical picture of pulmonary tuberculosis; sputum containing an acid-resisting streptothrix; no tubercle bacilli. *Am. Journ. Med. Sci.* (1904), n. s., **128**, 637 to 649.

Webber, H. W. Further notes on a case of abdominal actinomycosis. *Brit. Med. Journ.* (1903), **1**, 1084.

* Weber, L. Ein Fall von Actinomycosis. *Allg. Wien. med. Ztg.* (1901), **46**, 210. *Also N. Yorker med. Monatschr.* (1901), **13**, 115 to 118.

* Wedemeyer, Adolf. Beiträge zur Kenntniss der Aktinomykose des Menschen, *Inaug.-Diss.* Göttingen (1897).

* Weiss. Un cas d'actinomycose. *Rev. méd. l'est.* (1896), **28**, 262 to 267.—* Gross and Rhomer. Actinomycose cervico-faciale. *Gazette heb.* (1896), 537.

Welch, W. H. *See* Latimer and Welch.

Welleminsky. Ueber atypische Befunde bei Perlsucht und Aktinomykose des Rindes. *Deutsche med. Wchnsch.* (1902), **28**, Ver.-Beil., 196.

* Werner. Ein Fall von Actinomycosis. *Med. Cor.-Bl. d. württemb. ärztl. Ver.* (1889), **59**, 93.

* Wettergren, C. Actinomycosis regionis ileocaecalis. *Eira*, Stockholm (1903), **28**, 199 to 210.

* Whitney, W. F. A case of actinomycosis in a heifer. *Boston Med. & Sur. Journ.* (1884), **110**, 532.

* Wildermuth. Ein Fall von Actinomycosis. *Med. Cor.-Bl. d. württemb. ärztl. Ver.* (1886), **56**, 9.

* Williams, W. L. Actinomycosis in relation to meat inspection. *J. Comp. Med. & Vet. Arch.* (1892), **13**, 351 to 355.

Williamson, G. A. Interesting case of mycetoma in Cyprus. *Journ. Trop. Med.* (1905), **8**, 81 to 82.

* Williger, F. Ueber Aktinomykose in der Armee. *Deutsche Monatschr. f. Zahnh.* (1905), **23**, 257 to 277, 1 pl.

* Wingate, J. L. Actinomycosis, *Journ. Comp. Med. Vet. Arch.* (1896), **17**, 276 to 281.

* Von Winiwarter, A. Un cas d'actinomycose. *Ann. Soc. méd.-chir. de Liège* (1890), **29**, 129 to 131.

* Winkler, M. Ueber eine eigenartige benigne Streptomycosis bullosa in der Blindenanstalt Könitz bei Bern. *Cor.-Bl. f. schweiz. Aerzte* (1903), **33**, 569 to 571.

* Winter. Ein Fall von Aktinomykosis bei einem Soldaten. *Deutsche mil.-ärztl. Ztschr.* (1886), **15**, 188 to 193.—* Die Pilze *in* Rabenhorst's Cryptogamen-Flora von Deutschland, Oesterreich und der Schweiz. Pilze. Leipzig (1884), 59; 60; 66.

* Wolff, I. Ueber einem Fall von Actinomycose. *Jahresb. d. schles. Gesellsch. f. vaterl. Kult.* (1884) **62**, 113 to 121. *Also: Breslau. ärztl. Ztschr.* (1884), **6**, 284 to 286; (1885), **7**, 31.

Wolff, Max. Aetiologie der Aktinomykose. *Verhandl. d. X. internat. med. Cong.* (1890), Berl., 1891, **2**, 3. Abth., 57 to 62.—Cultur und Uebertragung des Actinomyces. *Münchener med. Wchnsch.* (1890), **37**, 224.—* Ueber Actinomycose, mit Demonstration. *Verhand. d. deutsch. Gesellsch. f. Chir.*, Berl. (1891), **20**, pt. 1, 148 to 155. *Also Centralbl. f. Chir.* (1891), **20**, 18, Beil. 67 to 75.—Ueber Aktinomykose. *Wien. med. Wchnsch.* (1891), **4**, 175.—Reinculturen des Strahlenpilzes. *Deutsche med. Wchnsch.* (1894), **20**, 208.—† and Israel, J. Ueber Erzeugung von Impfaktinomykose mittels Culturen des Strahlenpilzes. *Verhandl. d. Berl. med. Gesellsch.* (1890), (1891), **21**, 68 to 70. *Also: Deutsche med. Wchnsch.* (1890), **16**, 247 to 248; *Berl. klin. Wchnsch.* (1890), **27**, 309.—Ueber Reincultur des Actinomyces und seine Uebertragbarkeit auf Tiere. *Virch. Arch.* (1891), **126**, 11 to 59, 8 pl.—Zur Actinomyces-Frage. (Zugleich eine Entgegnung an Herrn Dr. van Niessen). *Ibid* (1898), **151**, 471 to 488.

Wolfler, A. Zur Behandlung der Aktinomykose. *Wien. klin. Wchnsch.* (1906), **19**, 1519 to 1520.

* Wolisch, A. Ein Fall von retrocoecaler Actinomycose. *Internat. klin. Rundschau* (1894), **8**, 369; 414; 445.

Wooldridge, G. H. Actinomycosis and botriomycosis. *Journ. Comp. Path. & Therap.* (1907), **20**, 189–202.

† Wortley Axe. Actinomycosis of the stomach. *Veterin.* (1882), **55**, 811; (1886), **59**, 313. Ref.: *Jahresb. f. gesammt. Med.* (1886), **1**, 550.

* Wrede, L. Ueber haematogene Osteomyelitis durch Aktinomykose. *Verhandl. d. deutsch. Gesellsch. f. Chir.* (1906), XXX **35**, pt. 2, 419–427.

Wright, J. H. A case of mycetoma (Madura foot). *Journ. Exp. Med.* (1898), **3**, 421–433, 2 pl. *Also; Trans. Ass. Am. Physicians*, Philadelphia (1898), **13**, 471–482, 4 pl.; *Journ. Bost. Soc. Med. Sci.*, (1897–98), **2**, 128–130, 1 pl.—Actinomycosis. *In:* Buck's Reference Handbook of the Medical Sciences (1900), **1**, 97–101.—Madura foot. *Ibid.* 101.—The biology of the microorganism of actinomycosis. *Pub. Mass. Gen. Hosp.* (1905), **1**, 1–56, 10 pl. *Also Journ. Med. Research* (1904–05), **13**, 349–404, 10 pl.—Actinomycosis *In:* Osler's Modern Medicine, (1907), **1**, 327–339.—Madura foot (Mycetoma). *Ibid.* 344–346.—Nocardiasis. *Ibid.* 340–352.

Wulff. Die Strahlenpilzkrankheit. *Berl. thierärztl. Woch.* (1900), 13–17.

Wyer, O. F. Actinomycosis. *Brit. Med. Journ.*, (1906), **2**, 1128.

* Wynne, E. T. Clinical and post-mortem notes of a case of actinomycosis. *St. Barth. Hosp. Rep.*, London (1889), **25**, 159–165.

* Yakimovitch, N. M. (Actinomycosis of lungs and pleura). *Vratch* (1888), **9**, 125; 170; 211; 228; 267.

* Zamuravkin, K. I. Sluchai aktinomikosa. (Case). *Voyenno-med. J.*, St. Petersb. (1905), **2**, med.-spec. pt., 233–235.

* Zaufal. Actinomycotische Abscesse in der Nähe des Warzenfortsatzes. *Prag. med. Woch.* (1894), 29.

* Zdanovski, A. (Actinomycosis of lungs). *Obsk. Tulsk. Vratch* (1890), **28**, 25–28.

* Zechmeister, H. Ein Fall von Aktinomykose mit Restitutio ad integrum nach innerlichem Gebrauche von Jodkalium, *Wien. med. Bl.* (1895), **18**, 203.

* Zederbaum, G. A case of actinomycosis. *Dental Reg.* (1907), **61**, 460–467. Also: *Dental Digest* (1907), **13**, 591–599.

Zeisler, Joseph. Notes on a case of actinomycosis. *Journ. Cutan. Dis.* (1906), **24**, 510–512.

* Zemann, A. Ueber die Actinomykose des Bauchfelles und der Baucheingeweide beim Menschen. *Med. Jahrb.*, Wien, (1883), 477–495. *Also, transl.: Riforma med.* (1885), **1**.

Ziegler, Ernest. Eng. transl. 9. rev. German ed. of Lehrbuch der allgemeine Pathologie, New York, (1900), Actinomycosis or ray-fungus, 515.

Ziegler, Paul. Aktinomykose des Gesichtes und Halses. *Münch, med. Woch.* (1892), **39**, 406–407.—Demonstration von Aktinomyces.—Reinculturen. *Münch. med. Woch.* (1895), **42**, 373.

* Zietzschmann, H. Primäre Aktinomykose der Haut mit Generalisation in verschiedenen Lymphdrusen vom Schwein. *Sachs. Vet.-Ber.* (1903), 260.

* Zimmermann, K. (A new acid pathogenic kind of streptothrix and the acid bacilli generally). *Orvosi hetil.*, Budapest (1903), **47**, 145. *Also Ungar. med. Presse* (1903), **8**, 189; *Pest. med.-chir. Presse* (1903) **39**, 704.

* Zondek, M. Beitrag zur Lehre von der Bauchaktinomykose. *Deutsche Ztschr. f. Chir.* (1903), **69**, 49–59.

* Zopf, W. Zur Morphologie des Spaltpflanzen (Spaltpilze und Spaltalgen), 74 pp., 7 pl., Leipzig, (1882), 2. ed. (1884), 3. ed. (1885).—* Die Pilze: *In* Handbuch der Botanik, Breslau (1890), 500 pp.

* Zwintz. Pseudoparametritis actinomycotica. *Wien. med. Presse*, (1905).

ILLUSTRATIONS.

(Colored photographs and photomicrographs are by Charles Martin, photographer, Bureau of Science, Manila.

PLATE I.

FIG. 1. Cultures of *S. eppingeri* from Foulerton's case.
2. Culture of *S. eppingeri* from original strain.

PLATE II.

FIG. 3. Culture of *S. maduræ* from Foulerton.
4. Culture of *S. maduræ* from Strong's culture.

PLATE III.

FIG. 5. Culture of *S. maduræ* from Vincent's strain.
6. Culture of *S. actinomyces* from Binot.

PLATE IV.

FIG. 7. Culture of *S. nocardii.*
8. Culture of S. *capræ.*

PLATE V.

FIG. 9. Culture of *S. canis.*
10. Culture of *S. freeri.*

PLATE VI.

FIG. 11. Culture of *S. freeri,* showing structure of individual colony.
12. Culture of *S. chalcea.*

PLATE VII.

FIG. 13. Section from experimental lesion in monkey produced by inoculation with culture of *S. eppingeri,* × 60.
14. Unstained smear preparation from culture of *S. eppingeri,* × 600.
15. Smear preparation from culture of *S. eppingeri* stained by gentian violet, × 800.
16. Section from experimental lesion in mesentery of monkey produced by inoculation with culture of *S. eppingeri,* × 600.

PLATE VIII.

FIG. 17. Smear preparation from culture of *S. maduræ* stained by gentian violet, × 400.
18. Same as No. 17, × 600.
19. Section from experimental lesion in monkey produced by inoculation with culture of *S. maduræ,* × 60.
20. Smear preparation from culture of *S. canis* stained by gentian violet, × 650.
21. Same as No. 20, × 650.

PLATE IX.

FIG. 22. Smear preparation from culture of *S. maduræ* (Strong's culture) stained by gentian violet, × 500.

23. Same as No. 22, × 700.

24. Same as No. 22 excepting that the preparation is unstained, × 500.

25. Smear preparation from culture of *S. actinomyces* (Pasteur Institute) stained by gentian violet, × 700.

26. Section from experimental lesion in monkey produced by inoculation with culture of *S. maduræ*.

PLATE X.

FIG. 27. Smear preparation from culture of *S. caprœ* unstained, × 600.

28. Section from experimental lesion in monkey produced by inoculation with culture of *S. actinomyces*, × 600.

29. Smear preparation from culture of *S. actinomyces* stained by gentian violet, × 650.

30. Smear preparation from unstained specimen of *S. actinomyces*, × 650.

PLATE XI.

FIG. 31. Smear preparation from culture of *S. caprœ* stained by gentian violet, × 650.

32. Section from experimental lesion from monkey produced by inoculation with culture of *S. caprœ*, × 600.

33. Smear preparation from culture of *S. caprœ* stained by gentian violet, × 1100.

34. Section from experimenal lesion in monkey produced by inoculation with culture of *S. caprœ*, × 60.

PLATE XII.

FIG. 35. Smear preparation from culture of *S. chalcea* stained by gentian violet, × 850.

36. Smear prepration from cultures of *S. nocardii* stained by gentian violet, × 850.

Fig. 2.

Fig. 1.

PLATE I.

Fig. 4.

Fig. 3.

PLATE II.

Fig. 6.

Fig. 5.

PLATE III.

Fig. 8.



Fig. 7.

PLATE IV.

Fig. 10.

Fig. 9.

PLATE V.

Fig. 12.

Fig. 11.

PLATE VI.



FIG. 13.



FIG. 14.



FIG. 15.



FIG. 16.

PLATE VII.



FIG. 17.



FIG. 18.

FIG. 19.



FIG. 20.



FIG. 21.

PLATE VIII.



FIG. 23.

FIG. 22.



FIG. 24.

FIG. 25.



FIG. 26.

PLATE IX.



FIG. 27.



FIG. 28.

FIG. 29.



FIG. 30.

PLATE X.



FIG. 31.



FIG. 32.



FIG. 33.



FIG. 34.

PLATE XI.



FIG. 35.



FIG. 36.

PLATE XII.

TRICHOCEPHALIASIS.

(WITH A REPORT OF FOUR CASES, INCLUDING ONE FATAL CASE.)

By W. E. Musgrave and M. T. Clegg, with a bibliography by Mary Polk.

(*From the Biological Laboratory and the Library, Bureau of Science.*)

OUTLINE.

Introduction.
Case Reports.
Literature of the Disease.
Trichocephaliasis.
Synonyms.
Definition.
Etiology.
Contributing causes.
Trichocephalidæ (by P. E. Garrison).
General discussion of the genus.
Trichuris trichiura.
Modes of transmission.
Direct.
Indirect.
Water.
Food.
Insects (flies, etc.).
Soil.
Dust, clothing.
Air.
Animals.
Trichocephaliasis—Continued.
Etiology—Continued.
Trichocephalidæ—Continued.
Pathogenesis (parasitism).
General.
Manner of infection.
Pathology of the disease.
Symptoms.
Incidence.
Clinical varieties.
Intestinal.
Appendicial.
Mild.
Severe.
Trichocephaliasis of animals.
Diagnosis.
Prognosis.
Prophylaxis.
Treatment.
Bibliography.

INTRODUCTION.

Our own experiences and a careful review of the literature of whipworm infections, convinced us that the subject has much more importance in human pathology than is generally assigned to it.

For this reason we have decided to publish four of our cases in detail, review the most important articles in the literature and then discuss the question of trichocephaliasis in monographic form with special reference to the disease-producing powers and clinical manifestations of infection with these parasites.

While it is true that the majority of investigators who have published the results of original work have attributed pathogenic properties to these parasites, and although these observations are supported by such authorities as Seifert, Stiles, Metchnikoff and others, to-day the majority of text books on medicine barely mention these infections and in general state that they are of little or no importance from a medical standpoint.

CASE REPORTS.

CASE I.—*Trichocephaliasis with severe progressive secondary anaemia followed by death; autopsy.*

L. G. N., male Filipino, 24 years of age, was admitted to St. Paul's Hospital August 9, 1908, died September 16. *Family history:* Father and mother are both living and in good health; two brothers died of smallpox. *Previous illness:* As a child the patient had chickenpox, a few mild attacks of "malaria," and five years ago suffered from an abscess of the parotid gland. *Present illness:* Three weeks before admission the patient began to feel dizziness, weakness, slight fever in the afternoons and ringing in the ears. This condition gradually grew worse and about ten days before admission he noticed beginning œdema of the ankles, hands, and face.

On admission the patient complained of shortness of breath, great weakness, ringing in the ears, palpitation of the heart, nausea, lack of appetite, dyspeptic symptoms and slight diarrhœa.

Examination.—The patient is a rather large, well-developed Filipino with marked œdema of the extremeties and face, marked pallor of the mucous membrane and rapid, shallow respiration. The circulatory system appears normal, except for a slight hæmic murmur heard at the apex of the heart and an increased pulse rate to 110, when the patient is in an upright position.

Blood examination as follows: Red blood cells, 1,800,000; leucocytes, 10,000; hæmoglobin, 20 per cent. Differential count: Mononuclear cells, 65 per cent; polynuclear cells, 30 per cent; transitional, 5 per cent. There are no nucleated red blood cells and but a moderate poikilocytosis. No eosinophile cells are present in a count of 500.

Respiratory system, negative. *Alimentary system* shows the following changes: Lips and other mucous membranes extremely pale, tongue large and flabby with a dark, almost black, streak down the center. Nothing abnormal is found in the abdomen, excepting a slight accumulation of fluid; the liver is slightly enlarged; the spleen is not palpable. The examination of the stomach contents shows no abnormality; examination of the stool after a saline cathartic shows enormous numbers of the ova of *Trichuris trichiura*. The urine is normal by chemical and microscopic examination. *Nervous system:* There is no pain, and the reflexes are normal. *Locomotor system:* Œdema and considerable flabbiness of the muscles are the only abnormalities noted. *Cutaneous system:* The skin appears normal and there is no enlargement of the superficial lymphatics. The patient's condition grew gradually worse from the time of admission with accentuation of the following symptoms: Œdema became general and progressed until death; the anæmia was also progressive, and the same general character of blood picture was maintained until the end. A count made a few days before death showed the following: Red cells, 844,000; hæmoglobin, 18 per cent. At this time there were still no nucleated red cells and no eosinophile cells to be found. Nausea, from which the patient suffered upon his admission, became more constant and toward the end vomiting was a very distressing and unmanageable symptom. The nervous and mental symptoms became more marked and during the last week of illness there was active delirium. The patient finally died of asthænia.

There were no hæmorrhages at any time and no severe diarrhœa. Repeated examinations of the stool always demonstrated very large numbers of *Trichuris* eggs. The temperature was practically normal throughout the course of the disease. Three separate attempts were made to dislodge the parasites, which were the only discoverable cause of the anæmia, the following substances being used for

this purpose: Thymol in doses of 8 grams on two occasions and the well-known eucalyptus-castor oil and chloroform mixture on one occasion. A few *Trichuris* were obtained with two of these treatments, about 15 or 20 in all being secured. The stool from the third treatment was unfortunately destroyed without examination. Other treatment administered consisted in stimulation and in attempts to build up the nutrition.

Autopsy was performed one hour after death by Dr. Gilman and his report of the findings is as follows:

Body of a sparely nourished Filipino 157 centimeters long; rigor mortis absent; body still warm; pupils moderately dilated; equal; distinct yellowish tinge of sclerae; the entire body has a yellowish tinge. There is no general glandular enlargement; abdomen flat; no scar on penis. Shins clear.

Subcutaneous fat is present in good amount, is of a lemon-yellow color, more brilliant than normal. Muscles are practically of normal color.

Peritoneum is everywhere smooth and glistening. The mesenteric and omental fat is present in good amount, similar in shade to the subcutaneous fat. The general visceral arrangement is normal. Coils of small bowel are in general contracted, small, appendix region is free from adhesions. Bladder is distended with urine extending 11.5 centimeters above the symphysis. Diaphragm at the upper border of the 5th rib on the left, lower border of the 4th rib on the right.

Pericardium is smooth and glistening, contains 1 or 2 cubic centimeters of clear fluid. Heart is firmly contracted, measures 10.5 centimeters from the auriculo-ventricular ring to apex, surface smooth, sub-epicardial fat well-developed. The right side of the heart is practically empty, but for a small amount of fluid blood. Valves are normal and competent. The left side of the heart contains a small, dark, post-mortem clot. The mitral valves are normal as are the aortic. Base of the aorta is smooth and elastic. The wall of the left ventricle measures 2 centimeters at its thickest portion and the right measures 0.6 centimeter. The heart muscle is of a firm, reddish-brown color. Heart weight 300 grams.

Lungs: The left pleural cavity is obliterated, the lung being adherent to the chest wall and diaphragm by firm, old, fibrous adhesions. The lung is apparently crepitant everywhere, is pale and on section is air-containing throughout, of a pasty yellowish-gray color. The right lung lies free in its pleural cavity except at the base, where it is adherent to the diaphragm; it presents a similar appearance to the left. Apparently air-containing throughout. On section similar to opposite lung.

Spleen is small, measuring 7.3 by 4.5 by about 2.5 centimeters, surface smooth, color dark grayish-purple. On section there is a decided increase in the connective tissue. The pulp is moist, of a brownish color. Spleen weighs 35 grams.

Kidneys: The left kidney measures 10 by 6 by 4 centimeters, its capsule strips readily, leaving a smooth, pale yellowish surface somewhat mottled with areas of red. On section the cortex measures, 0.6 centimeter, entire organ is of a pale yellowish color, contains a very small amount of blood. The right kidney measures 9.8, by 5.5 by 3.5 centimeters, is similar throughout to opposite organ. Combined weight of the two organs is 270 grams. Suprarenals are normal.

Pancreas is firm, of a yellowish color. On section it is somewhat pale, although apparently normal.

Liver measures 24 by 14 by 8 centimeters, surface is smooth, edge sharp, organ is somewhat soft and putty-like. On section the parenchyma is of a brownish color, homogeneous. Gall-bladder contains several cubic centimeters of dark, slightly inspissated bile. Duct patent. Weight of liver is 1,020 grams.

Bladder measures 15 by 12 centimeters, walls are stretched and thinned. It contains a large amount of clear, amber-colored urine. Mucosa is very pale, otherwise normal. Prostate apparently normal.

Stomach: Mucous membrane extremely pale with small, hæmorrhagic areas near the pyloric end; otherwise the organ appears normal.

Small intestine: Contents, greenish mucus. The mucosa shows the same pale condition. Peyer's patches and solitary follicles prominent, but not congested. Numerous *Trichuris trichiura* are present, the greater number being found in the lower part of the ileum.

Large intestine: On opening the ileocæcal valve two hundred trichiuris are counted. These parasites extend the entire length of the gut. The mucosa of the descending colon shows a few healed amœbic ulcers. With the exception of the extremely pale condition of the mucosa, the organ appears normal.

Anatomic diagnosis.—Severe grade of secondary anæmia.

Specimens of the worms were submitted to Mr. G. F. Richmond, chief of the chemical laboratory, for determination as to the presence of pigments of blood origin and he reports that no such pigments are found.

Dr. Philip E. Garrison, medical zoölogist, Biological Laboratory, Bureau of Science, reports as follows:

"The whip-worms received on September 16 have been carefully examined and compared with specimens identified in Dr. Stiles' laboratory in Washington. They appear to agree in every material character with *Trichuris trichiura* (Linnaeus 1761)=*Trichocephalus dispar* Rudolphi, 1801. The specimens are No. 258 in the Bureau of Science Helminthological Collection."

CASE II.—*Severe infection with whip-worms, with diarrhœa, muscular cramps, dizziness, œdema and indigestion; patient left hospital after ten days observation.*

F. E., male Filipino, age 22, was admitted to St. Paul's Hospital March 10, 1906, and left March 21. His *family history* and the *history of previous diseases* give no data of importance. *Present illness:* For about six months previous to admission the patient has suffered from diarrhœa of a somewhat intermittent character. The stools were only one or two on some days, but generally they were from five to ten or more in twenty-four hours, being soft and pasty. Other symptoms during this time were dizziness, weakness, palpition of the heart, tinnitus aurium, cramps in muscles of the legs and chest, anorexia and indigestion.

On admission the patient complained of weakness, shortness of breath, palpitation of the heart, nausea, diarrhœa, ringing in the ears, partial loss of voice and melancholic tendencies.

Examination shows a pale, somewhat emaciated Filipino with moderate œdema of the feet and legs. Examination of the circulatory system demonstrates a weak pulse of 110 in the upright position: the arteries are soft and there is a soft murmur at the apex of the heart. The spleen not palpable, and blood examination is as follows: Erythrocytes, 2,400,000; leucocytes, 9,000; hæmoglobin 23 per cent.

There are no nucleated red cells and the differential leucocyte count shows a slight relative increase in large mononuclears. There is no eosinophilia. The respiratory system appears normal. Examination of the alimentary system gives the following details: Tongue, large and flabby with teeth marks on the edges, but no dark streak in the center. The abdominal examination is negative and the stool contains very many ova of *Trichuris trichiura.* The urinary system (including urine examination), the reproductive, cutaneous, and locomotor systems appear normal, except for moderate œdema of the lower extremeties. The reflexes

are present and nothing abnormal is found upon examination of the nervous system.

During the patient's ten days' stay in the hospital, two unsuccessful attempts were made to remove the *Trichuris*. Both thymol in 6 gram and santonin and calomel in 0.250 gram doses were used. The patient left the hospital in a worse condition than when he entered.

In this case repeated and careful examinations were made and the data obtained could only be explained by assuming the cause of disease to be *Trichuris trichiura*.

CASE III.—*Trichocephaliasis with severe secondary anæmia; under observation and treatment for one month without improvement.*

D. V., a male Eurasian from Singapore came under our observation in September, 1904, and left the Philippine Islands in October of the same year. The family history and history of previous illnesses of this patient are unimportant.

Present illness.—About one year after arrival in the Philippine Islands the patient began to notice that he was becoming "anæmic," with shortness of breath and palpitation of the heart. About one month before he consulted us, diarrhœa developed and there was some œdema of the legs. At the time he came under our observation the principal subjective symptoms were dizziness, with shortness of breath and palpitation of the heart upon exertion, indifferent appetite, with nausea more marked in the morning, occasional vomiting, and moderate diarrhœa. There was a general feeling of "nervousness" and tendency toward melancholia.

Examination.—A rather severe secondary anæmia was present which could not be explained, except by the presence of a rather severe *Trichuris* infection. The stools after a saline cathartic often contained as many as 20 eggs in one field of the microscope, an AA objective and No. 4 ocular being used.

The blood examination was as follows: Erythrocytes, 2,800,000; leucocytes, 6,400; hæmoglobin, 36 per cent.

There were no nucleated red cells and the differential leucocyte count was approximately normal.

One unsuccessful attempt was made to dislodge the parasites by thymol. Good food, tonics, and general hygiene were prescribed, but the patient did not improve and he left the Philippine Islands about one month after we first saw him.

CASE IV.—*Embolism of the left coronary artery caused by an adult Trichuris trichiura.*[1]

It has not been possible to obtain a clinical history of very much importance in this case. The patient was a Filipino boy, 17 years old, and had been ill for about a week. For approximately the first five days he complained of pain in the chest, cough, shortness of breath, and he vomited once or twice. Slight diarrhœa developed and continued throughout the trouble. Pulse 130, temperature at first from 38° to 39°. Two days before death the symptoms became very much aggravated; vomiting more frequent, temperature 40°.5. Death occurred suddenly.

Autopsy (twenty-three hours after death).—Petechial hæmorrhages in skin of shoulders and occasional ones over the rest of the body. No glandular enlargements.

Lungs: Lower lobes much congested, dark, grayish-red in color. Center of

[1] This case is not reported as one of importance in elucidating the question of the pathogenicity of whip-worms, but it is mentioned as being a very curious finding for which we have no explanation to offer.

outer surface of left lobe shows a hæmorrhagic infarct about 4 centimeters in diameter, penetrating about 3 centimeters into the lung substance. Lower edge of right lower lobe has a large infarct about 11 centimeters in length and about 6 centimeters broad. On section no embolism could be found in the arteries.

Heart: Very slight atheromatous patches in arch of aorta. The left coronary artery contains an adult *Trichuris trichiura,* with about one-third of the posterior portion remaining free in the aorta.

Liver: The liver is friable, dark red in color (slight post-mortem change). The *spleen* is dark red, slightly mottled, and of firm consistency. *Kidneys* congested.

Moderate injection of the mucous membrane of the large intestine, which contains a small amount of mucus; a number of *Trichuris* were found in the cæcum.

Anatomic diagnosis: Embolism of the left coronary artery due to adult *Trichuris;* hæmorrhagic infarcts of the lower lobe of either lung.

Numerous other cases of *Trichuris* infection much less severe, but with suggestive symptoms, could be mentioned, but in them the diagnosis is not so clear and they will only be considered in the general discussion in the second part of this paper.

GENERAL REVIEW OF THE LITERATURE.

The first description of the parasite which we now recognize as *Trichuris trichiura* (Linnaeus 1761) was that of Morgagni(61) who found the worms in the cæcum of a man at autopsy. Röderer(73), in collaboration with his students, made additional contributions to the description of the parasite, and Büttner (1761) examining their specimens and mistaking the posterior for the anterior end of the parasite named it *Trichuris.* Linnæus (1771) introduced the name *Ascaris trichiura.* Goeze(34) recognized Büttner's error in thinking the smaller end to be the head of the parasite and changed the name from *Trichuris* to *Trichocephalos.* According to Blanchard(8) Morgagni, Röderer, Wagler, Happ, and Wristberg thought there were two species of these worms, but Müller (1778) showed the two kinds to be male and female; Schrank (1788) recognized the parasite under the name of *Trichocephalus hominis,* and the same name was used by Gmelin (1789); Rudolphi (1801) described the parasite under the name of *Trichocephalus dispar* and this name has been the one in most common use until within the last few years.

Owing to the adoption of different rules of nomenclature there is at present considerable difference of opinion among zoölogists as to the correct name of the parasite. Stiles maintains *Trichuris, trichiura,* while most of the European authors adopt *Trichocephalus trichuris.*

Without entering into the discussion, we have decided in this report to use the name recommended by Stiles.

Barth(4) reports a case of severe infection with *Trichuris* as the only explanation of severe meningitis-like symptoms in a patient. This is perhaps the first report in which a pathogenic rôle in human beings has been attributed to this parasite.

Davaine(19) examined a large series of stools in Paris and concluded that probably one-half the people of that city harbor this parasite.

Gibson(32) reports an interesting case of severe nervous symptoms with paralysis in a girl who had a very severe *Trichuris* infection. Calomel 0.065 gram and rhubarb powder 0.65 gram were given night and morning for some days. "On two occasions the child passed a small chamber vesselful of worms; species,

Trichocephalus dispar." More worms were passed on another date and following this, the child rapidly recovered.

Pascal (67) reports the case of a girl, 4 years old, who died of brain symptoms simulating meningitis. Many *Trichuris* were found in the intestine at autopsy and no pathologic changes were discovered. The author believed the parasites to have been the cause of death.

Burchardt (14) reports two cases in which he regarded *Trichocephalus* as the causative agent. Vomiting, diarrhœa, headache, dizziness, and marked anæmia were present in his first case. There was slight improvement upon treatment, followed by relapse with icterus, bloody vomit, aphonia, and marked anæmia. Complete recovery followed the expulsion of a large number of the worms. Burchardt's second case was similar to the first, excepting that the symptoms including the nervous ones were much less marked.

Mégnin (56) pointed out the fact that dogs which harbor large numbers of a *Trichuris* suffer from a pernicious anæmia.

Lutz (53) working in Brazil, found eggs in the stools of man, pigs, and cats. In man, about one-third of the specimens examined contained the ova.

Erni (24) believed the Trichuris to be one of the causes of beri-beri.

Grassi (35) describes the swallowing of *Trichuris* eggs by a student who 28 days later found the first eggs in his stools.

R. Blanchard (9) in a review of the subject of *Trichuris* infections states that Vix, Leukart, and Brumpt all give observation of the transfixation of the mucosa of the caecum by *Trichuris* and believes that this definite fastening of the worms explains their constant absence from the fæces in infected cases. However, Klebs (50), Heller (45) and Wichmann (89) do not believe that the worms attach themselves to the mucosa, but state that they are maintained in place by a mucus capsule resting upon the mucous membrane.

Moosbrugger (59) reports three cases. The first was in a boy a year and a half old who had a severe diarrhœa and with the usual symptoms accompanying the latter. During an attack of bronchitis with fever the eggs of *Trichuris*, which had been very numerous in the stools, disappeared. The author's second case was in a three year old boy with symptoms similar to those in the first. The patient died of croup and at the autopsy 442 male and 445 female worms were removed from the bowel. The mucous membrane was pale in some places and in others there were zones of congestion, with red spots and some erosions and ulcerations in others. The third case was that of a girl three and half years old, who had suffered for six months with severe diarrhœa of twenty or more stools daily. The stools were tough in consistency and usually contained some blood. There were 1,650 *Trichuris* eggs in one cubic centimeter of fluid stool. The child was very anæmic, but recovered under treatment. Moosbrugger considers the parasites to be the cause of the symptoms in his cases.

Boas (11) considered the *Trichuris* to be the cause of severe chronic diarrhœa, emaciation and anæmia in his patient, who was a man 71 years old.

Huber (47) considers the clinical significance of this parasite to be generally underestimated. He notes the presence of the worms in the cæcum, appendix, and occasionally in the ileum. The symptoms depend upon the severity of the infection and are local (diarrhœa) and reflex (aphasia, etc.).

Askanazy (3) demonstrated an iron-containing pigment in the intestinal epithelial cells of *Trichuris* and concluded it to be a blood-sucking parasite. He also showed that the parasite attached itself to the intestinal wall by piercing a loop of mucosa or by inserting the head into the glandular tubules of the mucosa. He believed that during certain diseases, or after death of the host, the parasites disengage themselved from the mucous membrane and that this fact explains why at autopsy they are usually found free in the intestine. Becker (5)

in accepting the hypothesis of Askanazy, explains his failure to secure the worms when his patient was treated with benzine, by stating that the worms were killed in place and so remained attached to the mucosa.

Morsasca (62) reports one case of severe progressive anæmia accompanied by abdominal pains, nervous symptoms, and diarrhœa and considers that *Trichuris*, which were present, to be the cause of the condition.

Federolf (27), a Russian military surgeon, notes that Russian authors have repeatedly called attention to the serious symptoms which may be produced by *Trichuris* infections. He reports one case in which the prominent symptoms were psychic depression, slight diarrhœa, retention of urine, fever of intermittent character, nausea, and palpitation of the heart. Treatment by santonin was followed by the disappearance of the eggs from the stools and the complete recovery of the patient.

Heine (43) encountered *Trichuris* in many sick deer and in two of the animals which he found dead the presence of these parasites was the only condition which could explain the result. Hausmann (42) believes that certain of the symptoms of infection with this parasite point to the production of some form of toxin. Guinard (41) reports finding nematode worms in the cæcum at operations for appendicitis.

Metchnikoff (57) considers both the clinical and pathologic facts to be convincing of the important rôle of eelworms and whip-worms in appendicitis. Guiart (39, 40) insists upon the important pathogenic rôle of *Trichuris* in human pathology. The parasite fixes itself on the mucous membrane of the cæcum or appendix and becomes an inoculating agent of bacteria. In one case of appendectomy an adult *Trichuris* was found attached to the mucosa of the appendix and the author believes it to have been the cause of the disease.

Girard (33) reports a case of appendicitis with general peritonitis due to infection with *Trichuris* (*Trichocephalus hominis*). The patient was a girl, 8 years old, who was recently convalescent from typhoid fever. She was re-admitted to the hospital suffering from diphtheria and a vaginal discharge. Under antitoxin treatment there was recovery from the diphtheria. Symptoms of acute, severe appendicitis developed and the appendix was resected on the fourth day. At operation a general peritonitis was encountered and after resection of the appendix and drainage, complete recovery resulted. The appendix was fixed *en masse* and sectioned. In general the organ showed but slight pathologic change except in the distal third where there was rather severe inflammation. Section in this area showed partial occlusion of the lumen caused by two of the adult *Trichuris* which were firmly attached to the mucosa. The author describes and illustrates the inflammatory zone around the attachment of the worms to the mucosa and, although no perforation is mentioned, he considers the peritonitis to be due to the condition of the appendix. Finally, Girard believes he has demonstrated the penetrating powers of the whip-worm; he states that the parasites play a septic rôle at least by transmission of pathogenic bacteria and he believes with Metchnikoff, that these parasites may cause inflammatory lesions which may lead to grave consequences.

Schiller (75) reports a case of heavy infection with *Trichuris* in which high fever was a symptom. Hemmeter (46) considers that single specimens of the parasite do not cause symptoms, but if larger numbers occur in the colon, diarrhœa and nervous symptoms may appear.

Becker (5) reports two cases of *Trichuris* infection. In the first the patient was a woman suffering from chronic progressive anæmia with dizziness, cramps in the muscles, abdominal pain, and palpitation of the heart and the stools contained many *Trichuris* eggs. Physical examination was negative. At the

time she came under the author's observation the blood examination showed as follows: Marked poikilocytosis and macrocytes were numerous; erythrocytes 3,456,000; leucocytes 6,250; hæmoglobin 35 per cent. After three week's observation it was: Erythrocytes 3,200,000, leucocytes 6,000, hæmoglobin 25 per cent. The differential count of the leucocytes showed: Polymorphonuclears 69, lymphocytes 26, mononuclears 3, eosinophiles 2. There were no nucleated red cells. Numerous *Trichuris* eggs were present in the stools. The patient was treated with eight enemas of solutions of benzin. No parasites were found during or after these treatments, but the eggs disappeared from the stools and were not again seen. The patient rapidly recovered after the disappearance of the eggs from the stools. The author's second case was that of a woman 26 years old suffering from chronic progressive anæmia, palpitation of the heart, dizziness, nausea, and diarrhœa. The stool contained numerous *Trichuris* eggs. The physical examination was negative. The blood examination showed: Erythrocytes, 2,852,000; leucocytes, 8,700; hæmoglobin, 25 per cent. Normoblasts were numerous but megoloblasts were not present. There was some improvement in this case under treatment by arsenic and iron. The patient left the hospital before further observation could be made. Becker believes the *Trichuris* to be responsible for the condition in both these cases and furthermore compares its action with that of *Agchylostoma* and the *Dibothriocephalus latus.*

Ward (88) states that since Askanazy has shown the occurrence of hæmoglobin in the alimentary canal of these worms, the fact that they nourish themselves on the blood of the host can not be doubted.

Sandler (74) reports a fatal case in a boy twelve years old. The patient when admitted to the service complained of fainting spells, anæmia, weakness, palpitation of the heart, fever of an irregular, intermittent type, yellow and brownish spots on the legs, headache, nausea, vomiting, diarrhœa with blood at times, abdominal pains and tinnitus aurium. On examination there was marked anæmia, hæmic murmur of the heart, and weak, rapid pulse. The spleen and lymphatics were not enlarged. Blood examination showed: Erythrocytes, 1,200,000; leucocytes, 30,600; hæmoglobin, 28 per cent. There was poikilocytosis, but no nucleated red cells. The stools contained eggs of *Trichuris.* After about two weeks of treatment and observation the blood examination was as follows: Erythrocytes, 690,000; leucocytes, 14,000; hæmoglobin, 20 per cent. The patient's condition grew progressively worse with accentuation of the principal symptoms mentioned, and others also developed. There was bleeding of the nose, the sense of fear became marked, fever became more severe, reaching 39° in the afternoons, and for some time before death which occurred in coma, there was delirium and very frequent vomiting. The diarrhœa improved so that the bowels were practically normal for some time before death. Autopsy could not be obtained, but the author is convinced that *Trichuris* was the etiologic factor in the condition.

French and Boycott (28) in reporting upon the prevalence of *Trichuris* infections in the patients in Guy's Hospital, London, makes the following observations. Stools of 500 patients were examined and *Trichuris* eggs were found 39 times or 7.8 per cent. The infection was rare in children under 5 years of age and was about equal in the sexes. The authors also compile the statistics as to prevalence from the following sources: Leuckart (52) in Erlangen 11.1 per cent, Erlangen insane 100 per cent, Dresden, 2.5 per cent, Kiel 32.2 per cent; Cobbold (17) in Dublin 90 per cent, Greenwich 69 per cent; Davaine (20) in Paris 50 per cent, Naples 100 per cent; Blanchard (9) in Bâle 23.6 per cent; Garrison, Ransom, and Stevenson (31) in 10.8 per cent of insane; Ashford, King, and Igaravidez (2) 6 per cent; Boycott (12) in Cornish Miners 79 per cent. Garrison (30) working in the Philippine Islands found the infection in 59 per cent of adult prisoners.

Osler (65) states that "The *Trichocephalus* rarely causes symptoms." He notes without comment the recent observations in which severe anæmia has occurred in connection with infections with this parasite. Allen J. Smith (80) considers the parasite to be of little pathologic importance, stating that apparently it does but little damage and practically never gives rise to appreciable symptoms of its presence. Emerson (23) states that "This parasite is harmless as a rule, but may cause enteritis and the severest and even fatal anæmia."

Poledne (68, 69) reports one case. The patient was a sailor who was admitted to the hospital suffering from fever and severe diarrhœa. The fever was typhoid in character, but careful examination by modern laboratory methods gave negative results for typhoid, paratyphoid, Malta fever, and tuberculosis. Anæmia, emaciation, and nervous symptoms were present. The blood showed a slight leucocytosis and a relative eosinophilia. There were numerous ova of *Trichuris trichiura* in the stools. The fever and diarrhœa subsided after treatment by thymol and by benzine enemas, the eggs disappeared from the stools and a rapid and permanent cure followed.

Vix (86) considers the parasite to be pathogenic and states that it may produce ulceration of the intestinal mucosa. Rippe (72) reviews the literature of *Trichuris* infections and reports two cases occurring in Russia. His first patient was a woman who had been ill for three years. The important symptoms were nausea and vomiting, anorexia, weakness, dizziness, muscular pains particularly in the legs, slight diarrhœa, and melancholia. On examination there was emaciation, marked pallor of the mucous membrane and some abdominal distension with tenderness over the cæcum. The stools contained mucus and numerous eggs of *Trichuris*. After treatment by thymol the eggs disappeared from the stools and the patient made a rapid and permanent recovery. The second case was in a man who was suffering from a purulent pleuritis, with frequent epileptoid seizures, diarrhœa, muscular cramps, nausea, and restlessness. There were numerous eggs of *Trichuris* in the stools. The patient died and at autopsy the brain was normal, except for slight œdema of the meninges; the mucous membrane of the cæcum and lower portion of the small intestine was hyperaemic and contained *Trichuris trichiura*. Rippe considers *Trichuris* to be a pathogenic parasite and that infection with it must be considered in making diagnoses in conditions showing intestinal and nervous symptoms. The author found the eggs in 42.93 per cent of 184 cases of parasitic infection of the intestine in Russia.

Stiles (82) treats the pathogenicity of this parasite in a conservative manner. He recognizes a lack of symptoms in the majority of light infections, but on the other hand this author makes the statement "that severe infections do not produce injury is not in accordance either with probability or with recorded observations." Again he states that "several cases of severe anæmia have been recorded within recent years which were, apparently justly, attributed to heavy infections with whip-worms." Kahane (48) considers the parasite to be pathogenic. He states that it can cause ulceration of the intestinal mucosa and that the head may penetrate the walls of the bowel to the muscular coat.

Sir Patrick Manson (54) considers *Trichuris* infestation of no practical moment and that, so far as known, the parasites give rise to no serious pathologic lesions. Manson and Shipley (55) state that so far as is known *Trichuris* has very little pathologic significance, except when it perforates the intestine. These authors call attention to reports in which nervous symptoms have been associated with a high degree of infection. According to the same authors "several cases of anæmia (some fatal) are reported as caused by the presence of this parasite, the blood showing a fall in the number of red cells, marked poikilocytosis, and the appearance of nucleated red cells principally of the normoblastic type." Eosinophilia has been noted, but is not the rule.

Braun (13) considers *Trichuris* to be pathogenic. Seifert (78) states that from the evidence of recent years it is necessary to remove *Trichuris* from the list of commensal parasites and place it among those pathogenic for man, and that the prognosis is grave in some severe infections.

TRICHOCEPHALIASIS.

SYNONYMS.—Whip-worm infection, Trichuriasis.

DEFINITION.—An infection with whip-worms, usually confined to the large intestine, less frequently to the appendix and ileum and occasionally by perforation invading other parts of the body. In light infections there are no pathologic changes or symptoms of importance, but the heavier infections are characterized by anæmia, which may be very severe and even fatal and by nervous and gastro-intestinal symptoms.

ETIOLOGY.—Contributing causes: The ordinary causes which usually are considered as influencing infections probably play a less important rôle here than with many other diseases.

The geographic distribution of the infection is cosmopolitan. It is particularly prevalent in the Philippine Islands.

Sex.—Practically all authors recognize the much more frequent infection in females. Stiles and Garrison (83) in 3,457 cases found 12.13 per cent of infections in females and 10.08 in males.

Age.—It is generally recognized that infections are most prevalent during childhood and gradually grow less with advancing age.

Garrison, Ransom, and Stevenson (31) in 500 examinations noted this gradual decrease of infection with age, and Stiles and Garrison (83) found the following percentages of infection according to age: Under 15 years 13.01; from 15 to 30 years 10.09; from 31 to 50 years 8.65 and over 50 years 4.65. Garrison's (30) statistics in the Philippines were based entirely upon examination of specimens from adults and in our service the number of recorded examinations for children is too small to make statistics of value. However, we have seen several instances of infections in breast-fed infants, in one case the child being less than 3 months old.

Race and nationality.—In general it may be stated that the infection is more frequent among the colored races than it is in the Caucasian. Stiles and Garrison noted particularly the preponderance of infections in the American negro over that of the white races of the same country. Our records show a much higher percentage of infection in native Filipinos than is found in the foreign residents of the Caucasian race resident in Manila. In Stiles and Garrison's statistics, Irish born patients showed a higher percentage of infection than did any other nationality resident in the United States.

Residence.—The infection is more common in rural districts than it is in cities, no doubt largely because of the better methods of disposal of fæces in cities.

Occupation.—The infection is more frequent among soldiers, laborers and others whose manner of life and occupation brings them into contact with faecal infected substances.

Overcrowding with its usual insanitary surroundings may readily be inferred to be a factor in incidence.

Climate.—It is generally stated that the infection is more frequent in warm climates.

Gastrointestinal condition.—There are no statistics available, but it is probable that it is not a factor in the incidence of this infection.

Environment and mode of life.—After all is said regarding the influence of race, sex, etc., upon the prevalence of this infection, it is probable that the question

of infection or noninfection is almost entirely due to the mode of life of any given individual or community. In other words, it is a question of exposure, which in the vast majority of instances occurs only in those who are ignorant, or whose personal hygiene is otherwise very faulty. *In a sense, it seems to us, that the prevalence of whip-worm infections in a community is, to a certain extent at least, an index of the intelligence and cleanliness of its citizens.* This statement makes the assumption of what is probably a fact, that physiologic predisposition to infection with this parasite (except perhaps in children), if it exists at all, surely does so to a very minor degree.

Stiles and Garrison in discussing the question of physiologic predisposition in females state that "the relative frequency of a given parasite, or of helminthiasis in general, among males and females is apparently not so much a question of physiologic predisposition of the sex toward the parasite, as it is a question of the relative intimacy existing between the average daily life of the sex and the more concentrated infectious material."

In the case of whip-worm infections it seems to us that this statement applies not only to a question of sex, but that it explains many of the differences noticed under predisposing causes. Carrying this statement to its logical conclusion, we may say that race, sex, occupation, etc., as such, probably are not factors in the incidence of the infection, except to the extent in which they harmonize with the person's exposure to sources of infection under a given mode of life.

TRICHURIS.[2]

This *genus* is easily recognized by its peculiar whip-worm shape; the thick posterior portion and the long, slender anterior portion representing the stock and the lash respectively. The anterior extremity bears a small, round, mouth opening, behind which lies a small, funnel-shaped oral cavity. The long, thread-like œsophagus extends throughout the greater portion of the anterior, slender part of the body and enters the intestine which, in a folded course, reaches the posterior extremity of the body to find exit at a terminal anus in the female, and at the common genito-anal orifice in the male. The cuticle is striated transversely, except on the ventral surface where a longitudinal band, set with minute papillæ, extends from the anterior to the posterior extremities. The thick posterior portion of the body contains the intestine and the genital organs. In the female it is straight or slightly bow-shaped; in the male it is more or less closely coiled.

Male organs.—The testicle is single and consists of a long, coiled and folded tube which fills the greater part of the caudal portion of the body. A short, broad *vas deferens* turning abruptly caudad from the anterior end of the testicle enters a dilated portion of the genital tube (*vesicula seminalis*) which, in turn extends posteriorly to the cirrus or spicule. The spicule is single, highly developed and lies in a sheath or cirrus pouch, both sheath and spicule being protrusible for a considerable distance through the cloacal orifice.

Female organs.—The ovary is simple, tubular, with its extremity attenuated into a slender oviduct which enters the posterior end of the more expanded uterus. From the anterior extremity of the uterus, the slender, more or less coiled vagina leads to the genital pore which is situated on the ventral surface at about the junction of the slender, anterior and the thick, posterior portions of the body.

[2] These diagnoses of this genus and species, furnished by Dr. Philip E. Garrison of this laboratory, are taken from various authors, chiefly Leuckart, with the purpose of furnishing for medical men a description of the parasite, not too technical, but rather more detailed than that usually given in medical publications.

The ova are barrel-shaped and are characterized, as are those of *Trichosoma*, by the little colorless plug which closes each end of the brownish shell.

Habitat.—The cæcum and, less frequently, the colon and ileum of certain mammals.

Several species of *Trichuris* are recognized for different hosts, namely, *Trichuris trichiura* of man and apes; *T. affinis* of sheep, goats, and cattle; *T. depressinsculus* of the dog and fox; *T. unginculatus* of the rabbit; *T. crenatus* of the pig. These species are distinguished morphologically chiefly by the size and structure of the spicule and its sheath.

Trichuris trichiura (Linnæus, 1761).

Male, 40 to 45 millimeters long; spicule, 2.5 millimeters long; sheath covered with small, evenly distributed papillæ. Female 40 to 50 millimeters long, length of slender anterior portion to length of thick posterior portion as 3 to 2. Mouth opening simple and with appendages. Ova yellow to brown, 50 to 54 millimeters long by 23 millimeters broad; unsegmented in fresh fæces.

Habitat.—Cæcum and, less frequently, colon and ileum of man, apes, and the lemur.

Development.—Simple, without intermediate host. Embryo develops in water or moist earth in from 3 to 6 months or longer and will remain alive in its shell for several years unless ingested by its final host when it hatches and reaches sexual maturity in the cæcum after about 16 days.

The resistance to the action of physical and chemical agents shown by both the parent and eggs is remarkable. The parasite is the most difficult of any to destroy in the intestine. The ordinary anthelmintics as a rule have little or no effect upon it. The eggs may be frozen or completely dried without doing them permanent injury.

Modes of transmission.—*Direct:* Inasmuch as an intermediate host is not necessary for the propagation of this parasite, and inasmuch as contact with fæcal discharges is necessary in order to receive the eggs into the gastro-intestinal canal, infection proves a closer personal contact with filth than is true of many other parasitic infections. However, because of the slow development of the eggs under outside influences, immediate transmission from person to person or auto-reinfection can not occur.

Water when contaminated with fæcal discharges, directly or indirectly, may become a transmitter of infection, but because of the rapidity with which the eggs are sedimented in water, it would seem as if this were a much less important medium of transmission than it is generally considered to be.

Food, particularly uncooked vegetables, and to a less extent, fruits, are undoubtedly a means of transmission in a considerable number of instances. On two occasions during our study of the washing from fresh vegetables for the presence of amœbæ, ova of *Trichuris* were found in the sedimented material. The danger from fresh vegetables is great in those countries where human fæces are used as fertilizer for the vegetable gardens. Although this procedure is a violation of the law, it is certainly practiced to a considerable extent by Chinese gardeners in the Philippines, as it is in several other oriental countries.

Insects, such as flies, water bugs, roaches, etc., may be factors of some importance as transmitters of the infection by mechanically carrying it from infected fæces to food and water supplies.

Soil, in localities where methods of sewage disposal are primitive, surely becomes infected and because of the great viability of the eggs in earth, such places become a menace in many obvious ways. So-called clay eaters would be especially liable to infection.

Air, clothing, dust and other substances may be considered as possible factors in the spread of the infection in a mechanical way.

Animals.—Several authors have called attention to the possible influence of animals in the transmission of this infection. When it is remembered that some domestic animals used as pets may harbor this parasite, and that from their habits several of them are likely to have the eggs in their hair, it would seem that animals might be expected to be occasional transmitters of the infection, particularly among children.

Pathogenicity.—The largely accepted commensalism of this parasite is not in accord with the majority of the carefully reported observations (see literature review), nor is it borne out by study of our cases. Belief in the harmless character of *Trichuris* is based largely upon two observations: first, the great, general prevalence of the parasite without symptoms, and second, no satisfactory explanation of the nature of its pathogenic action has been offered.

If we examine the first of these in comparison with some other parasitic infections we find it to be a difference in *degree only* and not a peculiar condition. For example, *Uncinaria* are accepted as pathogenic parasites and yet in our service more than 90 per cent of individuals harboring the worm show no symptoms whatever of the infection. The first symptoms of infection with amœbæ may be those of peritonitis from a perforating ulcer.

Strongyloides stercoralis is now generally accepted as a pathogenic parasite, and yet in the vast majority of instances no symptoms are produced by its presence, and similar statements may be made regarding *Balantidium coli, Dibothriocephalus latus* and a number of other parasites usually accepted as pathogenic.

The percentage of cases infected with *Trichuris* without symptoms probably is greater than for any of the above-named parasites, but infections do occur which produce symptoms and the degree of difference does not appear to be great enough to justify it as an argument for the commensalism of *Trichuris*.

As to the second of the two views, that no satisfactory explanation of the nature of the infection has been made, we may at least in part answer that here the difference from some of the other parasites is also one of degree. No satisfactory explanation of the exact manner of the production or character of the lesions, produced by the majority of the intestinal parasites of man, has been offered.

The pathogenic character of *Trichuris* has been explained by considering it to be a blood sucker; by believing it to produce a hæmolytic substance such as is also acredited to hook-worms; by the formation by it of a toxin; by mechanical action and by its producing perforations which open a way for the entrance of bacteria. None of these explanations is supported by sufficient data to justify its acceptance. The majority of the repeated statements that the parasite is a blood sucker are credited

to Askanazy. What Askanazy actually said was that he had isolated "an iron containing pigment from the epithelial cells of the intestine of the parasite." So far as we know, neither blood nor other positive evidence of blood absorption has been found in *Trichuris*.

These parasites do attach themselves to the intestinal wall during the life of the host by transfixation and direct penetration by the head end, and that these locations may show bright, hyperæmic spots and become eroded or even ulcerated, is proved beyond doubt. However, the lesions formed by this action are not as a rule sufficient to explain the symptoms, unless we accept some peculiar hæmolytic or other toxic action of the parasite. It is certain that the mechanical loss of blood from these infections is not sufficient to account for the symptoms and a definite toxic action has not been proved. The nervous symptoms and some of the gastro-intestinal ones might be explained from the mechanical action of the parasites by an hypothesis similar to that of Mayo's embryologic formation used for the explanation of similar symptoms in certain cases of chronic appendicitis and gall bladder diseases. However, this is only a hypothesis and does not account for the severe anæmia, which after all is the important manifestation. The nature of the action of these parasites in producing disease has not been satisfactorily explained, but this should not be accepted as proof of their commensalism.

Pathology.—The general pathologic changes in fatal cases of trichocephaliasis are severe, secondary anæmia. The special conditions are the presence of the worms and certain changes at their points of attachment to the mucous membrane of the bowel or appendix. Several changes have been described by different authors, which in the main are hyperæmia of the mucosa with hæmorrhagic points; erosions and superficial ulceration of the mucous membrane, surrounded by areas of cell infiltration; and in some instances deep ulceration or other inflammatory reactions extending to the muscular coat of the bowel. Several observers have shown that the worms are attached to the mucous membrane during the life of the patient. They attach themselves by transfixing a fold of mucous membrane, by penetrating the glandular follicles, and by direct perforation, the head being embedded in the deeper layers of the bowel wall. After the death of the patient, the worms detach themselves and at autopsy are usually found free in the bowel. Several authors have demonstrated this direct attachment to the intestine. Corroborative evidence is furnished by the fact that they do not appear in the discharges even after violent purgation. The worms were not passed in a number of cases of successful treatment, when the disappearance of eggs from the fæces showed that they had been destroyed. In these instances we can only assume that the parasites have been killed and remain attached to the mucous membrane of the intestine after death.

Incidence.—This is one of the most frequent of all parasitic infections of man and Garrison has shown it to be the most prevalent in the Philippine Islands.

Some of the statistics in percentages are as follows:

French & Boycott (1905), London 7.8; (1887), Erlangen, 11.1; (1887), Erlangen (Insane), 100. Leukart (1887), Dresden, 2.5; (1887), Kiel, 32.2; (1879), Dublin, 90. Cobbold (1879), Greenwich, 69; (1877), Paris, 50. Davaine (1877), Naples, 100. Blanchard (1889), Bale, 23.6. Garrison, Ranson, and Stevenson

(1903), United States (Insane), 16.8. Heisig (1893), Germany, 45.21. Gubareff (1896), Russia, 43. Boycott (1904), England (Cornwall), 38.78. Grechaninoff (1890), St. Petersburg, 26.41. Sievers (1887), Kiel, 19.81. Roth (1877–1880), Bale, 26.67. Gribbohm (1872–1877), Kiel, 32.20. Heller (1872–1875), Kiel, 30.60. Müller (1862–1873), Erlangen, 11.11. Müller (1852–1862), Dresden, 2.57. Stiles and Garrison (1906), United States (Insane), 7.72. Cima (1893 and 1896), Italy, 37.27. Anæmia Commission (1904), Porto Rico, 7.27. Daniels (1901), Africa, 2.79. Fearnside (1900), India, 6.95. Dobson (1893), India, 4.40. Garrison (1908), Philippine Islands, 59.

Some of the figures included in the series are from examinations of only a few cases or were compiled under conditions which prevented the findings being properly considered as an index of the prevalence of the infection in the general population of the community in which the examinations were made.

Garrison's large and carefully worked out statistics show the highest percentage of infection among the general population as yet reported for any country.

The *incubation* period in this disease would be the time for development of the ingested embryo to maturity which, to judge from the few available data, is more than four weeks.

Symptoms.—We recognize for convenience in discussion *mild* and *severe* intestinal trichocephaliasis and trichocephaliasis of the *appendix*. In general the symptoms may be said to depend upon the severity and location of the infection. In *mild* cases of intestinal infection there are no noticeable symptoms and the patient may harbor the parasites in considerable numbers for a long time without inconvenience. In *severe* infections the symptom complex is largely that of a severe secondary anæmia, with pronounced nervous symptoms and some gastro-intestinal disturbance. The early symptoms vary somewhat in different cases. Usually there is weakness, shortness of breath, nausea, vomiting, diarrhœa, cramps in the muscles, nervousness, insomnia, and a gradually developing anæmia. The condition is as a rule progressive, with accentuation of the principal symptoms as the disease advances.

Circulatory system.—Weakness, with shortness of breath and palpitation of the heart are often early symptoms and become more marked as the disease progresses. Evidences of anæmia, such as pallor of the mucous membrane, œdema, and hæmic murmur of the heart have been noted in several cases. As a rule the spleen is not enlarged and in fact, it may be smaller than normal. The blood picture is largely that of secondary anæmia. There is a decrease in the number of red cells, with poikilocytosis, but nucleated cells are not as a rule present. The number of the leucocytes remains approximately normal and the differential count usually shows a slight relative increase in mononuclears. Slight eosinophilia has been noted, but most observers agree that the eosin cells are not increased and in our cases eosinophilia was absent. No eosin cells were ever found in our Case No. 1. The hæmoglobin is reduced and may be very low before death.

The symptoms in the *respiratory system* may consist of some dyspnœa and cough, when severe anæmia is present.

The *alimentary system* as a rule suffers rather severely. Nausea and vomiting are frequent and annoying symptoms, anorexia and dyspeptic symptoms are also often encountered, although the appetite may remain good throughout the course of the disease. The mucous membranes are pale, the tongue often large and flabby and in two of our cases it showed a dark band down the center, similar to that which has been described for uncinariasis. There may, or may not, be tenderness and pain in the gastric region and in the two cases where it has been recorded, examination of the stomach contents showed nothing abnormal. More or less abdominal discomfort or even pain has been a feature of the majority of

cases which have been described and diarrhœa has been present in most of them. The character and number of the intestinal discharges varies, ranging from only one or two stiff, pasty stools in twenty-four hours, to the severest form of diarrhœa, even with bloody discharges. Again, the bowel movements may appear normal for considerable periods of time. Whatever the number and character of the bowel movements, tenesmus and other symptoms indicating involvement of the lower bowel are rare. Stool examinations demonstrate the eggs of the *Trichuris;* at times but few, and again large numbers may be seen in a single field of the microscope. The *urinary*, *reproductive*, and *locomotor systems* do not, as a rule, show any characteristic symptoms. Muscular cramps, particularly in the calves of the legs, have been repeatedly observed. The urine is usually normal. The *cutaneous system* shows the changes due to anæmia. There may be more or less œdema of dependent parts and even anasarca may develop. The superficial lymphatics are not enlarged.

The *nervous* symptoms are usually quite noticeable and consist of mental and nervous depression which may be melancholic in character; there is often great restlessness, headache, and sometimes insomnia. Objectively nothing of importance is found.

Among the symptoms of the *special senses* aphasia is frequent and it is difficult to explain. Tinnitus aurium and even partial deafness have also been noted by several observers.

Appendiceal trichocephaliasis has not heretofore received much attention. Metchinkoff urged its importance and reported a convincing case in detail. Very little is given about the clinical findings, but *a priori* there is no reason to assume that they would be materially different from appendicitis of other etiology.

Trichocephaliasis of animals has been observed by several authors.

Analysis of symptoms.—As pointed out by Becker, the symptoms in the majority of cases of this disease fall into three groups:

(1) Those of the blood, (2) nervous and (3) gastro-intestinal symptoms.

These are present to a greater or less extent in all cases of severe infection. The blood changes and the anæmia point positively to what we recognize as a secondary type of the latter. The absence of eosinophilia is quite generally recognized and is to be noted. French and Boycott paid particular attention to this condition and failed to find eosinophilia in 26 cases which they examined. In ours, eosinophilipænia was the rule, and in Case No. 1 no eosinophile cells were ever found after repeated blood examinations.

The nervous symptoms are difficult to explain, particularly the mental and nervous depression and the aphasia which has been so frequently noted, and it is probable that the nausea and vomiting are to a considerable extent of nervous origin.

Even the intestinal symptoms are difficult to explain as a result of the pathologic findings, certainly they are not to be considered as being due to the direct loss of blood caused by the mechanical action of the worms. Marsasca and others believe that the latter secrete a special toxin, and although the entire clinical picture supports such a view, it has not been demonstrated. Bacteria and other secondary invaders entering through the injured mucosa probably account for some of the symptoms and explain those cases having septicæmia. However, the manner of the production of the symptoms remains unexplained.

Diagnosis.—While the clinical picture in some of these cases is quite suggestive, diagnosis can only be determined by finding the eggs of the parasite in the stools. Not every patient having eggs in the stools will show symptoms of the infection, and for clinical purposes only a combination of some of the clinical phenomena and

the presence of eggs should be considered as a diagnosis from a practical therapeutic standpoint. The differentiation from most of the other recognized types of anæmia is readily made by the blood picture. Severe infections are only differentiated clinically from similar cases of uncinariasis by stool examinations and by the greater prevalence of eosinophilia in the hook-worm infections.

PROGNOSIS.—Certain light infections do not appear to do recognizable harm, even when the parasites remain for a long time in the bowel. The prognosis in severe infections is grave, largely because of the difficulty with which the parasites are destroyed. Barth, Pascal, Sandler, Moosbrugger and others have reported fatal cases. The disease may continue for a long time, and, on the other hand, as in our Case No. 1, the course may be fairly acute and its duration from the development of clinical symptoms short.

PROPHYLAXIS.—The prevention of this disease consists in personal hygiene and in public health measures. Because of the simple life cycle of the parasite and the long viability of the eggs under outside influences, personal hygiene, by avoiding the introduction of unsterilized substances into the mouth, makes protection fairly safe even in infected zones.

The hygienic side of the problem requires a safe and practical method for the disposal of fæces. This is very important, not only because of the presence of whip-worms, but of others as well. This subject has been covered by Garrison (30).

TREATMENT.—Whip-worms are exceedingly difficult to destroy in the intestine. In this respect out experiences have been in conformity with the general literature of the subject. However, several authors have had more or less success with various anthelmintics, *and in most cases where it has been possible to destroy the worms, rapid and complete recovery of the patient has followed.* Enemas of solutions of benzine have been the most successful in the hands of several observers, while others have reported success by the use of calomel, thymol, naphthol, santonin, or garlic. We have used thymol, the eucaliptus oil, chloroform, and the castor oil mixtures employed for uncinariasis, but without very much success. The benzine treatment has only just come to our notice, but to judge from the reports in the literature it deserves a more extended trial. Schiller (quoted by Sandler(74)) used thymol by mouth and benzine solutions as enemas and secured 2,000 worms from one case. This is remarkable because of the fact that the worms were passed. The killed worms in the experience of most authors do not pass from the bowel and their destruction is only made evident by the disappearance of the eggs from the stools. Other treatment of the disease consists in general hygienic measures with attempts to improve the nutrition of the patient.

BIBLIOGRAPHY.[3]

ANDRIKIDIS. *(1) Étude des troubles morbides attribuables au trichocéphale de l'homme, Paris, (1906), 119 pp.

ASHFORD, B. K., KING, W. W., and IGARAVIDEZ, P. G. *(2) Report of the Commission for the study and treatment of anemia in Porto Rico. Authorized by act of the Legislative Assembly. San Juan, (1904), 311 pp., 7 pl.

ASKANAZY, M. (3) Der Peitschenwurm ein blutsaugender Parasit. *Deutsches Arch. f. klin. Med.*, (1896), **57**, 104–117, 1 pl.

BARTH. *(4) *Ref.:* von Valleix, Guide de médicin practicien, Paris, (1845), **6**, 98.

[3] Unmarked references have been verified in the library of the Bureau of Science. References marked † have been partially verified—in most cases the original article has not been available, but reviews and abstracts have been verified as cited. References marked * have not been verified.

Becker, E. (5) Ueber die durch Trichocephalus dispar verursachten Krankheitszustände. *Deutsche med. Wchnsch.*, (1902), **28**, 468–470.

Bellingham, O'Bryan. †(6) On the frequency of the presence of trichocephalus dispar in the human intestines. *Dublin Journ. Med. Sci.*, (1837–8), **12**, 341–347; *also abstr.: Am. Journ. Sci. & Arts.* (1838), **34**, 10–11.

Berguin, J. E. *(7) Essai sur le traitement à employer contre les principaux helminthes de l'homme (ascarides, oxyures, trichocéphales). Paris, (1881), 64 pp.

Blanchard, Raphael. †(8) Trichocéphale. *In his:* Dictionnaire encyclopedique des sciences médicales (1887), III, **18**, 171–179; *review by* M. Braun: *Centralbl. f. Bakt. etc.*, (1888), **3**, 441.—(9) Trichocephalus hominis. *In his:* Traité de zoologie médicale, (1889), **1**, 774–787.—*(10) *In:* Bouchard's Traité de pathologie générale, (1895–7), **2**, 760.

Boas. (11) Demonstration eines Falles von Trichocephalus dispar. *Deutsche med. Wchnsch.*, (1895), **21**, Ver.-Beil., 97.

Boycott, A. E. (12) Further observations on the diagnosis of ankylostoma infection with special reference to the examination of the blood. *Journ. Hyg.*, (1904), **4**, 437–479.—*See also* French and Boycott.

Braun, Max. (13) Trichocephalus. *In his:* Tierischen Parasiten des Menschen, (1903), 3. ed., 278–279; (1908), 4. ed. 311–313.

Burchardt, Max. (14) Schwere durch Peitschenwürmer bedingte Erkrankung. *Deutsche med. Wchnsch.*, (1880), **6**, 633.

Cima, Francesco. *(15) Sulla elmintiasi dei bambini. *Pediatria*, (1893), **1**, 39–48.—*(16) Sull' elmintiasi dei bambini. Nuevo ricerche ed osservazioni cliniche. *Ibid*, (1896), **4**, 303–312.

Cobbold. *(17) Parasites, (1879), 179. *Cited by* French and Boycott (28).

Daniels, C. W. (18) Notes on intestinal worms in natives of British Central Africa. *Journ. Trop. Med.*, (1901), **4**, 199.

Davaine, C. †(19) Recherches sur le développement et la propagation du trichocéphale de l'homme et de l'ascaride lombricoïde. *Journ. physiol. de l'homme*, (1858), **2**, 295–300; also: *Compt. rend. Acad. sci.*, (1858), **46**, 1217–1219; *also transl.: Boston Med. & Surg. Journ.*, (1859), **59**, 157–159.—(20) Trichocéphale de l'homme. *In his.:* Traité des entozoaires, (1877), 2. ed., 206–210; 223–224.

Dobson, Edwin F. H. †(21) Indian helminthology. (Abstract of a series of articles in *Indian Med. Gaz.*, (1892), **27**, 354–357; (1893), **28**, 1–4; 40–43; 68–72; 262–267). *Brit. Med. Journ.*, (1893), **2**, 807–808.

Eberth, I. *(22) Beiträge zur Anatomie und Physiologie des Trichocephalus dispar. *Ztschr. f. wiss. Zool.*, (1860), **10**, 233; 383.

Emerson, Charles Phillips. (23) Trichocephalus trichiuria. Trichocephalus dispar. Trichiuris trichiura. *In his:* Clinical diagnosis, Philadelphia, (1906), 384–385.

Erni, H. (24) Trichocephalus dispar. Ein Beitrag zur Beri-Beri-Frage. *Berl. klin. Wchnsch.*, (1886), **23**, 614–616.

Fanyo, F. *(25) Infection with trichuris trichiura (trichocephalus dispar) with the report of a case. *Illinois Med. Journ.*, (1903–4), n. s., **5**, 765–767; (discussion), 683–685.

Fearnside, C. J. (26) A discussion on ankylostomiasis. *Brit. Med. Journ.*, (1900), **2**, 541–542.

Federolf, A. K. †(27) Ein schwerer Fall von Trichocephaluswirkung, (Russisch). *Voyenno-med. Journ.*, (1897), **189**, 430–433; *review by* Ucke: *Centralbl. f. Bakt.*, etc., (1898), **23**, 612.

FRENCH, H. S., and BOYCOTT, A. E. (28) The prevalence of Trichocephalus dispar, *Journ. Hyg.*, (1905), **5**, 274–279.

FRIEDRICH, ARTHUR. (29) Ueber die Häufigkeit der thierischen Darmparasiten bei Erwachsenen in München. *Münchener med. Wchnsch.*, (1887), **34**, 935–936; 955–956; *review by* Rockwitz: *Centralbl. f. Bakt., etc.*, (1888), **3**, 308–309.

GARRISON, P. E. (30) The prevalence and distribution of the animal parasites of man in the Philippine Islands, with a consideration of their possible influence upon the public health. *This Journal*, (1908), **3-B**, 191–209.—*See also* Stiles and Garrison.

GARRISON, P. E., RANSOM, B. H., and STEVENSON, E. C. (31) A statistical study of the intestinal parasites of 500 white male patients at the United States Government Hospital for the Insane. *Bull. Hyg. Lab. U. S. Pub. Health & Marine Hosp. Service*, no. 13, (1903), 1–13; review: *Med. Rec.*, (1903), **64**, 222.

GIBSON, DANIEL. (32) On a case of paralysis with loss of speech from intestinal irritation. *Lancet*, (1862), **2**, 139.

GIRARD, J. (33) Rôle des trichocéphales dans l'infection de l'appendice iléo-cæcal. *Ann. Inst. Pasteur*, (1901), **15**, 440–444, 1 pl.

GOEZE, J. A. E. *(34) Naturgeschichte der Eingeweidewürmer, (1872).

GRASSI, B. (35) Trichocephalus und Ascarisentwicklung. *Centralbl. f. Bakt., etc.*, (1887), **1**, 131–132.

GRECHANINOFF, V. N. *(36) K statistikle helminthiasis. (Statistics of helminthiasis), St. Petersburg, (1890), 39 pp.

GRIBBOHM, HARTWIG. *(37) Zur Statistik der menschlichen Entozoen, *Inaug. Dissert.*, Kiel, (1877), 12 pp; *also Schrift. d. Univ. zu Kiel*, (1877), **24**.

GUBAREFF, P. A. *(38) (Several statistics concerning the frequency of helminthiasis in the governments Tiflis and Kutais) (Russian text). *Vratch*, (1896), **17**, 269–271.

GUIART, J. (39) Le trichocéphale et les associations parasitaires. *Compt. rend. Soc. biol.*, (1901), **53**, 307–308.—(40) Rôle du trichocéphale dans l'etiologie de la fièvre typhoïde. *Semaine méd.*, (1904), **24**, 333.

GUINARD. *(41) Appendicite à répétition. Operation à froid. Ver intestinal vivant (trichocéphale mâle) dans l'appendice. *Bull. et mém. Soc. chir.*, (1900), **26**, 1009–1010.

HASSELL, ALBERT. *See* Stiles and Hassell.

HAUSMANN, TH. *(42) Zur Symptomatologie und Therapie der durch den Trichocephalus dispar verursachten Gesundheitsstörungen. *St. Petersb. med. Wchnsch.*, (1900), n. s., **17**, 301–304.

HEINE, PAUL. (43) Beiträge zur Anatomie und Histologie der Trichocephalen, insbesondere des Trichocephalus affinis. *Centralbl. f. Bakt.*, etc., (1900), **28**, 779–787; 809–817, 2 pl.

HEISIG, OSWALD. †(44) Beitrag zur Statistik menschlicher Entozoen. *Inaug. Dissert.*, Griefswald, (1893), 28 pp.; *review by* E. Roth: *Centralbl. f. Bakt., etc.*, (1894), **15**, 326–327.

HELLER, ARNOLD. *(45) Darmschmarotzer. *In:* Ziemssen's Handbuch d. spec. Path., (1876), **7**, pt. 2, 559–664; *also transl.:* Cycl. prac. med., (1876), **7**, 667–778.

HEMMETER, JOHN C. (46) Trichotrachelidae (Leuckart) (Whip-worm). Trichuris trichiura (old name, Trichocephalus dispar). *In his:* Diseases of the intestines, (1902), **2**, 580–582.

HUBER, J. CH. (47) Trichocephalus hominis. *In:* Twentieth century practice of medicine, (1896), **8**, 590–593.

IGARAVIDES, P. G. *See* Ashford, King and Igaravidez.

KAHANE. *(48) *Cor.-Bl. f. schweiz. Aerzte*, (1907). *Cited by* Seifert (78).

KESSLER, D. A. (49) Thierische Darmparasiten; *review by* Besser: *Centralbl. f. Bakt.*, etc., (1889), **5**, 522.

KING, W. W. *See* Ashford, King and Igaravidez.

KLEBS, E. (50) Trichocephalus dispar (Peitschenwurm). *In his:* Handbuch d. path. Anatomie, (1869), **1**, Abt. 1, 313–315.

KÜCHENMEISTER. *(51) Die Parasiten, Leipzig, (1855), 240.

LEUCKART, RUDOLF. †(52) Die menschlichen Parasiten, (1876), **2**, Trichocephalus, 463–465; Trichocephalus dispar, 465–509; *also transl.* by Hoyle, (1887), **1**, 151.

LUTZ, ADOLF. (53) Trichocephalus dispar. *Centralbl. f. Bakt., etc.*, (1888), **3**, 777–778.

MANSON, PATRICK. (54) Trichocephalus trichiuris. *In his* Tropical diseases, (1908), 4 ed. 706–710.

MANSON, PATRICK, and SHIPLEY, A. E. (55) Trichocephalus trichiurus. *In:* Allbutt's System of medicine, (1907), **2**, pt. 2, 906–908.

MÉGNIN, P. (56) Du rôle des ankylostomes et des trichocéphales dans le développement des anémies pernicieuses. *Compt. rend. Soc. biol.*, (1882), VII, **4**, 172–177.

METCHNIKOFF, E. (57) Sur l'appendicite; note helminthologique sur l'appendicite. *Bull. Acad. méd.*, (1901), III, **45**, 301–309.

MILLER, FRANK H. (58) Haemorrhagic colitis of the dog due to infection with trichocephalus depressiusculus (true whip worm). *Am. Vet. Rev.*, (1904–5), **28**, 722–729, 1 pl.

MOOSBRUGGER. *(59) Ueber Erkrankung an Trichocephalus dispar. *Med. Cor.-Bl. d. württemb. ärztl. Ver.*, (1890), **40**, 193–196.—(60) Ueber Trichocephaliasis. *Münchener med. Wchnsch.*, (1895), **42**, 1097–1099.

MORGAGNI, J. B. *(61) Epistolarum anatomicarum duodeviginti ad scripta pertinentium cel. viri. Ant. Mar. Valsalvae, Venetiis, (1740), Voir III, epistola XIV, § 42, p. 45. (*Cited by* Blanchard (9) in this form).

MORSASCA. (62) Cloro-anemia da trichocephalus dispar. *Centralbl. f. innere Med.*, (1897), **18**, 551.

MÜLLER, KARL. *(63) Statistik der menschlichen Entozoen. *Inaug. Dissert.*, (1874), 31 pp.

MYER, J. S. *(64) A case of double intestinal parasitism (trichocephalus dispar and ascaris lumbricoides) with enlargement of the spleen. *Quart. Bull. Med. Dept. Wash. Univ.* St. Louis, (1906–7), **5**, 132–137.

OSLER, WILLIAM. (65) Trichocephalus dispar. *In his:* Practice of medicine, (1906), 6. ed., 51.

OSTROVSKI, A. L. *(66) Le rôle du trichocephalus dispar dans la pathologie, *Vratch*, (1900), **21**, 1206–1207.

PASCAL. (67) *Ref.:* Leuckart (52).

POLEDNE, W. (68) Schwere Enteritis verursacht durch Trichocephalus dispar im Darme. *Wien. med. Wchnsch.*, (1906), **56**, 1821.—(69) Taenia solium, Ascaris und Trichocephalus dispar gleichzeitig im Darme. *Ibid*, 1822–1823.

PRUDDEN, T. M. *(70) Trichocephalus dispar. *Proc. New York Path. Soc.*, (1892), 82.

RAILLIET, A. *(71) *Bull. Soc. centr. méd. vétér.*, (1884), 449.

RANSOM, B. H. *See* Garrison, Ransom and Stevenson.

RIPPE, E. (72) Einiges über den Trichocephalus dispar. *St. Petersb. med. Wchnsch.*, (1907), **32**, 1–4.

ROEDERER, J. G. *(73) Nachr. v. d. Trichiuriden. *Göttinger gelehrte Anzeigen* (1761).

SANDLER, ARON. (74) Trichocephaliasis mit tödlichem Ausgang. *Deutsche med. Wchnsch.*, (1905), **31**, 95–98.

SCHILLER. *(75) *Beitr. klin. Chir.*, (1902), **34**. *Cited by* Seifert (78).

SCHRANK, F. v. P. *(76) Eingeweidewürmer, München, (1788).

SCHULTZE, W. (77) Ueber Trichocephalus dispar. *Deutsche med Wchnsch.*, (1905), **31**, Ver.-Beil., 1487.

SEIFERT, OTTO. (78) Trichocephalus trichiurus. *In* Braun's Tierische Parasiten des Menschen, (1908), 551–553.

SHIPLEY, A. E. *See* Manson and Shipley.

SIEVERS. *(79) Schmarotzer-Statistik aus den Sections-Befunden des pathologischen Instituts zu Kiel vom Jahre 1877 bis 1887, Kiel, (1887), 24 pp.

SMITH, ALLEN J. (80) Trichiuris trichiura. *In Tyson's* Practice of medicine, (1903), 3. ed., 1198; (1906), 4. ed., 1242.

SOMMER, H. OTTO. *(81) Fecal examination of 36 patients in the Children's Hospital of Washington, D. C., for ova of intestinal parasites. *Southern Journ. Homeop.*, (1895), **13**, 353–354.

STEVENSON, E. C. *See* Garrison, Ransom and Stevenson.

STILES, CH. WARDELL. (82) Trichocephaliasis. *In* Osler's Modern medicine, (1907), **1**, 602–605.

STILES, CH. WARDELL, and GARRISON, P. E. (83) A statistical study of the prevalence of intestinal worms in man. *Bull. Hyg. Lab. U. S. Pub. Health & Marine Hosp. Service*, no. 28, (1906), 1–77.

STILES, CH. WARDELL, and HASSELL, ALBERT. (84) The determination of generic names and a list of roundworm genera, with their original and type species. *Bull. U. S. Bur. An. Ind.*, no. 79, (1905), 1–150.

STRONG, R. P. (85) Animal parasites. Circ. on trop. diseases, no. 1, Manila, (1901), 25–26.

VIX. *(86) *Ztschr. f. Psychiatrie*, (1860), **17**, 305. *Cited by* Seifert (78).

WALTHER. *(87) Appendicite et trichocéphales. *Bull. et mém. Soc. chir.*, (1905), n. s. **31**, 355.

WARD, HENRY B. (88) Trichocephalus. *In:* Buck's Reference handbook of the medical sciences, (1903), **6**, 216–217.

WICHMANN, J. C. *(89) Ueber das Verhalten des Trichocephalus zur Darmschleimhaut, *Inaug.-Dissert*, Kiel, (1889).

ZINN, W., and JACOBY, MARTIN. (90) Ueber das regelmässige Vorkommen von Anchylostomum duodenale ohne secundäre Anämie bei Negern, nebst weiteren Beiträgen zur Fauna des Negerdarmes. *Berl. klin. Wchnsch.*, (1896), **33**, 797–801.

EDITORIAL.

NEOPLASMS AMONG THE FILIPINOS.

It has been variously stated by different writers that new growths and more especially malignant tumors, are not only less common than in temperate climates, but even rare in many tropical countries. There was a more or less well-defined idea prevalent among the profession in Manila that in the native inhabitants of the Philippines neoplasms are uncommon, but no complete records were available bearing on the point.

Malignant disease as a cause of death occurred twice among the first series of 100 necropsies at the Philippine Medical School; 2 cases of carcinoma of the greater curvature of the stomach, and 1 of the cheek and antrum.

Dudley,[1] writing on the prevalence of cancer in the Philippines after an experience of ten years spent in the various islands, believes that cancer exists in the Philippines to a greater extent than in the United States, and that the statistics available on malignant and nonmalignant growths are very incomplete.

The following figures illustrating the frequency of neoplasms in Dr. McDill's surgical clinic at the St. Paul's Hospital may be taken as fairly typical, as the clinic draws on all classes of free patients, the material coming not only from the city of Manila, but from the surrounding and often distant provinces.

The record of cases coming to operation from September 5, 1907, up to December 19, 1908, shows a total of 377. Sixty-five of these cases, or 17.4 per cent, were operations for new growths, 37 benign and 28 malignant.

Of the benign neoplasms, cystic ovarian disease leads in frequency, there being 16 cases, of which 4 were double. Of uterine fibroids there were 7 cases, the most frequent being the sub-serous variety. There were 5 cases of fibromata of the skin and 2 each of osteomata of the head and intracanalicular myxomata of the breast. There was but one case of adenoma of the breast; one dentigerous cyst of the lower jaw; one cyst of the neck (thyroglossal duct); one cyst of the anterior wall of the vagina, and one epulis.

[1] The Prevalence of Cancer in the Philippine Islands, *Journ. Amer. Med. Ass.* (1908), **50**, 1663 to 1665.

Of the 28 cases of malignant growths there were 13 sarcomata, 12 carcinomata, and 3 of Hodgkin's disease. The sarcomata involved the scalp, testicle, femur, and dorsum of the foot twice; the forearm, eye, parotid, tibia, and skin once.

There were 6 cases of carcinoma of the stomach. The breast was carcinomatous in 2 cases, the pancreas, uterus, liver, and skin one each. The 3 cases of Hodgkin's disease all showed involvement of the supraclavicular and axillary glands.

PHILIP K. GILMAN.

REVIEW.

A Text-Book of the Principles of Animal Histology. By Ulric Dahlgren, M. S. and William A. Kepner, A. B. Cloth. Pp. xiii+515. Price, $3.75 net. New York: The Macmillan Company, 1908.

This work of more than 500 pages, containing nearly 500 illustrations of which about 300 are original, has had such a flattering reception and has been so favorably reviewed that it is unnecessary further to discuss its value. The authors state that the book "is intended to be a work that teaches general principles and teaches histology as a pure science and for its own sake. It is believed that it will serve as a broad foundation for future studies of morphology and embryology as well as for the medical studies." This object is well fulfilled. Rich fields heretofore untouched by any text-book, such as the tissues that produce electricity, light, gases, etc., are opened, offering the results of original research work on these subjects.

The illustrations are not only largely original, but they explain the text by their clearness and detail. Many of them are diagrammatic, thus furnishing an ideal method of instruction without sacrificing reality. In this way the structure of protoplasm is made plain, chromatin changes become distinct, the amplification of body surface is illustrated, the arrangements of muscles' cells is explained, the relation of absorptive surface is indicated, and many other obscure points are made clear. A bit of muscle fiber shows relaxation in the segment at one end and successive stages of contraction in all the other segments, and this is only one of many unusual illustrations.

The comprehensive character of the work is made clear by the following list of tissues, to each of which a chapter is devoted:

Epithelium.
Supporting and connecting tissues.
Tissues of motion.
Electric tissues.
Tissues of light production.
Tissues of heat production.
Circulatory tissues.
Nerve tissues.
Pigment tissues.
Alimentary tissues.
Ductless glands.
Tissues of respiration.
Gas-secreting tissues.
Excretory tissues.
Protective tissues.
Tissues of reproduction.
Nidamental and erectile tissues.
Nourishing membranes.

Improvement in style and in grammatical structure are to be desired and will doubtless be found in future editions of the work, of which it is to be hoped there will be many. If references are given they should be in greater number and form a complete bibliography. However, such a wide range of subjects is covered that an undertaking of this kind is scarcely feasible.

The book is not altogether suitable for beginners, but it is the best that I have seen and I most heartily recommend it to students preparing for medical work.

R. B. B.

INDEX.

[NOTE.—The heavy-faced type is used in this index only in cases where the species is new, that is, pertaining to this year.]

Actinomyces, 458.
Acute bacillary dysentery, 216.
Acute dysentery, 71.
African tick fever from the examination of the blood, the diagnosis of, 231; animal experiments with the African spirochæta, 237–242; with the American spirochæta, 237–242.
Agchylostoma, number of cases of at Bilibid Prison during 1908, 71; present in the Philippine Islands, 195.
Agglutinin, agglutinoid, and agglutinable substances and their quantitative relationships, 47.
ALBERT, JOSÉ, A case of infantile beriberi with autopsy report, 345.
Alimentary canal, lesions of the, 216.
Amœbæ, per cent of in water examined in the Philippine Islands, 122; presence among soldiers, 262; prevalence and distribution in the Philippine Islands, 191, 195, 200, 201; results of 300 examinations of fæces with reference to the presence of, 417.
Amœbic dysentery, 71; 212; 123; 188.
Animal and bacterial symbiosis, 84.
Animal parasites of man, prevalence and distribution in the Philippine Islands, 191–209.
ASHBURN, P. M., & CRAIG, CHARLES F., A comparative study of tsutsugamushi disease and spotted or tick fever of Montana, 1.
ATKINSON, J. M., Plague procedure in Hongkong, 39.
Atrophy, per cent in 100 cases of acute, yellow, 218.
Autopsies (100) at the Philippine Medical School, 211; diseases of the kidneys, 215; disease of the nervous system, 220; lesions of the alimentary canal, 216; of the liver, 217; of the spleen, 218; lymphatic system, 219; carcinoma, 219; peritoneum, 218; pneumonia, 213; tuberculosis, 212.
Bacillary dysentery, 31.
Bacillus cabildonis, 125.
Bacillus coli, 47.
Bacillus dysenteriæ, 47.
Bacillus paratyphosus, 47.
Bacillus pyocyaneus, 47.
Bacillus typhosus, 47.
Bacterial symbiosis, 79.
Balantidium, present in the Philippine Islands, 195, 201; 71.
BANKS, CHARLES S., A mosquito which breeds in salt and fresh water, 335; Discussion on the paper by Dr. T. Kitajima (151), 188.
Beriberi, autopsy report of infantile, 345; discovery of the existence of, 347; number of cases at Bilibid Prison during 1908, 71, 212.
Blastomycosis of the skin in the Philippine Islands, 395; different forms of, 396; other forms of, 402; organism and cultures of, 400; prevalence of, 395.
BOWMAN, FRED B., A series of cases of tropical infantile dysentery with a hitherto undescribed bacillus as the causative factor (preliminary report), 31.
CALDERON, FERNANDO, Obstetrics in the Philippine Islands, 245.
Carcinoma, per cent among 100 cases of, 219.
CARPENTER, DUDLEY N., Discussion of the paper by Dr. Edwards (121), 187.
CASTELLANI, ALDO, Observations on a palliative treatment of elephantiasis, 311.
Ceratophyllus fasciatus Bosc., 45.
Cercomonas, present in the Philippine Islands, 195, 201.
Cerebellar hæmorrhage, per cent among 100 cases of, 212.
Cerebral abscess, per cent among 100 cases of, 212.
Cholera, cases resembling, 186; cause of the reappearance in the Philippine Islands of, 89; cholera-like cases preceding true outbreaks, 92; differences during routine examinations, 110; dissemination by flies contaminating food, 184; factors suggesting hypothesis, 93; habits of Filipinos conducive to, 185, 90; hypothesis, 95; number of cases at Bilibid Prison during 1908 of, 71; other observations, 92; past epidemics of, 90; statistics of, 97–104; studies of, 107; variations in virulence; 107.
Cholera organisms inoculated into soda water, 187.
Chronic cardiac disease and arterial change, per cent among 100 cases of, 214.

Ciliates, per cent of, in water examined in the Philippine Islands, 122; prevalence and distribution of, in the Philippine Islands, 191.
Cladothrix, 458.
CLEGG, M. T. See MUSGRAVE, W. E.
CLEMENTS, PAUL, Discussion on the papers by Dr. Heiser (89) and Dr. Marshall (107), 185.
Complement fixation, technique and interpretation of results in syphilis, 358; experimental work, 373; in the bacterial diseases, 371; in the forensic diagnosis of blood stains, 365.
Conjunctivitis, number of cases at Bilibid Prison during 1908, 71.
CRAIG, CHARLES F. See ASHBURN, P. M.
Ctenocephalus canis Curtis, 44.
Dysenteric abscess of the liver, 349; table of cases at Colombo, 351; in French Indo-China, 351; at Manila, 349; at Singapore, 352.
Dysentery, acute, 71; acute bacillary, 216, amœbic, 71; 212; 123; 417; bacillary, 31.
EDWARDS, RALPH T., A biological study of the water supply of the Philippine Islands, with a description of a new pathogenic organism, 121.
Elephantiasis, observations on a palliative treatment of, 311; cases treated, 313–315; filariasis and, 293; technique of the treatment, 312; thiosinamin injections and methodical bandaging, 315.
Endocarditis, 212.
Excreta, resolution concerning the disposal in the Philippine Islands of, 209.
Fæces examination with reference to the presence of amœbæ, 417.
Fascioletta ilocana, 385.
Filaria nocturna, notes on the distribution of, 305.
Filariasis and elephantiasis, 293; mosquitoes as carriers of, 299; treatment and prophylaxis, 300.
Filtration experiments with virus of cattle plague, 165; further, 319.
Flagellates, prevalence and distribution in the Philippine Islands of, 191; per cent in water examined of, 122.
Fleas encountered in Hongkong: Ceratophyllus fasciatus Bosc., 45; Ctenocephalus canis Curtis, 44; Pulex cheopis Rotschild or pallidus Tasch., 44; Pulex irritans Linn., 45.
Flukes. See Trematoda.
FRASER, HENRY, Discussion on Dr. Jackson's paper (431), 443.
FREER, PAUL C., Editorials: The Free Dispensary of the Philippine Medical School, 352; The Philippine Islands Medical Association, 183.
GARRISON, PHILIP E., A new intestinal trematode of man. **Fascioletta ilocana** n. g., n. sp., 385; Discussions on the papers of Dr. Garrison (191) and Dr. Gilman (211), 263; Editorial: Intestinal parasites in the Philippines, 71; The prevalence and distribution of the animal parasites of man in the Philippine Islands, with a consideration of their possible influence upon the public health, 191.
GILMAN, PHILIP K., A report on the first 100 autopsies at the Philippine Medical School, 211; Editorial: Neoplasms among the Filipinos, 567; Editorial: Occurrence of tuberculosis in 100 autopsies in the Philippine Islands, 74.
GREENLEAF, HENRY S., Discussions on the papers of Dr. Garrison (191) and Dr. Gilman (211), 262.
Habu venom and its serum therapy, 151; action upon animals, 154; agglutination, 161; hæmolysis, 156; immunization and immunized serum, 162; physical and chemical constitution, 152; resistance to physical and chemical influences, 153; symptoms in man, 156; therapy, 163; Trimeresurus flavoviridis (Hallowell), 152; Trimesurus hypnale D & B, 188; Trimeresurus riukiuanus (Hilgendorf), 152.
Health, conditions of, in the Philippines, 269; rules published by the Bureau of Health on hygiene measures against cholera, 281.
HEISER, VICTOR G., Discussions on the papers of Dr. Garrison (191) and Dr. Gilman (211), 262; Discussion of Dr. Jackson's paper (431), 439; Discussion on the papers of Dr. Heiser (89) and Dr. Marshall (107), 186; Editorial: Prophylactic medicine as applied to the hygiene of Bilibid Prison, 69; Some considerations with regard to the cause of the frequent reappearance of cholera in the Philippine Islands (with statistics beginning with the outbreak in 1902 to January 1, 1908), 89.
Hepatitis, the prevention of tropical abscess of the liver by the early diagnosis and treatment of the presuppurative stage of amœbic, 285; case reports, 287–289; method of giving ipecacuanha, 289.
HERZOG, MAXIMILIAN, Peculiar cases of traumatism of internal organs, some due to tropical conditions and practices, 55.
Hook-worms, prevalence and distribution of, in the Philippine Islands, 191, 195, 198, 199.
HOYT, R. E., Results of 300 examinations of fæces with reference to the presence of amœbæ, 417.
Hymenolepis, prevalence and distribution in the Philippine Islands of, 191, 195, 205.

Immune serum, 382; animal experiments with, 382; clinical application of, 383; internal administration of, 384.
Infantile beriberi, 345.
Infantile dysentery, 31; case reports, 31–33.
Intestinal protozoa, prevalence and distribution in the Philippine Islands of, 191, 201.
Intestinal worms, prevalence and distribution in the Philippine Islands of, 191, 206; possible influence upon the public health of, 206–209; predisposing to disease, 206–207, 263, 264; sanitation against, 208–209, 264.
Ipecacuanha, method of giving, 289.
ISHIGAMI, T., Tuberculo-toxoidin and immunization serum, 379.
JACKSON, THOMAS W., Discussion of the papers of Dr. Garrison (191) and Dr. Gilman (211), 261; Discussion of Dr. Jackson's paper (431), 443; Sanitary conditions and needs in provincial towns, 431.
Jellyfish (Scyphozoa), a report of several cases with unusual symptoms caused by contact with some unknown variety of, 329.
KEE, TEE HAN, The habitual use of opium as a factor in the production of diseases, 63.
Kidney diseases, 215.
KITAJIMA, T., On "habu" venom and its serum therapy, 151.
Lamblia, present in the Philippine Islands, 195, 201.
Liver, lesions of the, 217.
Lymphatic system, disease of the, 219.
Malaria, number of cases at Bilibid Prison during 1908 of, 71.
MARSHALL, HARRY T., Discussion on the papers by Dr. Heiser (89) and Dr. Marshall (107), 187; Discussion of the paper by Dr. Kitajima (151), 188; Studies of cholera, 107.
MARSHALL, HARRY T., & TEAGUE, OSCAR, A study of the precipitin and complement fixation reactions, 357.
MAUS, L. M., Discussion of the papers by Dr. Heiser (89) and Dr. Marshall (107), 184.
McDILL, JOHN R., Editorial: Dysenteric abscess of the liver, 349.
Medical Association of the Philippine Islands, 183.
Meningococcus intracellularis, 47.
Mosquito which breeds in salt and fresh water, 335; character of the river and spring water of breeding places of the, 337; climate and temperature, 337; geography and topography of breeding area, 336; observations on larvæ of, 337.
Mus decumanus Pallas., 45.
Mus musculus Linn., 46.
Mus rattus Linn., 45.
MUSGRAVE, W. E., Discussions on the papers of Dr. Garrison (191) and Dr. Gilman (211), 261; The influence of symbiosis upon the pathogenicity of micro̊rganisms, 77.
MUSGRAVE, W. E., CLEGG, M. T., & POLK, MARY, Streptothricosis with special reference to the etiology and classification of mycetoma, with a bibliography, 447; Trichocephaliasis, with a report of four cases, including one fatal case, with a bibliography, 545.
Mycetoma, classification of, 449.
Necator, present in the Philippine Islands, 195.
Nephritis, per cent among 100 cases of, 215.
Neoplasms among the Filipinos, 567.
Nervous system, disease of the, 220.
NICHOLS, HENRY J., Discussions on the papers of Dr. Garrison (191) and Dr. Gilman (211), 262; Discussion on Dr. Jackson's paper (431), 442. See PHALEN, JAMES M.
Nocardia, 459.
Obstetrics in the Philippine Islands, 245; early superstitions regards, 246; modern advances in, 252; outside influences upon, 247, 250.
OHNO, Y. K., An investigation of the quantitative relationships between agglutinin, agglutinoid, and agglutinable substance, 47.
OLD, EDWARD H. H., A report of several cases with unusual symptoms caused by contact with some unknown variety of jellyfish (Scyphozoa), 329.
Oösphora, 458.
Opisthorchis, prevalence and distribution in the Philippine Islands of, 191, 195, 204, 205; 71.
Opium as a factor in the production of diseases, 63; reasons for the use, 63; results of the habitual use and the diseases produced, 65; treatment of chronic poisoning, 66.
Oxyuris, prevalence and distribution in the Philippine Islands of, 191, 195, 201, 202.
Paragonimus, prevalence and distribution in the Philippine Islands of, 191, 195, 203, 204; 71.
Parasites. See Animal parasites.
Peritonitis, 212.
PERRY, ALLAN, Discussion of Dr. Jackson's paper (431), 440.
PHALEN, JAMES M., & NICHOLS, HENRY J., Blastomycosis of the skin in the Philippine Islands, 395; Filariasis and elephantiasis in southern Luzon, 293; Notes on the condition of the liver in schistosomiasis, 223; Notes on the distribution of Filaria nocturna in the Philippine Islands, 305.

Philippine Medical School, free dispensary of the, 352; 100 autopsies at the, 211.
Plague, procedure in Hongkong against, 39; cleansing operations during, 40; disinfection of infected premises during, 4.; fleas encountered during, 44; measures in 1906-7 regards, 42; notification of, 39; procedure in 1904 against, 41; rats encountered during, 45; segregation of contacts during, 40; table of cases from 1895 to 1904 of, 46.
Plague animals and insects. See Fleas and Rats.
Plague serum, a reduction in the cost of anticattle, 407.
Pleurisy, acute, 212.
Pneumonia, per cent among 100 cases of, 213; number of cases at Bilibid Prison during 1908 of, 71.
POLK, MARY. See MUSGRAVE, W. E.
Precipitin and complement fixation reactions, 357; comparison of the two methods in the differentiation of the blood of the different races of men and of the different varieties of cattle found in the Philippine Islands, 373–376; complement fixation, technique and interpretation of results in syphilis, 358; in bacterial diseases, 371; in the forensic diagnosis of blood stains, 365; precipitin reaction, technique, 372.
Prophylactic medicine as applied to the hygiene of Bilibid Prison, 69.
Protozoa, without specification of identity, Per cent in water examined in the Philippine Islands of, 122; see Intestinal protozoa.
Pulex cheopis Rotschild (pallidus Tasch.), 44.
Pulex irritans Linn., 45.
Rats encountered in Hongkong: Mus decumanus Pallas., 45; Mus musculus Linn., 46; Mus rattus Linn., 45; Sorex giganteus Is. Geoffr., 46.
REVIEWS (book):
Dahlgren, Ulric. A Text-Book of the Principles of Animal Histology, 569.
Emery, W. D'Este. Clinical Bacteriology and Hæmatology for Practitioners, 445.
Parkes, Louis C., and Kenwood, H. R., Hygiene and Public Health, 265.
Tyrode, Maurice Vejux. Pharmacology: The Action and Uses of Drugs, 355.
Wiedersheim, Robert, Adapted from the German by Parker W. N., Comparative Anatomy of Vertebrates, 189.
Wilcox, Reynold Webb. The Treatment of Disease, a Manual of Practical Medicine (Notice only), 189.
ROGERS, LEONARD. The prevention of tropical abscess of the liver by the early diagnosis and treatment of the presuppurative stage of amœbic hepatitis, 285.
RUEDIGER, E. H., A reduction in the cost of anticattle-plague serum, 407; Filtration experiments with the virus of cattle plague, 165; Further filtration experiments with virus of cattle plague, 319.
RYLEY, C., Discussion of the paper by Dr. Edwards (121), 187.
SALEEBY, N. M., Discussions on the papers of Dr. Garrison (191) and Dr. Gilman (211), 261; Discussion on Dr. Jackson's paper (431), 442.
Sanitary conditions and needs in provincial towns, 431.
Schistosoma, prevalence and distribution in the Philippine Islands of, 191, 195, 204; 71.
Schistosomiasis, notes on the condition of the liver in schistosomiasis, 223; case record of, 223; comparison of infection with Schistosomum japonicum and Schistosomum hæmatobium, 225; geographical location, 227; pathogenesis of cirrhosis of the liver, 225.
Schistosomum hæmatobium, 225.
Schistosomum japonicum, 225.
Septicæmia, per cent among 100 cases of, 216.
Sorex giganteus Is. Geoffr., 46.
Sphærotilus, 459.
Spirochæta, African, 237–242; American, 237–242.
Spirillum cholerae, 47.
Spleen, lesions of the, 218.
Splenomegaly, per cent among 100 cases of, 212.
Spotted or Tick Fever. See Tsutsugamushi Disease.
Staphylococcus pyogenes aureus, 47.
Streptothricosis, with special reference to the etiology and classification of mycetoma, 447; animal experiments in, 455; bibliography of, 485; clinical types of, 482; complications of, 485; definition of, 477; diagnosis of, 479; etiology of, 478; history of, 477; pathology of, 481; prognosis in, 484; prophylaxis, 484; review of literature as to genus, 457; as to species, 460; symptomatology, 482; treatment, 485; synonyms of, 477, 479.
Streptothrix actinomyces, 452.
Streptothrix of bovine Actinomyces, 450.
Streptothrix canis Levy, 450, 455, 457.
Streptothrix capræ Silberschmidt, 450, 454, 457.
Streptothrix chalcea Foulerton, 450, 455, 457.
Streptothrix of Deci, 450.
Streptothrix eppingeri "A" and "B", 450, 453, 456.
Streptothrix of Eppinger ("A") and (Kl), 450.
Streptothrix of "farcin du boeuf", 450, 452.
Streptothrix freeri Musgrave & Clegg, 449, 451.
Streptothrix maduræ Vincent, 449, 450, 451, 452.
Streptothrix nocardii, 450, 452, 455.
Streptothrix "32".
STRONG, RICHARD P., The diagnosis of African tick fever from the examination of the blood, 231.
Strongyloides, prevalence and distribution of in the Philippine Islands, 191, 195, 201.

Symbiosis and its influence upon the pathogenicity of microörganisms, 77; animal and bacterial, 84; bacterial, 79.
Tænia, prevalence and distribution in the Philippine Islands of, 191, 195, 202; 7[illegible].
TEAGUE, OSCAR. See MARSHALL, HARRY T.
Thiosinamin injections in elephantiasis, 312.
Traumatism of internal organs, 55; perforation of the heart, 57; rupture of the liver due to obstetrical practice in the Philippine Islands, 59; two cases of spleen rupture in splenomegaly, 55.
Trematoda, prevalence and distribution in the Philippine Islands of, 191.
Trichocephaliasis, bibliography of, 562; case reports of, 546; definition of, 555; diagnosis of, 561; etiology of, 555; literature of, 550; pathogenesis of, 558; pathology of, 559; prognosis of, 562; prophylaxis of, 562; symptoms of, 560; synonyms of, 555; treatment of, 562.
Trichuris, female organs, 556; habitat of, 557; male organs, 556; prevalence and distribution in the Philippine Islands of, 191, 195, 197.
Trichuris trichiura, development, 557; female, 557; male, 557; habitat, 557; resistance to the action of physical and chemical agents, 557.
Trimeresurus flavoviridis (Hallowell), 152.
Trimeresurus riukiuanus (Hilgendorf), 152.
Trimesurus hypnale D. & B., 188.
Tsutsugamushi disease and spotted or tick fever, 1; blood examination of, 9; diagnosis of, 12; etiology of, 4; history of, 3; immunity conferred by attack of, 11; mortality of, 11; pathological anatomy of, 12; prognosis of, 12; prophylaxis of, 15; susceptibility of animals to, 11; symptoms of, 7; synonymy of, 1; treatment of, 15.
Tuberculosis, 212; occurrence in 100 autopsies of, 74.
Tuberculo-toxoidin, 379; clinical application of, 381; internal administration of, 384; methods of preparing, 380; prophylactic test, 381; test of toxicity, 380; therapeutic experiments, 381.
Typhus fever in Indo-China, 131; experimental infection, 133; natural infection, 132.
Vaccines, the inoculation of bacterial, as a practical method for the treatment of bacterial diseases, with special reference to the treatment of infections due to the gonococcus, 421.
VASSAL, J. J. See YERSIN, A.
WASHBURN, WILLIAM S., Health conditions in the Philippines, 269.
Water supply of the Philippine Islands, with a description of a new pathogenic organism, 121; amœbæ, 122; **Bacillus cabildonis**, 125; ciliata, 122; artesian wells, 123; flagellata, 122; protozoa, without specification of identity, 122; results obtained at Cavite from artesian well water, 188; wells and cisterns, 124.
Wells and cisterns, per cent of rejections in, 124.
WHITMORE, EUGENE R., The inoculation of bacterial vaccines as a practical method for the treatment of bacterial diseases, with special reference to the treatment of infections due to the gonococcus, 421.
WORCESTER, DEAN C., Discussions on the papers of Dr. Garrison (191) and Dr. Gilman (211), 262; Discussion of Dr. Jackson's paper (431), 440, 443.
YERSIN, A., & VASSAL, J. J., Typhus fever in Indo-China, 131.

190

VOL. III JANUARY, 1908 No. 1

THE PHILIPPINE JOURNAL OF SCIENCE

EDITED BY

PAUL C. FREER, M. D., PH. D.

CO-EDITOR

RICHARD P. STRONG, PH. B., M. D.

WITH THE COLLABORATION OF

VICTOR G. HEISER, M. D.; W. E. MUSGRAVE, M. D.
HARRY T. MARSHALL, A. B., M. D.
JOHN R. McDILL, M. D.; FERNANDO CALDERON, M. D.
JOSE ALBERT, M. D.; PHILIP K. GILMAN, A. B., M. D.
PHILIP E. GARRISON, B. A., M. D.

PUBLISHED BY

THE BUREAU OF SCIENCE

OF THE

GOVERNMENT OF THE PHILIPPINE ISLANDS

B. MEDICAL SCIENCES



MANILA
BUREAU OF PRINTING
1908

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES.

No. 1, 1902, Biological Laboratory.—Preliminary Report of the Appearance in the Philippine Islands of a Disease Clinically Resembling Glanders. By R. P. Strong, M. D.

No. 2, 1902, Chemical Laboratory.—The Preparation of Benzoyl-Acetyl Peroxide and Its Use as an Intestinal Antiseptic in Cholera and Dysentery. Preliminary Notes. By Paul C. Freer, M. D., Ph. D.

No. 3, 1903, Biological Laboratory.—A Preliminary Report on Trypanosomiasis of Horses in the Philippine Islands. By W. E. Musgrave, M. D., and Norman E. Williamson.

No. 4, 1903, Serum Laboratory.—Preliminary Report on the Study of Rinderpest of Cattle and Carabaos in the Philippine Islands. By James W. Jobling, M. D.

No. 5, 1903, Biological Laboratory.—Trypanosoma and Trypanosomiasis, with Special Reference to Surra in the Philippine Islands. By W. E. Musgrave, M. D., and Moses T. Clegg.

No. 6, 1903.—New or Noteworthy Plants, I. The American Element in the Philippine Flora. By Elmer D. Merrill, Botanist. (Issued January 20, 1904.)

No. 7, 1903, Chemical Laboratory.—The Gutta Percha and Rubber of the Philippine Islands. By Penoyer L. Sherman, jr., Ph. D.

No. 8, 1903.—A Dictionary of the Plant Names of the Philippine Islands. By Elmer D. Merrill, Botanist.

No. 9, 1903, Biological and Serum Laboratories.—A Report on Hæmorrhagic Septicæmia in Animals in the Philippine Islands. By Paul G. Woolley, M. D., and J. W. Jobling, M. D.

No. 10, 1903, Biological Laboratory.—Two Cases of a Peculiar Form of Hand Infection (Due to an Organism Resembling the Koch-Weeks Bacillus). By John R. McDill, M. D., and Wm. B. Wherry, M. D.

No. 11, 1903, Biological Laboratory.—Entomological Division, Bulletin No. 1: Preliminary Bulletin on Insects of the Cacao. (Prepared Especially for the Benefit of Farmers.) By Charles S. Banks, Entomologist.

No. 12, 1903, Biological Laboratory.—Report on Some Pulmonary Lesions Produced by the Bacillus of Hæmorrhagic Septicæmia of Carabaos. By Paul G. Woolley, M. D.

No. 13, 1904, Biological Laboratory.—A Fatal Infection by a Hitherto Undescribed Chromogenic Bacterium: Bacillus Aureus Fœtidus. By Maximilian Herzog, M. D.

No. 14, 1904.—Serum Laboratory: Texas Fever in the Philippine Islands and the Far East. By J. W. Jobling, M. D., and Paul G. Woolley, M. D. *Biological Laboratory:* Entomological Division, Bulletin No. 2: The Australian Tick (Boophilus Australis Fuller) in the Philippine Islands. By Charles S. Banks, Entomologist.

No. 15, 1904, Biological and Serum Laboratories.—Report on Bacillus Violaceus Manilæ: A Pathogenic Micro-Organism. By Paul G. Woolley, M. D.

No. 16, 1904, Biological Laboratory.—Protective Inoculation Against Asiatic Cholera: An Experimental Study. By Richard P. Strong, M. D.

No. 17, 1904.—New or Noteworthy Philippine Plants, II. By Elmer D. Merrill, Botanist.

No. 18, 1904, Biological Laboratory.—I. Amebas: Their Cultivation and Etiologic Significance. By W. E. Musgrave, M. D., and Moses T. Clegg. II. The Treatment of Intestinal Amœbiasis (Amœbic Dysentery) in the Tropics. By W. E. Musgrave, M. D.

No. 19, 1904, Biological Laboratory.—Some Observations on the Biology of the Cholera Spirillum. By W. B. Wherry, M. D.

No. 20, 1904.—Biological Laboratory: I. Does Latent or Dormant Plague Exist Where the Disease is Endemic? By Maximilian Herzog, M. D., and Charles B. Hare. *Serum Laboratory:* II. Broncho-Pneumonia of Cattle: Its Association with B. Bovisepticus. By Paul G. Woolley, M. D., and Walter Sorrell, D. V. S. III. Pinto (Paño Blanco). By Paul G. Woolley, M. D. *Chemical Laboratory:* IV. Notes on Analysis of the Water from the Manila Water Supply. By Charles L. Bliss, M. S. *Serum Laboratory:* V. Framboesia: Its Occurrence in Natives in the Philippine Islands. By Paul G. Woolley, M. D.

No. 21, 1904, Biological Laboratory.—Some Questions Relating to the Virulence of Micro-Organisms with Particular Reference to Their Immunizing Powers. By Richard P. Strong, M. D.

No. 22, 1904, Bureau of Government Laboratories.—I. A Description of the New Buildings of the Bureau of Government Laboratories. By Paul C. Freer, M. D., Ph. D. II. A Catalogue of the Library of the Bureau of Government Laboratories. By Mary Polk, Librarian.

No. 23, 1904, Biological Laboratory.—Plague: Bacteriology, Morbid Anatomy, and Histopathology (Including a Consideration of Insects as Plague Carriers). By Maximilian Herzog, M. D.

No. 24, 1904, Biological Laboratory.—Glanders: Its Diagnosis and Prevention (Together with a Report on Two Cases of Human Glanders Occurring in Manila and Some Notes on the Bacteriology and Polymorphism of Bacterium Mallei). By William B. Wherry, M. D.

No. 25, 1904.[1]—Birds from the Islands of Romblon, Sibuyan, and Cresta de Gallo. By Richard C. McGregor.

No. 26, 1904, Biological Laboratory.—The Clinical and Pathological Significance of Balantidium Coli. By Richard P. Strong, M. D.

No. 27, 1904.—A Review of the Identification of the Species Described in Blanco's Flora de Filipinas. By Elmer D. Merrill, Botanist.

No. 28, 1904.—I. The Polypodiaceæ of the Philippine Islands. II. Edible Philippine Fungi. By Edwin B. Copeland, Ph. D.

No. 29, 1904.—I. New or Noteworthy Philippine Plants, III. II. The Source of Manila Elemi. By Elmer D. Merrill, Botanist.

No. 30, 1905, Chemical Laboratory.—I. Autocatalytic Decomposition of Silver Oxide. II. Hydration in Solution. By Gilbert N. Lewis, Ph. D.

No. 31, 1905, Biological Laboratory.—I. Notes on a Case of Hæmatochyluria (Together with Some Observations on the Morphology of the Embryo Nematode, Filaria Nocturna). By William B. Wherry, M. D., and John R. McDill, M. D., Manila, P. I. II. A Search Into the Nitrate and Nitrite Content of Witte's "Peptone," with Special Reference to Its Influence on the Demonstration of the Indol and Cholera-Red Reactions. By William B. Wherry, M. D.

(Concluded on third page of cover.)

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES—Concluded.

(Concluded from second page of cover.)

No. 32, 1905.—*Biological Laboratory:* I. Intestinal Hæmorrhage as a Fatal Complication in Amœbic Dysentery and Its Association with Liver Abscess. By Richard P. Strong, M. D. II. The Action of Various Chemical Substances upon Cultures of Amœbæ. By J. B. Thomas, M. D., Baguio, Benguet. *Biological and Serum Laboratories:* III. The Pathology of Intestinal Amœbiasis. By Paul G. Woolley, M. D., and W. E. Musgrave, M. D.

No. 33, 1905, Biological Laboratory.—Further Observations on Fibrin Thrombosis in the Glomerular and in Other Renal Vessels in Bubonic Plague. By Maximilian Herzog, M. D.

No. 34, 1905.—I. Birds from Mindoro and Small Adjacent Islands. II. Notes on Three Rare Luzon Birds. By Richard C. McGregor.

No. 35, 1905.—I. New or Noteworthy Philippine Plants, IV. II. Notes on Cuming's Philippine Plants in the Herbarium of the Bureau of Government Laboratories. III. Hackel, "Notes on Philippine Grasses." IV. Ridley, "Scitimineæ Philippinenses." V. Clarke, "Philippine Acanthaceæ." By Elmer D. Merrill, Botanist.

No. 36, 1905.—A Hand-List of the Birds of the Philippine Islands. By Richard C. McGregor and Dean C. Worcester.

The previous publications of the Bureau were given out as bulletins in serial number pertaining to the entire Bureau. These publications, if they are desired, can be obtained by applying to the librarian of the Bureau of Science, Manila, P. I., or to the Director of the Bureau of Science, Manila, P. I. Correspondents will confer a favor by returning to the Bureau any previous publications which they may have in duplicate, as a number of bulletins are now out of print.

LIST OF PREVIOUS PUBLICATIONS OF THE MINING BUREAU (NOW DIVISION OF MINES OF THE BUREAU OF SCIENCE).

1890.—Descripción física, geológica y minera en bosquejo de la Isla de Panay por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1890.—Memoria descriptiva de los manantiales minero-medicinales de la Isla de Luzon, estudiados por la comisión compuesta de los Señores D. José Centano, Ingeniero de Minas y Vocal Presidente, D. Anacleto del Rosario y Sales, Vocal Farmacéutico, y D. José de Vera y Gómez, Vocal Médico.

1893.—Estudio Descriptivo de algunas manantiales minerales de Filipinas ejecutado por la comisión formada por D. Enrique Abella y Casariego, Inspector General de Minas, D. José de Vera y Gómez, Médico, y D. Anacleto del Rosario y Sales, Farmacéutico; precidido de un prólogo escrito por el Excmo. Sr. D. Angel de Avilés, Director General de Administración Civil.

1893.—Terremotos experimentados en la Isla de Luzón durante los meses de Marzo y Abril de 1892, especialmente desastrosos en Pangasinán, Unión y Benguet. Estudio ejecutado por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1901.—The Coal Measures of the Philippines. Charles H. Burritt.

1902.—Abstract of the Mining Laws (in force in the Philippines, 1902). Charles H. Burritt.

1902, Bulletin No. 1.—Platinum and Associated Rare Metals in Placer Formations. H. D. McCaskey, B. S.

1903.—Report of the Chief of the Mining Bureau of the Philippine Islands. Charles H. Burritt.

1903, Bulletin No. 2.—Complete List of Spanish Mining Claims Recorded in the Mining Bureau. Charles H. Burritt.

1903, Bulletin No. 3.—Report on a Geological Reconnoissance of the Iron Region of Angat, Bulacan. H. D. McCaskey, B. S.

1904.—Fifth Annual Report of the Mining Bureau. H. D. McCaskey.

1905.—Sixth Annual Report of the Chief of the Mining Bureau. H. D. McCaskey.

1905, Bulletin No. 4.—A Preliminary Reconnoissance of the Mancayan-Suyoc Mineral Region, Lepanto, P. I. A. J. Eveland, Geologist.

1905, Bulletin No. 5.—The Coal Deposits of Batan Island. Warren D. Smith, B. S., M. A., Geologist.

LIST OF PREVIOUS PUBLICATIONS OF THE ETHNOLOGICAL SURVEY (NOW DIVISION OF ETHNOLOGY, BUREAU OF SCIENCE).

(For sale at Bureau of Printing.)

Vol. I.—The Bontoc Igorot, by Albert Ernest Jenks. Paper, ₱4.50; half Morocco, ₱7.

Vol. II, Part 1.—Negritos of Zambales, by William Allen Reed. Paper, ₱1.25; half Morocco, ₱3.75.

Vol. II, Part 2 and Part 3.—The Nabaloi Dialect, by Otto Scheerer. The Bataks of Palawan, by Edward Y. Miller. (Bound also in one volume with Part 1, Negritos of Zambales.) Paper, ₱1.25; half Morocco, ₱3.75. Combined half Morocco, ₱5.

Vol. III.—Relaciones Agustinianas de las razas del Norte de Luzon, by Perez. Not listed by Bureau of Printing.

Vol. IV, Part 1.—Studies in Moro History, Law, and Religion, by Najeeb M. Saleeby. Paper, ₱0.75; half Morocco, ₱3.25.

---

[1] The first four bulletins in the ornithological series were published by The Ethnological Survey under the title "Bulletins of the Philippine Museum." The other ornithological publications of the Government appeared as publications of the Bureau of Government Laboratories.

CONTENTS.

Page.

ASHBURN, P. M., and CRAIG, CHARLES F. A Comparative Study of Tsutsugamushi Disease and Spotted or Tick Fever of Montana 1

BOWMAN, FRED B. A Series of Cases of Tropical Infantile Dysentery with a Hitherto Undescribed Bacillus as the Causative Factor (Preliminary Report) 31

ATKINSON, J. M. Plague Procedure in Hongkong 39

OHNO, Y. K. An Investigation of the Quantitative Relationships Between Agglutinin, Agglutinoid, and Agglutinable Substance 47

HERZOG, MAXIMILIAN. Peculiar Cases of Traumatism of Internal Organs, Some Due to Tropical Conditions and Practices 55

KEE, TEE HAN. The Habitual Use of Opium as a Factor in the Production of Diseases 63

Editorial 69

The "Philippine Journal of Science" is issued as follows:

Section A. General Science, $2, United States currency, per year.
Section B. Medical Sciences, $3, United States currency, per year.
Section C. Botany, $2, United States currency, per year.

The entire "Journal," $5, United States currency, per year.
Single numbers, 50 cents, United States currency.
Authors receive 100 copies of their paper free.
The numbers in each section will appear as rapidly as material is available. Each section will be separately paged and indexed. Subscriptions may be sent to the DIRECTOR OF PRINTING, Manila, P. I.

FOREIGN AGENTS.

THE MACMILLAN COMPANY, 64-66 Fifth Avenue, New York.
Messrs. WM. WESLEY & SON, 28 Essex Street, Strand, London, W. C.
MARTINUS NIJHOFF, Nobelstraat 18, 's Gravenhage, Holland.
Messrs. MAYER & MÜLLER, Prinz Louis Ferdinandstrasse 2, Berlin, N. W.
Messrs. KELLY & WALSH, LIMITED, 32 Raffles Place, Singapore, S. S.
Messrs. A. M. & J. FERGUSON, 19 Baillie Street, Colombo, Ceylon.

(Entered at the Post-Office at Manila, P. I., as second-class matter.)

A limited number of complete copies of Volume I, Supplement to Volume I, and Volume II, all sections, are still available for sale. Price of the complete volumes unbound, each, $5, United States currency; the Botanical Supplements to Volume I unbound, $2.50, United States currency. Volume I was not divided into sections.

VOL. III APRIL, 1908 No. 2

THE PHILIPPINE JOURNAL OF SCIENCE

EDITED BY

PAUL C. FREER, M. D., PH. D.

CO-EDITOR

RICHARD P. STRONG, PH. B., M. D.

WITH THE COLLABORATION OF

VICTOR G. HEISER, M. D.; W. E. MUSGRAVE, M. D.
HARRY T. MARSHALL, A. B., M. D.
JOHN R. McDILL, M. D.; FERNANDO CALDERON, M. D.
JOSE ALBERT, M. D.; PHILIP K. GILMAN, A. B., M. D.
PHILIP E. GARRISON, B. A., M. D.

PUBLISHED BY

THE BUREAU OF SCIENCE

OF THE

GOVERNMENT OF THE PHILIPPINE ISLANDS

B. MEDICAL SCIENCES



MANILA
BUREAU OF PRINTING
1908

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES.

No. 1, 1902, Biological Laboratory.—Preliminary Report of the Appearance in the Philippine Islands of a Disease Clinically Resembling Glanders. By R. P. Strong, M. D.

No. 2, 1902, Chemical Laboratory.—The Preparation of Benzoyl-Acetyl Peroxide and Its Use as an Intestinal Antiseptic in Cholera and Dysentery. Preliminary Notes. By Paul C. Freer, M. D., Ph. D.

No. 3, 1903, Biological Laboratory.—A Preliminary Report on Trypanosomiasis of Horses in the Philippine Islands. By W. E. Musgrave, M. D., and Norman E. Williamson.

No. 4, 1903, Serum Laboratory.—Preliminary Report on the Study of Rinderpest of Cattle and Carabaos in the Philippine Islands. By James W. Jobling, M. D.

No. 5, 1903, Biological Laboratory.—Trypanosoma and Trypanosomiasis, with Special Reference to Surra in the Philippine Islands. By W. E. Musgrave, M. D., and Moses T. Clegg.

No. 6, 1903.—New or Noteworthy Plants, I. The American Element in the Philippine Flora. By Elmer D. Merrill, Botanist. (Issued January 20, 1904.)

No. 7, 1903, Chemical Laboratory.—The Gutta Percha and Rubber of the Philippine Islands. By Penoyer L. Sherman, jr., Ph. D.

No. 8, 1903.—A Dictionary of the Plant Names of the Philippine Islands. By Elmer D. Merrill, Botanist.

No. 9, 1903, Biological and Serum Laboratories.—A Report on Hæmorrhagic Septicæmia in Animals in the Philippine Islands. By Paul G. Woolley, M. D., and J. W. Jobling, M. D.

No. 10, 1903, Biological Laboratory.—Two Cases of a Peculiar Form of Hand Infection (Due to an Organism Resembling the Koch-Weeks Bacillus). By John R. McDill, M. D., and Wm. B. Wherry, M. D.

No. 11, 1903, Biological Laboratory.—Entomological Division, Bulletin No. 1: Preliminary Bulletin on Insects of the Cacao. (Prepared Especially for the Benefit of Farmers.) By Charles S. Banks, Entomologist.

No. 12, 1903, Biological Laboratory.—Report on Some Pulmonary Lesions Produced by the Bacillus of Hæmorrhagic Septicæmia of Carabaos. By Paul G. Woolley, M. D.

No. 13, 1904, Biological Laboratory.—A Fatal Infection by a Hitherto Undescribed Chromogenic Bacterium: Bacillus Aureus Fœtidus. By Maximilian Herzog, M. D.

No. 14, 1904.—Serum Laboratory: Texas Fever in the Philippine Islands and the Far East. By J. W. Jobling, M. D., and Paul G. Woolley, M. D. *Biological Laboratory:* Entomological Division, Bulletin No. 2: The Australian Tick (Boophilus Australis Fuller) in the Philippine Islands. By Charles S. Banks, Entomologist.

No. 15, 1904, Biological and Serum Laboratories.—Report on Bacillus Violaceus Manilæ: A Pathogenic Micro-Organism. By Paul G. Woolley, M. D.

No. 16, 1904, Biological Laboratory.—Protective Inoculation Against Asiatic Cholera: An Experimental Study. By Richard P. Strong, M. D.

No. 17, 1904.—New or Noteworthy Philippine Plants, II. By Elmer D. Merrill, Botanist.

No. 18, 1904, Biological Laboratory.—I. Amebas: Their Cultivation and Etiologic Significance. By W. E. Musgrave, M. D., and Moses T. Clegg. II. The Treatment of Intestinal Amœbiasis (Amœbic Dysentery) in the Tropics. By W. E. Musgrave, M. D.

No. 19, 1904, Biological Laboratory.—Some Observations on the Biology of the Cholera Spirillum. By W. B. Wherry, M. D.

No. 20, 1904.—Biological Laboratory: I. Does Latent or Dormant Plague Exist Where the Disease is Endemic? By Maximilian Herzog, M. D., and Charles B. Hare. *Serum Laboratory:* II. Broncho-Pneumonia of Cattle: Its Association with B. Bovisepticus. By Paul G. Woolley, M. D., and Walter Sorrell, D. V. S. III. Pinto (Paño Blanco). By Paul G. Woolley, M. D. *Chemical Laboratory:* IV. Notes on Analysis of the Water from the Manila Water Supply. By Charles L. Bliss, M. S. *Serum Laboratory:* V. Frambœsia: Its Occurrence in Natives in the Philippine Islands. By Paul G. Woolley, M. D.

No. 21, 1904, Biological Laboratory.—Some Questions Relating to the Virulence of Micro-Organisms with Particular Reference to Their Immunizing Powers. By Richard P. Strong, M. D.

No. 22, 1904, Bureau of Government Laboratories.—I. A Description of the New Buildings of the Bureau of Government Laboratories. By Paul C. Freer, M. D., Ph. D. II. A Catalogue of the Library of the Bureau of Government Laboratories. By Mary Polk, Librarian.

No. 23, 1904, Biological Laboratory.—Plague: Bacteriology, Morbid Anatomy, and Histopathology (Including a Consideration of Insects as Plague Carriers). By Maximilian Herzog, M. D.

No. 24, 1904, Biological Laboratory.—Glanders: Its Diagnosis and Prevention (Together with a Report on Two Cases of Human Glanders Occurring in Manila and Some Notes on the Bacteriology and Polymorphism of Bacterium Mallei). By William B. Wherry, M. D.

No. 25, 1904.[1]—Birds from the Islands of Romblon, Sibuyan, and Cresta de Gallo. By Richard C. McGregor.

No. 26, 1904, Biological Laboratory.—The Clinical and Pathological Significance of Balantidium Coli. By Richard P. Strong, M. D.

No. 27, 1904.—A Review of the Identification of the Species Described in Blanco's Flora de Filipinas. By Elmer D. Merrill, Botanist.

No. 28, 1904.—I. The Polypodiaceæ of the Philippine Islands. II. Edible Philippine Fungi. By Edwin B. Copeland, Ph. D.

No. 29, 1904.—I. New or Noteworthy Philippine Plants, III. II. The Source of Manila Elemi. By Elmer D. Merrill, Botanist.

No. 30, 1905, Chemical Laboratory.—I. Autocatalytic Decomposition of Silver Oxide. II. Hydration in Solution. By Gilbert N. Lewis, Ph. D.

No. 31, 1905, Biological Laboratory.—I. Notes on a Case of Hæmatochyluria (Together with Some Observations on the Morphology of the Embryo Nematode, Filaria Nocturna). By William B. Wherry, M. D., and John R. McDill, M. D., Manila, P. I. II. A Search Into the Nitrate and Nitrite Content of Witte's "Peptone," with Special Reference to Its Influence on the Demonstration of the Indol and Cholera-Red Reactions. By William B. Wherry, M. D.

(Concluded on third page of cover.)

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES—Concluded.

(Concluded from second page of cover.)

No. 32, 1905.—Biological Laboratory: I. Intestinal Hæmorrhage as a Fatal Complication in Amœbic Dysentery and Its Association with Liver Abscess. By Richard P. Strong, M. D. II. The Action of Various Chemical Substances upon Cultures of Amœbæ. By J. B. Thomas, M. D., Baguio, Benguet. *Biological and Serum Laboratories:* III. The Pathology of Intestinal Amœbiasis. By Paul G. Woolley, M. D., and W. E. Musgrave, M. D.

No. 33, 1905, Biological Laboratory.—Further Observations on Fibrin Thrombosis in the Glomerular and in Other Renal Vessels in Bubonic Plague. By Maximilian Herzog, M. D.

No. 34, 1905.—I. Birds from Mindoro and Small Adjacent Islands. II. Notes on Three Rare Luzon Birds. By Richard C. McGregor.

No. 35, 1905.—I. New or Noteworthy Philippine Plants, IV. II. Notes on Cuming's Philippine Plants in the Herbarium of the Bureau of Government Laboratories. III. Hackel, "Notes on Philippine Grasses." IV. Ridley, "Scitimineæ Philippinenses." V. Clarke, "Philippine Acanthaceæ." By Elmer D. Merrill, Botanist.

No. 36, 1905.—A Hand-List of the Birds of the Philippine Islands. By Richard C. McGregor and Dean C. Worcester.

The previous publications of the Bureau were given out as bulletins in serial number pertaining to the entire Bureau. These publications, if they are desired, can be obtained by applying to the librarian of the Bureau of Science, Manila, P. I., or to the Director of the Bureau of Science, Manila, P. I. Correspondents will confer a favor by returning to the Bureau any previous publications which they may have in duplicate, as a number of bulletins are now out of print.

LIST OF PREVIOUS PUBLICATIONS OF THE MINING BUREAU (NOW DIVISION OF MINES OF THE BUREAU OF SCIENCE).

1890.—Descripción física, geológica y minera en bosquejo de la Isla de Panay por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1890.—Memoria descriptiva de los manantiales minero-medicinales de la Isla de Luzon, estudiados por la comisión compuesta de los Señores D. José Centano, Ingeniero de Minas y Vocal Presidente, D. Anacleto del Rosario y Sales, Vocal Farmacéutico, y D. José de Vera y Gómez, Vocal Médico.

1893.—Estudio Descriptivo de algunas manantiales minerales de Filipinas ejecutado por la comisión formada por D. Enrique Abella y Casariego, Inspector General de Minas, D. José de Vera y Gómez, Médico, y D. Anacleto del Rosario y Sales, Farmacéutico; precidido de un prólogo escrito por el Excmo. Sr. D. Angel de Avilés, Director General de Administración Civil.

1893.—Terremotos experimentados en la Isla de Luzón durante los meses de Marzo y Abril de 1892, especialmente desastrosos en Pangasinán, Unión y Benguet. Estudio ejecutado por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1901.—The Coal Measures of the Philippines. Charles H. Burritt.

1902.—Abstract of the Mining Laws (in force in the Philippines, 1902). Charles H. Burritt.

1902, Bulletin No. 1.—Platinum and Associated Rare Metals in Placer Formations. H. D. McCaskey, B. S.

1903.—Report of the Chief of the Mining Bureau of the Philippine Islands. Charles H. Burritt.

1903, Bulletin No. 2.—Complete List of Spanish Mining Claims Recorded in the Mining Bureau. Charles H. Burritt.

1903, Bulletin No. 3.—Report on a Geological Reconnoissance of the Iron Region of Angat, Bulacan. H. D. McCaskey, B. S.

1904.—Fifth Annual Report of the Mining Bureau. H. D. McCaskey.

1905.—Sixth Annual Report of the Chief of the Mining Bureau. H. D. McCaskey.

1905, Bulletin No. 4.—A Preliminary Reconnoissance of the Mancayan-Suyoc Mineral Region, Lepanto, P. I. A. J. Eveland, Geologist.

1905, Bulletin No. 5.—The Coal Deposits of Batan Island. Warren D. Smith, B. S., M. A., Geologist.

LIST OF PREVIOUS PUBLICATIONS OF THE ETHNOLOGICAL SURVEY (NOW DIVISION OF ETHNOLOGY, BUREAU OF SCIENCE).

(For sale at Bureau of Printing.)

Vol. I.—The Bontoc Igorot, by Albert Ernest Jenks. Paper, ₱4.50; half Morocco, ₱7.

Vol. II, Part 1.—Negritos of Zambales, by William Allen Reed. Paper, ₱1.25; half Morocco, ₱3.75.

Vol. II, Part 2 and Part 3.—The Nabaloi Dialect, by Otto Scheerer. The Bataks of Palawan, by Edward Y. Miller. (Bound also in one volume with Part 1, Negritos of Zambales.) Paper, ₱1.25; half Morocco, ₱3.75. Combined half Morocco, ₱5.

Vol. III.—Relaciones Agustinianas de las razas del Norte de Luzon, by Perez. Not listed by Bureau of Printing.

Vol. IV, Part 1.—Studies in Moro History, Law, and Religion, by Najeeb M. Saleeby. Paper, ₱0.75; half Morocco, ₱3.25.

---

[1] The first four bulletins in the ornithological series were published by The Ethnological Survey under the title "Bulletins of the Philippine Museum." The other ornithological publications of the Government appeared as publications of the Bureau of Government Laboratories.

CONTENTS

	Page.
MUSGRAVE, W. E. The Influence of Symbiosis Upon the Pathogenicity of Microörganisms	77
HEISER, VICTOR G. Some Considerations with Regard to the Cause of the Frequent Reappearance of Cholera in the Philippine Islands (with Statistics Beginning with the Outbreak in 1902 to January 1, 1908)	89
MARSHALL, HARRY T. Studies of Cholera	107
EDWARDS, RALPH T. A Biological Study of the Water Supply of the Philippine Islands, with a Description of a New Pathogenic Organism	121
YERSIN, A., and VASSAL, J. J. Typhus Fever in Indo-China	131
KITAJIMA, T. On "Habu" Venom and its Serum Therapy	151
RUEDIGER, E. H. Filtration Experiments with the Virus of Cattle Plague	165
EDITORIAL	183
REVIEWS	189

The "Philippine Journal of Science" is issued as follows:

Section A. General Science, $2, United States currency, per year.
Section B. Medical Sciences, $3, United States currency, per year.
Section C. Botany, $2, United States currency, per year.

The entire "Journal," $5, United States currency, per year.
Single numbers, 50 cents, United States currency.
Authors receive 100 copies of their paper free.
The numbers in each section will appear as rapidly as material is available. Each section will be separately paged and indexed. Subscriptions may be sent to the DIRECTOR OF PRINTING, Manila, P. I.

FOREIGN AGENTS.

THE MACMILLAN COMPANY, 64-66 Fifth Avenue, New York.
Messrs. WM. WESLEY & SON, 28 Essex Street, Strand, London, W. C.
MARTINUS NIJHOFF, Nobelstraat 18, 's Gravenhage, Holland.
Messrs. MAYER & MÜLLER, Prinz Louis Ferdinandstrasse 2, Berlin, N. W.
Messrs. KELLY & WALSH, LIMITED, 32 Raffles Place, Singapore, S. S.
Messrs. A. M. & J. FERGUSON, 19 Baillie Street, Colombo, Ceylon.

(Entered at the Post-Office at Manila, P. I., as second-class matter.)

A limited number of complete copies of Volume I, Supplement to Volume I, and Volume II, all sections, are still available for sale. Price of the complete volumes unbound, each, $5, United States currency; the Botanical Supplements to Volume I unbound, $2.50, United States currency. Volume I was not divided into sections.

VOL. III JULY, 1908 No. 3

THE PHILIPPINE JOURNAL OF SCIENCE

EDITED BY

PAUL C. FREER, M. D., PH. D.

CO-EDITOR

RICHARD P. STRONG, PH. B., M. D.

WITH THE COLLABORATION OF

VICTOR G. HEISER, M. D.; W. E. MUSGRAVE, M. D.
HARRY T. MARSHALL, A. B., M. D.
JOHN R. McDILL, M. D.; FERNANDO CALDERON, M. D.
JOSE ALBERT, M. D.; PHILIP K. GILMAN, A. B., M. D.
PHILIP E. GARRISON, B. A., M. D.

PUBLISHED BY

THE BUREAU OF SCIENCE

OF THE

GOVERNMENT OF THE PHILIPPINE ISLANDS

B. MEDICAL SCIENCES



MANILA
BUREAU OF PRINTING
1908

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES.

[1] *No. 1, 1902, Biological Laboratory.*—Preliminary Report of the Appearance in the Philippine Islands of a Disease Clinically Resembling Glanders. By R. P. Strong, M. D.

No. 2, 1902, Chemical Laboratory.—The Preparation of Benzoyl-Acetyl Peroxide and Its Use as an Intestinal Antiseptic in Cholera and Dysentery. Preliminary Notes. By Paul C. Freer, M. D., Ph. D.

[1] *No. 3, 1903, Biological Laboratory.*—A Preliminary Report on Trypanosomiasis of Horses in the Philippine Islands. By W. E. Musgrave, M. D., and Norman E. Williamson.

[1] *No. 4, 1903, Serum Laboratory.*—Preliminary Report on the Study of Rinderpest of Cattle and Carabaos in the Philippine Islands. By James W. Jobling, M. D.

[1] *No. 5, 1903, Biological Laboratory.*—Trypanosoma and Trypanosomiasis, with Special Reference to Surra in the Philippine Islands. By W. E. Musgave, M. D., and Moses T. Clegg.

[1] *No. 6, 1903.*—New and Noteworthy Plants, I. The American Element in the Philippine Flora. By Elmer D. Merrill, Botanist. (Issued January 20, 1904.)

[1] *No. 7, 1903, Chemical Laboratory.*—The Gutta Percha and Rubber of the Philippine Islands. By Penoyer L. Sherman, jr., Ph. D.

[1] *No. 8, 1903.*—A Dictionary of the Plant Names of the Philippine Islands. By Elmer D. Merrill, Botanist.

[1] *No. 9, 1903, Biological and Serum Laboratories.*—A Report on Hæmorrhagic Septicæmia in Animals in the Philippine Islands. By Paul G. Woolley, M. D., and J. W. Jobling, M. D.

[1] *No. 10, 1903, Biological Laboratory.*—Two Cases of a Peculiar Form of Hand Infection (Due to an Organism Resembling the Koch-Weeks Bacillus). By John R. McDill, M. D., and Wm. B. Wherry, M. D.

[1] *No. 11, 1903, Biological Laboratory.*—Entomological Division, Bulletin No. 1: Preliminary Bulletin on Insects of the Cacao. (Prepared Especially for the Benefit of Farmers.) By Charles S. Banks, Entomologist.

[1] *No. 12, 1903, Biological Laboratory.*—Report on Some Pulmonary Lesions Produced by the Bacillus of Hæmorrhagic Septicæmia of Carabaos. By Paul G. Woolley, M. D.

No. 13, 1904, Biological Laboratory.—A Fatal Infection by a Hitherto Undescribed Chromogenic Bacterium: Bacillus Aureus Fœtidus. By Maximilian Herzog, M. D.

[1] *No. 14, 1904.*—*Serum Laboratory:* Texas Fever in the Philippine Islands and the Far East. By J. W. Jobling, M. D., and Paul G. Woolley, M. D. *Biological Laboratory:* Entomological Division, Bulletin No. 2: The Australian Tick (Boophilus Australis Fuller) in the Philippine Islands. By Charles S. Banks, Entomologist.

No. 15, 1904, Biological and Serum Laboratories.—Report on Bacillus Violaceus Manilæ: A Pathogenic Micro-Organism. By Paul G. Woolley, M. D.

[1] *No. 16, 1904, Biological Laboratory.*—Protective Inoculation Against Asiatic Cholera: An Experimental Study. By Richard P. Strong, M. D.

No. 17, 1904.—New or Noteworthy Philippine Plants, II. By Elmer D. Merrill, Botanist.

[1] *No. 18, 1904, Biological Laboratory.*—I. Amebas: Their Cultivation and Etiologic Significance. By W. E. Musgrave, M. D., and Moses T. Clegg. II. The Treatment of Intestinal Amœbiasis (Amœbic Dysentery) in the Tropics. By W. E. Musgrave, M. D.

No. 19, 1904, Biological Laboratory.—Some Observations on the Biology of the Cholera Spirillum. By W. B. Wherry, M. D.

No. 20, 1904.—*Biological Laboratory:* I. Does Latent or Dormant Plague Exist Where the Disease is Endemic? By Maximilian Herzog, M. D., and Charles B. Hare. *Serum Laboratory:* II. Broncho-Pneumonia of Cattle: Its Association with B. Boviseptícus. By Paul G. Woolley, M. D., and Walter Sorrell, D. V. S. III. Pinto (Paño Blanco). By Paul G. Woolley, M. D. *Chemical Laboratory:* IV. Notes on Analysis of the Water from the Manila Water Supply. By Charles L. Bliss, M. S. *Serum Laboratory:* V. Frambœsia: Its Occurrence in Natives in the Philippine Islands. By Paul G. Woolley, M. D.

No. 21, 1904, Biological Laboratory.—Some Questions Relating to the Virulence of Micro-Organisms with Particular Reference to Their Immunizing Powers. By Richard P. Strong, M. D.

No. 22, 1904, Bureau of Government Laboratories.—I. A Description of the New Buildings of the Bureau of Government Laboratories. By Paul C. Freer, M. D., Ph. D. II. A Catalogue of the Library of the Bureau of Government Laboratories. By Mary Polk, Librarian.

[1] *No. 23, 1904, Biological Laboratory.*—Plague: Bacteriology, Morbid Anatomy, and Histopathology (Including a Consideration of Insects as Plague Carriers). By Maximilian Herzog, M. D.

No. 24, 1904, Biological Laboratory.—Glanders: Its Diagnosis and Prevention (Together with a Report on Two Cases of Human Glanders Occurring in Manila and Some Notes on the Bacteriology and Polymorphism of Bacterium Mallei). By William B. Wherry, M. D.

No. 25, 1904.[2]—Birds from the Islands of Romblon, Sibuyan, and Cresta de Gallo. By Richard C. McGregor.

No. 26, 1904, Biological Laboratory.—The Clinical and Pathological Significance of Balantidium Coli. By Richard P. Strong, M. D.

No. 27, 1904.—A Review of the Identification of the Species Described in Blanco's Flora de Filipinas. By Elmer D. Merrill, Botanist.

No. 28, 1904.—I. The Polypodiaceæ of the Philippine Islands. II. Edible Philippine Fungi. By Edwin B. Copeland, Ph. D.

No. 29, 1904.—I. New or Noteworthy Philippine Plants, III. II. The Source of Manila Elemi. By Elmer D. Merrill, Botanist.

No. 30, 1905, Chemical Laboratory.—I. Autocalytic Decomposition of Silver Oxide. II. Hydration in Solution. By Gilbert N. Lewis, Ph. D.

No. 31, 1905, Biological Laboratory.—I. Notes on a Case of Hæmatochyluria (Together with Some Observations on the Morphology of the Embryo Nematode, Filaria Nocturna). By William B. Wherry, M. D., and John R. McDill, M. D., Manila, P. I. II. A Search Into the Nitrate and Nitrite Content of Witte's "Peptone," with Special Reference to Its Influence on the Demonstration of the Indol and Cholera-Red Reactions. By William B. Wherry, M. D.

---

[1] Out of print.

[2] The first four bulletins in the ornithological series were published by the Ethnological Survey under the title "Bulletins of the Philippine Museum." Later ornithological publications of the Government appeared as publications of the Bureau of Government Laboratories.

(Concluded on third page of cover.)

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES—Continued.

(Concluded from second page of cover.)

No. 32, 1905.—Biological Laboratory: I. Intestinal Hæmorrhage as a Fatal Complication in Amœbic Dysentery and Its Association with Liver Abscess. By Richard P. Strong, M. D. II. The Action of Various Chemical Substances upon Cultures of Amœbæ. By J. B. Thomas, M. D., Baguio, Benguet. *Biological and Serum Laboratories:* III. The Pathology of Intestinal Amœbiasis. By Paul G. Woolley, M. D., and W. E. Musgrave, M. D.

No. 33, 1905, Biological Laboratory.—Further Observations on Fibrin Thrombosis in the Glomerular and in Other Renal Vessels in Bubonic Plague. By Maximilian Herzog, M. D.

[1] *No. 34, 1905.*—I. Birds from Mindoro and Small Adjacent Islands. II. Notes on Three Rare Luzon Birds. By Richard C. McGregor.

No. 35, 1905.—I. New or Noteworthy Philippine Plants, IV. II. Notes on Cuming's Philippine Plants in the Herbarium of the Bureau of Government Laboratories. III. Hackel, "Notes on Philippine Grasses." IV. Ridley, "Scitamineæ Philippinenses." V. Clarke, "Philippine Acanthaceæ." By Elmer D. Merrill, Botanist.

No. 36, 1905.—A Hand-List of the Birds of the Philippine Islands. By Richard C. McGregor and Dean C. Worcester.

LIST OF PREVIOUS PUBLICATIONS OF THE MINING BUREAU (NOW DIVISION OF MINES OF THE BUREAU OF SCIENCE).

1890.—Descripción física, geológica y minera en bosquejo de la Isla de Panay por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1890.—Memoria descriptiva de los manantiales minero-medicinales de la Isla de Luzon, estudiados por la comisión compuesta de los Señores D. José Centeno, Ingeniero de Minas y Vocal Presidente, D. Anacleto del Rosario y Sales, Vocal Farmacéutico, y D. José de Vera y Gómez, Vocal Médico.

1893.—Estudio descriptivo de algunos manantiales minerales de Filipinas ejecutado por la comisión formada por D. Enrique Abella y Casariego, Inspector General de Minas D. José de Vera y Gómez, Médico, y D. Anacleto del Rosario y Sales, Farmacéutico; precedido de un prólogo escrito por el Excmo. Sr. D. Angel de Avilés, Director General de Administración Civil.

1893.—Terremotos experimentados en la Isla de Luzón durante los meses de Marzo y Abril de 1892, especialmente desastrosos en Pangasinán, Unión y Benguet. Estudio ejecutado por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1901.—The Coal Measures of the Philippines. Charles H. Burritt.

1902.—Abstract of the Mining Laws (in force in the Philippines, 1902). Charles H. Burritt.

1902, Bulletin No. 1.—Platinum and Associated Rare Metals in Placer Formations. H. D. McCaskey, B. S.

1903.—Report of the Chief of the Mining Bureau of the Philippine Islands. Charles H. Burritt.

1903, Bulletin No. 2.—Complete List of Spanish Mining Claims Recorded in the Mining Bureau. Charles H. Burritt.

1903, Bulletin No. 3.—Report on a Geological Reconnoissance of the Iron Region of Angat, Bulacan. H. D. McCaskey, B. S.

1904.—Fifth Annual Report of the Mining Bureau. H. D. McCaskey.

1905.—Sixth Annual Report of the Chief of the Mining Bureau. H. D. McCaskey.

1905, Bulletin No. 4.—A Preliminary Reconnoissance of the Mancayan-Suyoc Mineral Region, Lepanto, P. I. A. J. Eveland, Geologist.

1905, Bulletin No. 5.—The Coal Deposits of Batan Island. Warren D. Smith, B. S., M. A., Geologist.

DIVISION OF MINES.

1908.—The Mineral Resources of the Philippine Islands, with a Statement of the Production of Commercial Mineral Products during the year 1907, issued by Warren D. Smith, Chief of the Division of Mines.

LIST OF PREVIOUS PUBLICATIONS OF THE ETHNOLOGICAL SURVEY (NOW DIVISION OF ETHNOLOGY, BUREAU OF SCIENCE).

Vol. I.—The Bontoc Igorot, by Albert Ernest Jenks. Paper, ₱2; half morocco, ₱3.

Vol. II, Part 1.—Negritos of Zambales, by William Allen Reed. Paper, ₱0.50; half morocco, ₱1.50.

Vol. II, Part 2 and Part 3.—The Nabaloi Dialect, by Otto Scheerer. The Bataks of Palawan, by Edward Y. Miller. (Bound also in one volume with Part 1, Negritos of Zambales.) Paper, ₱0.50; half morocco, ₱1.50. Combined, half morocco, ₱3.

[1] *Vol. III.*—Relaciones Agustinianas de las razas del Norte de Luzón, by Pérez. Paper, ₱0.75; half morocco, ₱2.

Vol. IV, Part 1.—Studies in Moro History, Law, and Religion, by Najeeb M. Saleeby. Paper, ₱0.50; half morocco, ₱1.50.

DIVISION OF ETHNOLOGY.

Vol. IV, Part 2.—The History of Sulu, by Najeeb M. Saleeby. Paper, ₱1.50.

Vol. V, Part 1 and Part 2.—The Batan Dialect as a Member of the Philippine Group of Malayan Languages, by Otto Scheerer. The F and V in the Philippine Languages, by C. E. Conant. Paper, ₱1.80.

Directions for Ethnographic Observations and Collections, 1908. *For free distribution.*

All the above publications which are for sale may be obtained from the Director of Printing, Manila, P. I. All publications for free distribution may be obtained from the Librarian, Bureau of Science, Manila, P. I. Correspondents will confer a favor by returning to the Library of the Bureau of Science any bulletins which they may have in duplicate, as there is a considerable demand for bulletins out of print.

---

[1] Out of print.

CONTENTS.

Page.

GARRISON, PHILIP E. The Prevalence and Distribution of the Animal Parasites of Man in the Philippine Islands, with a Consideration of their Possible Influence upon the Public Health 191

GILMAN, PHILIP K. A Report on the First One Hundred Autopsies at the Philippine Medical School 211

PHALEN, J. N., and NICHOLS, HENRY J. Notes on the Condition of the Liver in Schistosomiasis 223

STRONG, RICHARD P. The Diagnosis of African Tick Fever from the Examination of the Blood 231

CALDERON, FERNANDO. Obstetrics in the Philippine Islands 245

EDITORIAL 261

REVIEWS 265

The "Philippine Journal of Science" is issued as follows:

Section A. General Science, $2, United States currency, per year.
Section B. Medical Sciences, $3, United States currency, per year.
Section C. Botany, $2, United States currency, per year.

The entire "Journal" $5, United States currency, per year.
Single numbers, 50 cents, United States currency.
Authors receive 100 copies of their papers free.
The numbers in each section will appear as rapidly as material is available. Each section will be separately paged and indexed. Subscriptions may be sent to the DIRECTOR OF PRINTING, Manila, P. I.

FOREIGN AGENTS.

THE MACMILLAN COMPANY, 64-66 Fifth Avenue, New York.
Messrs. WM. WESLEY & SON, 28 Essex Street, Strand, London, W. C.
MARTINUS NIJHOFF, Nobelstraat 18, 's Gravenhage, Holland.
Messrs. MAYER & MÜLLER, Prinz Louis Ferdinandstrasse 2, Berlin, N. W.
Messrs. KELLY & WALSH, LIMITED, 32 Raffles Place, Singapore, S. S.
Messrs. A. M. & J. FERGUSON, 19 Baillie Street, Colombo, Ceylon.

(Entered at the post-office at Manila, P. I., as second-class matter.)

A limited number of complete copies of Volume I and Supplement to Volume I and Volume II, all sections, are still available for sale. Price of the complete volumes, unbound, $5, United States currency; of the Botanical Supplements to Volume I, unbound, $2.50, United States currency. Volume I was not divided into sections.

VOL. III SEPTEMBER, 1908 No. 4

THE PHILIPPINE JOURNAL OF SCIENCE

EDITED BY

PAUL C. FREER, M. D., PH. D.

CO-EDITOR

RICHARD P. STRONG, PH. B., M. D.

WITH THE COLLABORATION OF

VICTOR G. HEISER, M. D.; W. E. MUSGRAVE, M. D.
JOHN R. McDILL, M. D.; FERNANDO CALDERON, M. D.
JOSE ALBERT, M. D.; PHILIP K. GILMAN, A. B., M. D.
PHILIP E. GARRISON, B. A., M. D.

PUBLISHED BY

THE BUREAU OF SCIENCE

OF THE

GOVERNMENT OF THE PHILIPPINE ISLANDS

B. MEDICAL SCIENCES

MANILA
BUREAU OF PRINTING
1908

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES.

[1] *No. 1, 1902, Biological Laboratory.*—Preliminary Report of the Appearance in the Philippine Islands of a Disease Clinically Resembling Glanders. By R. P. Strong, M. D.

No. 2, 1902, Chemical Laboratory.—The Preparation of Benzoyl-Acetyl Peroxide and Its Use as an Intestinal Antiseptic in Cholera and Dysentery. Preliminary Notes. By Paul C. Freer, M. D., Ph. D.

[1] *No. 3, 1903, Biological Laboratory.*—A Preliminary Report on Trypanosomiasis of Horses in the Philippine Islands. By W. E. Musgrave, M. D., and Norman E. Williamson.

[1] *No. 4, 1903, Serum Laboratory.*—Preliminary Report on the Study of Rinderpest of Cattle and Carabaos in the Philippine Islands. By James W. Jobling, M. D.

[1] *No. 5, 1903, Biological Laboratory.*—Trypanosoma and Trypanosomiasis, with Special Reference to Surra in the Philippine Islands. By W. E. Musgave, M. D., and Moses T. Clegg.

[1] *No. 6, 1903.*—New and Noteworthy Plants, I. The American Element in the Philippine Flora. By Elmer D. Merrill, Botanist. (Issued January 20, 1904.)

[1] *No. 7, 1903, Chemical Laboratory.*—The Gutta Percha and Rubber of the Philippine Islands. By Penoyer L. Sherman, jr., Ph. D.

[1] *No. 8, 1903.*—A Dictionary of the Plant Names of the Philippine Islands. By Elmer D. Merrill, Botanist.

[1] *No. 9, 1903, Biological and Serum Laboratories.*—A Report on Hæmorrhagic Septicæmia in Animals in the Philippine Islands. By Paul G. Woolley, M. D., and J. W. Jobling, M. D.

[1] *No. 10, 1903, Biological Laboratory.*—Two Cases of a Peculiar Form of Hand Infection (Due to an Organism Resembling the Koch-Weeks Bacillus). By John R. McDill, M. D., and Wm. B. Wherry, M. D.

[1] *No. 11, 1903, Biological Laboratory.*—Entomological Division, Bulletin No. 1: Preliminary Bulletin on Insects of the Cacao. (Prepared Especially for the Benefit of Farmers.) By Charles S. Banks, Entomologist.

[1] *No. 12, 1903, Biological Laboratory.*—Report on Some Pulmonary Lesions Produced by the Bacillus of Hæmorrhagic Septicæmia of Carabaos. By Paul G. Woolley, M. D.

No. 13, 1904, Biological Laboratory.—A Fatal Infection by a Hitherto Undescribed Chromogenic Bacterium: Bacillus Aureus Fœtidus. By Maximilian Herzog, M. D.

[1] *No. 14, 1904.—Serum Laboratory:* Texas Fever in the Philippine Islands and the Far East. By J. W. Jobling, M. D., and Paul G. Woolley, M. D. *Biological Laboratory:* Entomological Division, Bulletin No. 2: The Australian Tick (Boophilus Australis Fuller) in the Philippine Islands. By Charles S. Banks, Entomologist.

No. 15, 1904, Biological and Serum Laboratories.—Report on Bacillus Violaceus Manilæ: A Pathogenic Micro-Organism. By Paul G. Woolley, M. D.

[1] *No. 16, 1904, Biological Laboratory.*—Protective Inoculation Against Asiatic Cholera: An Experimental Study. By Richard P. Strong, M. D.

No. 17, 1904.—New or Noteworthy Philippine Plants, II. By Elmer D. Merrill, Botanist.

[1] *No. 18, 1904, Biological Laboratory.*—I. Amœbas: Their Cultivation and Etiologic Significance. By W. E. Musgrave, M. D., and Moses T. Clegg. II. The Treatment of Intestinal Amœbiasis (Amœbic Dysentery) in the Tropics. By W. E. Musgrave, M. D.

No. 19, 1904, Biological Laboratory.—Some Observations on the Biology of the Cholera Spirillum. By W. B. Wherry, M. D.

No. 20, 1904.—Biological Laboratory: I. Does Latent or Dormant Plague Exist Where the Disease is Endemic? By Maximilian Herzog, M. D., and Charles B. Hare. *Serum Laboratory:* II. Broncho-Pneumonia of Cattle: Its Association with B. Bovisepticus. By Paul G. Woolley, M. D., and Walter Sorrell, D. V. S. III. Pinto (Paño Blanco). By Paul G. Woolley, M. D. *Chemical Laboratory:* IV. Notes on Analysis of the Water from the Manila Water Supply. By Charles L. Bliss, M. S. *Serum Laboratory:* V. Frambœsia: Its Occurrence in Natives in the Philippine Islands. By Paul G. Woolley, M. D.

No. 21, 1904, Biological Laboratory.—Some Questions Relating to the Virulence of Micro-Organisms with Particular Reference to Their Immunizing Powers. By Richard P. Strong, M. D.

No. 22, 1904, Bureau of Government Laboratories.—I. A Description of the New Buildings of the Bureau of Government Laboratories. By Paul C. Freer, M. D., Ph. D. II. A Catalogue of the Library of the Bureau of Government Laboratories. By Mary Polk, Librarian.

[1] *No. 23, 1904, Biological Laboratory.*—Plague: Bacteriology, Morbid Anatomy, and Histopathology (Including a Consideration of Insects as Plague Carriers). By Maximilian Herzog, M. D.

No. 24, 1904, Biological Laboratory.—Glanders: Its Diagnosis and Prevention (Together with a Report on Two Cases of Human Glanders Occurring in Manila and Some Notes on the Bacteriology and Polymorphism of Bacterium Mallei). By William B. Wherry, M. D.

No. 25, 1904.[2]—Birds from the Islands of Romblon, Sibuyan, and Cresta de Gallo. By Richard C. McGregor.

No. 26, 1904, Biological Laboratory.—The Clinical and Pathological Significance of Balantidium Coli. By Richard P. Strong, M. D.

No. 27, 1904.—A Review of the Identification of the Species Described in Blanco's Flora de Filipinas. By Elmer D. Merrill, Botanist.

No. 28, 1904.—I. The Polypodiaceæ of the Philippine Islands. II. Edible Philippine Fungi. By Edwin B. Copeland, Ph. D.

No. 29, 1904.—I. New or Noteworthy Philippine Plants, III. II. The Source of Manila Elemi. By Elmer D. Merrill, Botanist.

No. 30, 1905, Chemical Laboratory.—I. Autocalytic Decomposition of Silver Oxide. II. Hydration in Solution. By Gilbert N. Lewis, Ph. D.

No. 31, 1905, Biological Laboratory.—I. Notes on a Case of Hæmatochyluria (Together with Some Observations on the Morphology of the Embryo Nematode, Filaria Nocturna). By William B. Wherry, M. D., and John R. McDill, M. D., Manila, P. I. II. A Search Into the Nitrate and Nitrite Content of Witte's "Peptone," with Special Reference to Its Influence on the Demonstration of the Indol and Cholera-Red Reactions. By William B. Wherry, M. D.

---

[1] Out of print.

[2] The first four bulletins in the ornithological series were published by the Ethnological Survey under the title "Bulletins of the Philippine Museum." Later ornithological publications of the Government appeared as publications of the Bureau of Government Laboratories.

(Concluded on third page of cover.)

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES—Continued.

(Concluded from second page of cover.)

No. 32, 1905.—Biological Laboratory: I. Intestinal Hæmorrhage as a Fatal Complication in Amœbic Dysentery and Its Association with Liver Abscess. By Richard P. Strong, M. D. II. The Action of Various Chemical Substances upon Cultures of Amœbæ. By J. B. Thomas, M. D., Baguio, Benguet. *Biological and Serum Laboratories:* III. The Pathology of Intestinal Amœbiasis. By Paul G. Woolley, M. D., and W. E. Musgrave, M. D.

No. 33, 1905, Biological Laboratory.—Further Observations on Fibrin Thrombosis in the Glomerular and in Other Renal Vessels in Bubonic Plague. By Maximilian Herzog, M. D.

[1] *No. 34, 1905.*—I. Birds from Mindoro and Small Adjacent Islands. II. Notes on Three Rare Luzon Birds. By Richard C. McGregor.

No. 35, 1905.—I. New or Noteworthy Philippine Plants, IV. II. Notes on Cuming's Philippine Plants in the Herbarium of the Bureau of Government Laboratories. III. Hackel, "Notes on Philippine Grasses." IV. Ridley, "Scitimineæ Philippinenses." V. Clarke, "Philippine Acanthaceæ." By Elmer D. Merrill, Botanist.

No. 36, 1905.—A Hand-List of the Birds of the Philippine Islands. By Richard C. McGregor and Dean C. Worcester.

LIST OF PREVIOUS PUBLICATIONS OF THE MINING BUREAU (NOW DIVISION OF MINES OF THE BUREAU OF SCIENCE).

1890.—Descripción física, geológica y minera en bosquejo de la Isla de Panay por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1890.—Memoria descriptiva de los manantiales minero-medicinales de la Isla de Luzon, estudiados por la comisión compuesta de los Señores D. José Centeno, Ingeniero de Minas y Vocal Presidente, D. Anacleto del Rosario y Sales, Vocal Farmacéutico, y D. José de Vera y Gómez, Vocal Médico.

1893.—Estudio descriptivo de algunos manantiales minerales de Filipinas ejecutado por la comisión formada por D. Enrique Abella y Casariego, Inspector General de Minas D. José de Vera y Gómez, Médico, y D. Anacleto del Rosario y Sales, Farmacéutico; precedido de un prólogo escrito por el Excmo. Sr. D. Angel de Avilés, Director General de Administración Civil.

1893.—Terremotos experimentados en la Isla de Luzón durante los meses de Marzo y Abril de 1892, especialmente desastrosos en Pangasinán, Unión y Benguet. Estudio ejecutado por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1901.—The Coal Measures of the Philippines. Charles H. Burritt.

1902.—Abstract of the Mining Laws (in force in the Philippines, 1902). Charles H. Burritt.

1902, Bulletin No. 1.—Platinum and Associated Rare Metals in Placer Formations. H. D. McCaskey, B. S.

1903.—Report of the Chief of the Mining Bureau of the Philippine Islands. Charles H. Burritt.

1903, Bulletin No. 2.—Complete List of Spanish Mining Claims Recorded in the Mining Bureau. Charles H. Burritt.

1903, Bulletin No. 3.—Report on a Geological Reconnoissance of the Iron Region of Angat, Bulacan. H. D. McCaskey, B. S.

1904.—Fifth Annual Report of the Mining Bureau. H. D. McCaskey.

1905.—Sixth Annual Report of the Chief of the Mining Bureau. H. D. McCaskey.

1905, Bulletin No. 4.—A Preliminary Reconnoissance of the Mancayan-Suyoc Mineral Region, Lepanto, P. I. A. J. Eveland, Geologist.

1905, Bulletin No. 5.—The Coal Deposits of Batan Island. Warren D. Smith, B. S., M. A., Geologist.

DIVISION OF MINES.

1908.—The Mineral Resources of the Philippine Islands, with a Statement of the Production of Commercial Mineral Products during the year 1907, issued by Warren D. Smith, Chief of the Division of Mines.

LIST OF PREVIOUS PUBLICATIONS OF THE ETHNOLOGICAL SURVEY (NOW DIVISION OF ETHNOLOGY, BUREAU OF SCIENCE).

Vol. I.—The Bontoc Igorot, by Albert Ernest Jenks. Paper, ₱2; half morocco, ₱3.

Vol. II, Part 1.—Negritos of Zambales, by William Allen Reed. Paper, ₱0.50; half morocco, ₱1.50.

Vol. II, Part 2 and Part 3.—The Nabaloi Dialect, by Otto Scheerer. The Bataks of Palawan, by Edward Y. Miller. (Bound also in one volume with Part 1, Negritos of Zambales.) Paper, ₱0.50; half morocco, ₱1.50. Combined, half morocco, ₱3.

[1] *Vol. III.*—Relaciones Agustinianas de las razas del Norte de Luzón, by Pérez. Paper, ₱0.75; half morocco, ₱2.

Vol. IV, Part 1.—Studies in Moro History, Law, and Religion, by Najeeb M. Saleeby. Paper, ₱0.50; half morocco, ₱1.50.

DIVISION OF ETHNOLOGY.

Vol. IV, Part 2.—The History of Sulu, by Najeeb M. Saleeby. Paper, ₱1.50.

Vol. V, Part 1 and Part 2.—The Batan Dialect as a Member of the Philippine Group of Malayan Languages, by Otto Scheerer. The F and V in the Philippine Languages, by C. E. Conant. Paper, ₱1.60.

Directions for Ethnographic Observations and Collections, 1908. *For free distribution.*

All the above publications which are for sale may be obtained from the Director of Printing, Manila, P. I. All publications for free distribution may be obtained from the Librarian, Bureau of Science, Manila, P. I. Correspondents will confer a favor by returning to the Library of the Bureau of Science any bulletins which they may have in duplicate, as there is a considerable demand for bulletins out of print.

[1] Out of print.

CONTENTS.

	Page.
WASHBURN, WILLIAM S. Health conditions in the Philippines	269
ROGERS, LEONARD. The Prevention of Tropical Abscess of the Liver by the Early Diagnosis and Treatment of the Presuppurative Stage of Amœbic Hepatitis	285
PHALEN, JAMES M., and NICHOLS, HENRY J. Filariasis and Elephantiasis in Southern Luzon	293
PHALEN, JAMES M., and NICHOLS, HENRY J. Notes on the Distribution of Filaria Nocturna in the Philippine Islands	305
CASTELLANI, ALDO. Observations on a Palliative Treatment of Elephantiasis	311
RUEDIGER, E. H. Further Filtration Experiments with Virus of Cattle Plague	319
OLD, EDWARD H. H. A Report of Several Cases with Unusual Symptoms Caused by Contact with some Unknown Variety of Jelly-Fish (Scyphozoa)	329
BANKS, CHARLES S. A Mosquito which Breeds in Salt and Fresh Water	335
ALBERT, JOSÉ. A Case of Infantile Beriberi with Autopsy Report	345
EDITORIAL	349
REVIEW	355

The "Philippine Journal of Science" is issued as follows:

Section A. General Science, $2, United States currency, per year.
Section B. Medical Sciences, $3, United States currency, per year.
Section C. Botany, $2, United States currency, per year.

The entire "Journal" $5, United States currency, per year.
Single numbers, 50 cents, United States currency.
Authors receive 100 copies of their papers free.
The numbers in each section will appear as rapidly as material is available. Each section will be separately paged and indexed. Subscriptions may be sent to the DIRECTOR OF PRINTING, Manila, P. I.

FOREIGN AGENTS.

THE MACMILLAN COMPANY, 64-66 Fifth Avenue, New York.
Messrs. WM. WESLEY & SON, 28 Essex Street, Strand, London, W. C.
MARTINUS NIJHOFF, Nobelstraat 18, 's Gravenhage, Holland.
Messrs. MAYER & MÜLLER, Prinz Louis Ferdinandstrasse 2, Berlin, N. W.
Messrs. KELLY & WALSH, LIMITED, 32 Raffles Place, Singapore, S. S.
Messrs. A. M. & J. FERGUSON, 19 Baillie Street, Colombo, Ceylon.

(Entered at the post-office at Manila, P. I., as second-class matter.)

A limited number of complete copies of Volume I and Supplement to Volume I and Volume II, all sections, are still available for sale. Price of the complete volumes, unbound, $5, United States currency; of the Botanical Supplements to Volume I, unbound, $2.50, United States currency. Volume I was not divided into sections.

VOL. III NOVEMBER, 1908 No. 5

THE PHILIPPINE JOURNAL OF SCIENCE

EDITED BY

PAUL C. FREER, M. D., PH. D.

CO-EDITOR

RICHARD P. STRONG, PH. B., M. D.

WITH THE COLLABORATION OF

VICTOR G. HEISER, M. D.; W. E. MUSGRAVE, M. D.
JOHN R. McDILL, M. D.; FERNANDO CALDERON, M. D.
JOSE ALBERT, M. D.; PHILIP K. GILMAN, A. B., M. D.
PHILIP E. GARRISON, B. A., M. D.

PUBLISHED BY

THE BUREAU OF SCIENCE

OF THE

GOVERNMENT OF THE PHILIPPINE ISLANDS

B. MEDICAL SCIENCES



MANILA
BUREAU OF PRINTING
1908

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES.

---

[1] *No. 1, 1902, Biological Laboratory.*—Preliminary Report of the Appearance in the Philippine Islands of a Disease Clinically Resembling Glanders. By R. P. Strong, M. D.

No. 2, 1902, Chemical Laboratory.—The Preparation of Benzoyl-Acetyl Peroxide and Its Use as an Intestinal Antiseptic in Cholera and Dysentery. Preliminary Notes. By Paul C. Freer, M. D., Ph. D.

[1] *No. 3, 1903, Biological Laboratory.*—A Preliminary Report on Trypanosomiasis of Horses in the Philippine Islands. By W. E. Musgrave, M. D., and Norman E. Williamson.

[1] *No. 4, 1903, Serum Laboratory.*—Preliminary Report on the Study of Rinderpest of Cattle and Carabaos in the Philippine Islands. By James W. Jobling, M. D.

[1] *No. 5, 1903, Biological Laboratory.*—Trypanosoma and Trypanosomiasis, with Special Reference to Surra in the Philippine Islands. By W. E. Musgave, M. D., and Moses T. Clegg.

[1] *No. 6, 1903.*—New and Noteworthy Plants, I. The American Element in the Philippine Flora. By Elmer D. Merrill, Botanist. (Issued January 20, 1904.)

[1] *No. 7, 1903, Chemical Laboratory.*—The Gutta Percha and Rubber of the Philippine Islands. By Penoyer L. Sherman, jr., Ph. D.

[1] *No. 8, 1903.*—A Dictionary of the Plant Names of the Philippine Islands. By Elmer D. Merrill, Botanist.

[1] *No. 9, 1903, Biological and Serum Laboratories.*—A Report on Hæmorrhagic Septicæmia in Animals in the Philippine Islands. By Paul G. Woolley, M. D., and J. W. Jobling, M. D.

[1] *No. 10, 1903, Biological Laboratory.*—Two Cases of a Peculiar Form of Hand Infection (Due to an Organism Resembling the Koch-Weeks Bacillus). By John R. McDill, M. D., and Wm. B. Wherry, M. D.

[1] *No. 11, 1903, Biological Laboratory.*—Entomological Division, Bulletin No. 1: Preliminary Bulletin on Insects of the Cacao. (Prepared Especially for the Benefit of Farmers.) By Charles S. Banks, Entomologist.

[1] *No. 12, 1903, Biological Laboratory.*—Report on Some Pulmonary Lesions Produced by the Bacillus of Hæmorrhagic Septicæmia of Carabaos. By Paul G. Woolley, M. D.

No. 13, 1904, Biological Laboratory.—A Fatal Infection by a Hitherto Undescribed Chromogenic Bacterium: Bacillus Aureus Fœtidus. By Maximilian Herzog, M. D.

[1] *No. 14, 1904.—Serum Laboratory:* Texas Fever in the Philippine Islands and the Far East. By J. W. Jobling, M. D., and Paul G. Woolley, M. D. *Biological Laboratory:* Entomological Division, Bulletin No. 2: The Australian Tick (Boophilus Australis Fuller) in the Philippine Islands. By Charles S. Banks, Entomologist.

No. 15, 1904, Biological and Serum Laboratories.—Report on Bacillus Violaceus Manilæ: A Pathogenic Micro-Organism. By Paul G. Woolley, M. D.

[1] *No. 16, 1904, Biological Laboratory.*—Protective Inoculation Against Asiatic Cholera: An Experimental Study. By Richard P. Strong, M. D.

No. 17, 1904.—New or Noteworthy Philippine Plants, II. By Elmer D. Merrill, Botanist.

[1] *No. 18, 1904, Biological Laboratory.*—I. Amebas: Their Cultivation and Etiologic Significance. By W. E. Musgrave, M. D., and Moses T. Clegg. II. The Treatment of Intestinal Amœbiasis (Amœbic Dysentery) in the Tropics. By W. E. Musgrave, M. D.

No. 19, 1904, Biological Laboratory.—Some Observations on the Biology of the Cholera Spirillum. By W. B. Wherry, M. D.

No. 20, 1904.—Biological Laboratory: I. Does Latent or Dormant Plague Exist Where the Disease is Endemic? By Maximilian Herzog, M. D., and Charles B. Hare. *Serum Laboratory:* II. Broncho-Pneumonia of Cattle: Its Association with B. Bovisepticus. By Paul G. Woolley, M. D., and Walter Sorrell, D. V. S. III. Pinto (Paño Blanco). By Paul G. Woolley, M. D. *Chemical Laboratory:* IV. Notes on Analysis of the Water from the Manila Water Supply. By Charles L. Bliss, M. S. *Serum Laboratory:* V. Framboesia: Its Occurrence in Natives in the Philippine Islands. By Paul G. Woolley, M. D.

No. 21, 1904, Biological Laboratory.—Some Questions Relating to the Virulence of Micro-Organisms with Particular Reference to Their Immunizing Powers. By Richard P. Strong, M. D.

No. 22, 1904, Bureau of Government Laboratories.—I. A Description of the New Buildings of the Bureau of Government Laboratories. By Paul C. Freer, M. D., Ph. D. II. A Catalogue of the Library of the Bureau of Government Laboratories. By Mary Polk, Librarian.

[1] *No. 23, 1904, Biological Laboratory.*—Plague: Bacteriology, Morbid Anatomy, and Histopathology (Including a Consideration of Insects as Plague Carriers). By Maximilian Herzog, M. D.

No. 24, 1904, Biological Laboratory.—Glanders: Its Diagnosis and Prevention (Together with a Report on Two Cases of Human Glanders Occurring in Manila and Some Notes on the Bacteriology and Polymorphism of Bacterium Mallei). By William B. Wherry, M. D.

No. 25, 1904.[2]—Birds from the Islands of Romblon, Sibuyan, and Cresta de Gallo. By Richard C. McGregor.

No. 26, 1904, Biological Laboratory.—The Clinical and Pathological Significance of Balantidium Coli. By Richard P. Strong, M. D.

No. 27, 1904.—A Review of the Identification of the Species Described in Blanco's Flora de Filipinas. By Elmer D. Merrill, Botanist.

No. 28, 1904.—I. The Polypodiaceæ of the Philippine Islands. II. Edible Philippine Fungi. By Edwin B. Copeland, Ph. D.

No. 29, 1904.—I. New or Noteworthy Philippine Plants, III. II. The Source of Manila Elemi. By Elmer D. Merrill, Botanist.

No. 30, 1905, Chemical Laboratory.—I. Autocalytic Decomposition of Silver Oxide. II. Hydration in Solution. By Gilbert N. Lewis, Ph. D.

No. 31, 1905, Biological Laboratory.—I. Notes on a Case of Hæmatochyluria (Together with Some Observations on the Morphology of the Embryo Nematode, Filaria Nocturna). By William B. Wherry, M. D., and John R. McDill, M. D., Manila, P. I. II. A Search Into the Nitrate and Nitrite Content of Witte's "Peptone," with Special Reference to Its Influence on the Demonstration of the Indol and Cholera-Red Reactions. By William B. Wherry, M. D.

---

[1] Out of print.

[2] The first four bulletins in the ornithological series were published by the Ethnological Survey under the title "Bulletins of the Philippine Museum." Later ornithological publications of the Government appeared as publications of the Bureau of Government Laboratories.

(Concluded on third page of cover.)

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES—Continued.

(Concluded from second page of cover.)

No. 32, 1905.—Biological Laboratory: I. Intestinal Hæmorrhage as a Fatal Complication in Amœbic Dysentery and Its Association with Liver Abscess. By Richard P. Strong, M. D. II. The Action of Various Chemical Substances upon Cultures of Amœbæ. By J. B. Thomas, M. D., Baguio, Benguet. *Biological and Serum Laboratories:* III. The Pathology of Intestinal Amœbiasis. By Paul G. Woolley, M. D., and W. E. Musgrave, M. D.

No. 33, 1905, Biological Laboratory.—Further Observations on Fibrin Thrombosis in the Glomerular and in Other Renal Vessels in Bubonic Plague. By Maximilian Herzog, M. D.

[1] *No. 34, 1905.*—I. Birds from Mindoro and Small Adjacent Islands. II. Notes on Three Rare Luzon Birds. By Richard C. McGregor.

No. 35, 1905.—I. New or Noteworthy Philippine Plants, IV. II. Notes on Cuming's Philippine Plants in the Herbarium of the Bureau of Government Laboratories. III. Hackel, "Notes on Philippine Grasses." IV. Ridley, "Scitaminæ Philippinenses." V. Clarke, "Philippine Acanthaceæ." By Elmer D. Merrill, Botanist.

No. 36, 1905.—A Hand-List of the Birds of the Philippine Islands. By Richard C. McGregor and Dean C. Worcester.

LIST OF PREVIOUS PUBLICATIONS OF THE MINING BUREAU (NOW DIVISION OF MINES OF THE BUREAU OF SCIENCE).

1890.—Descripción física, geológica y minera en bosquejo de la Isla de Panay por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1890.—Memoria descriptiva de los manantiales minero-medicinales de la Isla de Luzon, estudiados por la comisión compuesta de los Señores D. José Centeno, Ingeniero de Minas y Vocal Presidente, D. Anacleto del Rosario y Sales, Vocal Farmacéutico, y D. José de Vera y Gómez, Vocal Médico.

1893.—Estudio descriptivo de algunos manantiales minerales de Filipinas ejecutado por la comisión formada por D. Enrique Abella y Casariego, Inspector General de Minas D. José de Vera y Gómez, Médico, y D. Anacleto del Rosario y Sales, Farmacéutico; precedido de un prólogo escrito por el Excmo. Sr. D. Angel de Avilés, Director General de Administración Civil.

1893.—Terremotos experimentados en la Isla de Luzón durante los meses de Marzo y Abril de 1892, especialmente desastrosos en Pangasinán, Unión y Benguet. Estudio ejecutado por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1901.—The Coal Measures of the Philippines. Charles H. Burritt.

1902.—Abstract of the Mining Laws (in force in the Philippines, 1902). Charles H. Burritt.

1902, Bulletin No. 1.—Platinum and Associated Rare Metals in Placer Formations. H. D. McCaskey, B. S.

1903.—Report of the Chief of the Mining Bureau of the Philippine Islands. Charles H. Burritt.

1903, Bulletin No. 2.—Complete List of Spanish Mining Claims Recorded in the Mining Bureau. Charles H. Burritt.

1903, Bulletin No. 3.—Report on a Geological Reconnoissance of the Iron Region of Angat, Bulacan. H. D. McCaskey, B. S.

1904.—Fifth Annual Report of the Mining Bureau. H. D. McCaskey.

1905.—Sixth Annual Report of the Chief of the Mining Bureau. H. D. McCaskey.

1905, Bulletin No. 4.—A Preliminary Reconnoissance of the Mancayan-Suyoc Mineral Region, Lepanto, P. I. A. J. Eveland, Geologist.

1905, Bulletin No. 5.—The Coal Deposits of Batan Island. Warren D. Smith, B. S., M. A., Geologist.

DIVISION OF MINES.

1908.—The Mineral Resources of the Philippine Islands, with a Statement of the Production of Commercial Mineral Products during the year 1907, issued by Warren D. Smith, Chief of the Division of Mines.

LIST OF PREVIOUS PUBLICATIONS OF THE ETHNOLOGICAL SURVEY (NOW DIVISION OF ETHNOLOGY, BUREAU OF SCIENCE).

Vol. I.—The Bontoc Igorot, by Albert Ernest Jenks. Paper, ₱2; half morocco, ₱3.

Vol. II, Part 1.—Negritos of Zambales, by William Allen Reed. Paper, ₱0.50; half morocco, ₱1.50.

Vol. II, Part 2 and Part 3.—The Nabaloi Dialect, by Otto Scheerer. The Bataks of Palawan, by Edward Y. Miller. (Bound also in one volume with Part 1, Negritos of Zambales.) Paper, ₱0.50; half morocco, ₱1.50. Combined, half morocco, ₱3.

[1] *Vol. III.*—Relaciones Agustinianas de las razas del Norte de Luzón, by Pérez. Paper, ₱0.75; half morocco, ₱2.

Vol. IV, Part 1.—Studies in Moro History, Law, and Religion, by Najeeb M. Saleeby. Paper, ₱0.50; half morocco, ₱1.50.

DIVISION OF ETHNOLOGY.

Vol. IV, Part 2.—The History of Sulu, by Najeeb M. Saleeby. Paper, ₱1.50.

Vol. V, Part 1 and Part 2.—The Batan Dialect as a Member of the Philippine Group of Malayan Languages, by Otto Scheerer. The F and V in the Philippine Languages, by C. E. Conant. Paper, ₱1.60.

Directions for Ethnographic Observations and Collections, 1908. *For free distribution.*

All the above publications which are for sale may be obtained from the Director of Printing, Manila, P. I. All publications for free distribution may be obtained from the Librarian, Bureau of Science, Manila, P. I. Correspondents will confer a favor by returning to the Library of the Bureau of Science any bulletins which they may have in duplicate, as there is a considerable demand for bulletins out of print.

---

[1] Out of print.

CONTENTS.

	Page.
MARSHALL, HARRY T., AND TEAGUE, OSCAR. A Study of the Precipitin and Complement Fixation Reactions	357
ISHIGAMI, T. Tuberculo-toxoidin and Immunization Serum	379
GARRISON, PHILIP E. A New Intestinal Trematode of Man. Fascioletta ilocana n. g., n. sp	385
PHALEN, JAMES M., AND NICHOLS, HENRY J. Blastomycosis of the Skin in the Philippine Islands	395
RUEDIGER, E. H. A Reduction in the Cost of Anti Cattleplague Serum	407
HOYT, R. E. Results of Three Hundred Examinations of Fæces with Reference to the Presence of Amœbæ	417
WHITMORE, EUGENE R. The Inoculation of Bacterial Vaccines as a Practical Method for the Treatment of Bacterial Diseases, with Special Reference to the Treatment of Infections due to the Gonococcus	421
JACKSON, THOMAS W. Sanitary Conditions and Needs in Provincial Towns	431
EDITORIAL	439
REVIEW	445

The "Philippine Journal of Science" is issued as follows:

Section A. General Science, $2, United States currency, per year.
Section B. Medical Sciences, $3, United States currency, per year.
Section C. Botany, $2, United States currency, per year.

The entire "Journal" $5, United States currency, per year.
Single numbers, 50 cents, United States currency.
Authors receive 100 copies of their papers free.
The numbers in each section will appear as rapidly as material is available. Each section will be separately paged and indexed. Subscriptions may be sent to the DIRECTOR OF PRINTING, Manila, P. I.

FOREIGN AGENTS.

THE MACMILLAN COMPANY, 64-66 Fifth Avenue, New York.
Messrs. WM. WESLEY & SON, 28 Essex Street, Strand, London, W. C.
MARTINUS NIJHOFF, Nobelstraat 18, 's Gravenhage, Holland.
Messrs. MAYER & MÜLLER, Prinz Louis Ferdinandstrasse 2, Berlin, N. W.
Messrs. KELLY & WALSH, LIMITED, 32 Raffles Place, Singapore, S. S.
Messrs. A. M. & J. FERGUSON, 19 Baillie Street, Colombo, Ceylon.

(Entered at the post-office at Manila, P. I., as second-class matter.)

A limited number of complete copies of Volume I and Supplement to Volume I and Volume II, all sections, are still available for sale. Price of the complete volumes, unbound, $5, United States currency; of the Botanical Supplements to Volume I, unbound, $2.50, United States currency. Volume I was not divided into sections.

VOL. III DECEMBER, 1908 No. 6

117

THE PHILIPPINE JOURNAL OF SCIENCE

EDITED BY

PAUL C. FREER, M. D., PH. D.

CO-EDITOR

RICHARD P. STRONG, PH. B., M. D.

WITH THE COLLABORATION OF

VICTOR G. HEISER, M. D.; W. E. MUSGRAVE, M. D.
JOHN R. McDILL, M. D.; FERNANDO CALDERON, M. D.
JOSE ALBERT, M. D.; PHILIP K. GILMAN, A. B., M. D.
PHILIP E. GARRISON, B. A., M. D.

PUBLISHED BY

THE BUREAU OF SCIENCE

OF THE

GOVERNMENT OF THE PHILIPPINE ISLANDS

B. MEDICAL SCIENCES



MANILA
BUREAU OF PRINTING
1908

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES.

[1] *No. 1, 1902, Biological Laboratory.*—Preliminary Report of the Appearance in the Philippine Islands of a Disease Clinically Resembling Glanders. By R. P. Strong, M. D.

No. 2, 1902, Chemical Laboratory.—The Preparation of Benzoyl-Acetyl Peroxide and Its Use as an Intestinal Antiseptic in Cholera and Dysentery. Preliminary Notes. By Paul C. Freer, M. D., Ph. D.

[1] *No. 3, 1903, Biological Laboratory.*—A Preliminary Report on Trypanosomiasis of Horses in the Philippine Islands. By W. E. Musgrave, M. D., and Norman E. Williamson.

[1] *No. 4, 1903, Serum Laboratory.*—Preliminary Report on the Study of Rinderpest of Cattle and Carabaos in the Philippine Islands. By James W. Jobling, M. D.

[1] *No. 5, 1903, Biological Laboratory.*—Trypanosoma and Trypanosomiasis, with Special Reference to Surra in the Philippine Islands. By W. E. Musgave, M. D., and Moses T. Clegg.

[1] *No. 6, 1903.*—New and Noteworthy Plants, I. The American Element in the Philippine Flora. By Elmer D. Merrill, Botanist. (Issued January 20, 1904.)

[1] *No. 7, 1903, Chemical Laboratory.*—The Gutta Percha and Rubber of the Philippine Islands. By Penoyer L. Sherman, jr., Ph. D.

[1] *No. 8, 1903.*—A Dictionary of the Plant Names of the Philippine Islands. By Elmer D. Merrill, Botanist.

[1] *No. 9, 1903, Biological and Serum Laboratories.*—A Report on Hæmorrhagic Septicæmia in Animals in the Philippine Islands. By Paul G. Woolley, M. D., and J. W. Jobling, M. D.

[1] *No. 10, 1903, Biological Laboratory.*—Two Cases of a Peculiar Form of Hand Infection (Due to an Organism Resembling the Koch-Weeks Bacillus). By John R. McDill, M. D., and Wm. B. Wherry, M. D.

[1] *No. 11, 1903, Biological Laboratory.*—Entomological Division, Bulletin No. 1: Preliminary Bulletin on Insects of the Cacao. (Prepared Especially for the Benefit of Farmers.) By Charles S. Banks, Entomologist.

[1] *No. 12, 1903, Biological Laboratory.*—Report on Some Pulmonary Lesions Produced by the Bacillus of Hæmorrhagic Septicæmia of Carabaos. By Paul G. Woolley, M. D.

No. 13, 1904, Biological Laboratory.—A Fatal Infection by a Hitherto Undescribed Chromogenic Bacterium: Bacillus Aureus Fœtidus. By Maximilian Herzog, M. D.

[1] *No. 14, 1904.—Serum Laboratory:* Texas Fever in the Philippine Islands and the Far East. By J. W. Jobling, M. D., and Paul G. Woolley, M. D. *Biological Laboratory:* Entomological Division, Bulletin No. 2: The Australian Tick (Boophilus Australis Fuller) in the Philippine Islands. By Charles S. Banks, Entomologist.

No. 15, 1904, Biological and Serum Laboratories.—Report on Bacillus Violaceus Manilæ: A Pathogenic Micro-Organism. By Paul G. Woolley, M. D.

[1] *No. 16, 1904, Biological Laboratory.*—Protective Inoculation Against Asiatic Cholera: An Experimental Study. By Richard P. Strong, M. D.

No. 17, 1904.—New or Noteworthy Philippine Plants, II. By Elmer D. Merrill, Botanist.

[1] *No. 18, 1904, Biological Laboratory.*—I. Amebas: Their Cultivation and Etiologic Significance. By W. E. Musgrave, M. D., and Moses T. Clegg. II. The Treatment of Intestinal Amœbiasis (Amœbic Dysentery) in the Tropics. By W. E. Musgrave, M. D.

No. 19, 1904, Biological Laboratory.—Some Observations on the Biology of the Cholera Spirillum. By W. B. Wherry, M. D.

No. 20, 1904.—Biological Laboratory: I. Does Latent or Dormant Plague Exist Where the Disease is Endemic? By Maximilian Herzog, M. D., and Charles B. Hare. *Serum Laboratory:* II. Broncho-Pneumonia of Cattle: Its Association with B. Bovisepticus. By Paul G. Woolley, M. D., and Walter Sorrell, D. V. S. III. Pinto (Paño Blanco). By Paul G. Woolley, M. D. *Chemical Laboratory:* IV. Notes on Analysis of the Water from the Manila Water Supply. By Charles L. Bliss, M. S. *Serum Laboratory:* V. Frambœsia: Its Occurrence in Natives in the Philippine Islands. By Paul G. Woolley, M. D.

No. 21, 1904, Biological Laboratory.—Some Questions Relating to the Virulence of Micro-Organisms with Particular Reference to Their Immunizing Powers. By Richard P. Strong, M. D.

No. 22, 1904, Bureau of Government Laboratories.—I. A Description of the New Buildings of the Bureau of Government Laboratories. By Paul C. Freer, M. D., Ph. D. II. A Catalogue of the Library of the Bureau of Government Laboratories. By Mary Polk, Librarian.

[1] *No. 23, 1904, Biological Laboratory.*—Plague: Bacteriology, Morbid Anatomy, and Histopathology (Including a Consideration of Insects as Plague Carriers). By Maximilian Herzog, M. D.

No. 24, 1904, Biological Laboratory.—Glanders: Its Diagnosis and Prevention (Together with a Report on Two Cases of Human Glanders Occurring in Manila and Some Notes on the Bacteriology and Polymorphism of Bacterium Mallei). By William B. Wherry, M. D.

No. 25, 1904.[2]—Birds from the Islands of Romblon, Sibuyan, and Cresta de Gallo. By Richard C. McGregor.

No. 26, 1904, Biological Laboratory.—The Clinical and Pathological Significance of Balantidium Coli. By Richard P. Strong, M. D.

No. 27, 1904.—A Review of the Identification of the Species Described in Blanco's Flora de Filipinas. By Elmer D. Merrill, Botanist.

No. 28, 1904.—I. The Polypodiaceæ of the Philippine Islands. II. Edible Philippine Fungi. By Edwin B. Copeland, Ph. D.

No. 29, 1904.—I. New or Noteworthy Philippine Plants, III. II. The Source of Manila Elemi. By Elmer D. Merrill, Botanist.

No. 30, 1905, Chemical Laboratory.—I. Autocalytic Decomposition of Silver Oxide. II. Hydration in Solution. By Gilbert N. Lewis, Ph. D.

No. 31, 1905, Biological Laboratory.—I. Notes on a Case of Hæmatochyluria (Together with Some Observations on the Morphology of the Embryo Nematode, Filaria Nocturna). By William B. Wherry, M. D., and John R. McDill, M. D., Manila, P. I. II. A Search Into the Nitrate and Nitrite Content of Witte's "Peptone," with Special Reference to Its Influence on the Demonstration of the Indol and Cholera-Red Reactions. By William B. Wherry, M. D.

---

[1] Out of print.

[2] The first four bulletins in the ornithological series were published by the Ethnological Survey under the title "Bulletins of the Philippine Museum." Later ornithological publications of the Government appeared as publications of the Bureau of Government Laboratories.

(Concluded on third page of cover.)

PREVIOUS PUBLICATIONS OF THE BUREAU OF GOVERNMENT LABORATORIES—Continued.

(Concluded from second page of cover.)

No. 32, 1905.—Biological Laboratory: I. Intestinal Hæmorrhage as a Fatal Complication in Amœbic Dysentery and Its Association with Liver Abscess. By Richard P. Strong, M. D. II. The Action of Various Chemical Substances upon Cultures of Amœbæ. By J. B. Thomas, M. D., Baguio, Benguet. *Biological and Serum Laboratories:* III. The Pathology of Intestinal Amœbiasis. By Paul G. Woolley, M. D., and W. E. Musgrave, M. D.

No. 33, 1905, Biological Laboratory.—Further Observations on Fibrin Thrombosis in the Glomerular and in Other Renal Vessels in Bubonic Plague. By Maximilian Herzog, M. D.

[1] *No. 34, 1905.*—I. Birds from Mindoro and Small Adjacent Islands. II. Notes on Three Rare Luzon Birds. By Richard C. McGregor.

No. 35, 1905.—I. New or Noteworthy Philippine Plants, IV. II. Notes on Cuming's Philippine Plants in the Herbarium of the Bureau of Government Laboratories. III. Hackel, "Notes on Philippine Grasses." IV. Ridley, "Scitimineæ Philippinenses." V. Clarke, "Philippine Acanthaceæ." By Elmer D. Merrill, Botanist.

No. 36, 1905.—A Hand-List of the Birds of the Philippine Islands. By Richard C. McGregor and Dean C. Worcester.

LIST OF PREVIOUS PUBLICATIONS OF THE MINING BUREAU (NOW DIVISION OF MINES OF THE BUREAU OF SCIENCE).

1890.—Descripción física, geológica y minera en bosquejo de la Isla de Panay por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1890.—Memoria descriptiva de los manantiales minero-medicinales de la Isla de Luzon, estudiados por la comisión compuesta de los Señores D. José Centeno, Ingeniero de Minas y Vocal Presidente, D. Anacleto del Rosario y Sáles, Vocal Farmacéutico, y D. José de Vera y Gómez, Vocal Médico.

1893.—Estudio descriptivo de algunos manantiales minerales de Filipinas ejecutado por la comisión formada por D. Enrique Abella y Casariego, Inspector General de Minas D. José de Vera y Gómez, Médico, y D. Anacleto del Rosario y Sales, Farmacéutico; precedido de un prólogo escrito por el Excmo. Sr. D. Angel de Avilés, Director General de Administración Civil.

1893.—Terremotos experimentados en la Isla de Luzón durante los meses de Marzo y Abril de 1892, especialmente desastrosos en Pangasinán, Unión y Benguet. Estudio ejecutado por D. Enrique Abella y Casariego, Inspector General de Minas del Archipiélago.

1901.—The Coal Measures of the Philippines. Charles H. Burritt.

1902.—Abstract of the Mining Laws (in force in the Philippines, 1902). Charles H. Burritt.

1902, Bulletin No. 1.—Platinum and Associated Rare Metals in Placer Formations. H. D. McCaskey, B. S.

1903.—Report of the Chief of the Mining Bureau of the Philippine Islands. Charles H. Burritt.

1903, Bulletin No. 2.—Complete List of Spanish Mining Claims Recorded in the Mining Bureau. Charles H. Burritt.

1903, Bulletin No. 3.—Report on a Geological Reconnoissance of the Iron Region of Angat, Bulacan. H. D. McCaskey, B. S.

1904.—Fifth Annual Report of the Mining Bureau. H. D. McCaskey.

1905.—Sixth Annual Report of the Chief of the Mining Bureau. H. D. McCaskey.

1905, Bulletin No. 4.—A Preliminary Reconnoissance of the Mancayan-Suyoc Mineral Region, Lepanto, P. I. A. J. Eveland, Geologist.

1905, Bulletin No. 5.—The Coal Deposits of Batan Island. Warren D. Smith, B. S., M. A., Geologist.

DIVISION OF MINES.

1908.—The Mineral Resources of the Philippine Islands, with a Statement of the Production of Commercial Mineral Products during the year 1907, issued by Warren D. Smith, Chief of the Division of Mines.

LIST OF PREVIOUS PUBLICATIONS OF THE ETHNOLOGICAL SURVEY (NOW DIVISION OF ETHNOLOGY, BUREAU OF SCIENCE).

Vol. I.—The Bontoc Igorot, by Albert Ernest Jenks. Paper, ₱2; half morocco, ₱3.

Vol. II, Part 1.—Negritos of Zambales, by William Allen Reed. Paper, ₱0.50; half morocco, ₱1.50.

Vol. II, Part 2 and Part 3.—The Nabaloi Dialect, by Otto Scheerer. The Bataks of Palawan, by Edward Y. Miller. (Bound also in one volume with Part 1, Negritos of Zambales.) Paper, ₱0.50; half morocco, ₱1.50. Combined, half morocco, ₱3.

[1] *Vol. III.*—Relaciones Agustinianas de las razas del Norte de Luzón, by Pérez. Paper, ₱0.75; half morocco, ₱2.

Vol. IV, Part 1.—Studies in Moro History, Law, and Religion, by Najeeb M. Saleeby. Paper, ₱0.50; half morocco, ₱1.50.

DIVISION OF ETHNOLOGY.

Vol. IV, Part 2.—The History of Sulu, by Najeeb M. Saleeby. Paper, ₱1.50.

Vol. V, Part 1 and Part 2.—The Batan Dialect as a Member of the Philippine Group of Malayan Languages, by Otto Scheerer. The F and V in the Philippine Languages, by C. E. Conant. Paper, ₱1.60.

Directions for Ethnographic Observations and Collections, 1908. *For free distribution.*

All the above publications which are for sale may be obtained from the Director of Printing, Manila, P. I. All publications for free distribution may be obtained from the Librarian, Bureau of Science, Manila, P. I. Correspondents will confer a favor by returning to the Library of the Bureau of Science any bulletins which they may have in duplicate, as there is a considerable demand for bulletins out of print.

---

[1] Out of print.

CONTENTS.

Page.

MUSGRAVE, W. E., CLEGG, M. T., and POLK, MARY. Streptothricosis with Special Reference to the Etiology and Classification of Mycetoma, with a Bibliography 447

MUSGRAVE, W. E., CLEGG, M. T., and POLK, MARY. Trichocephaliasis, with a Report of Four Cases, Including One Fatal Case, with a Bibliography 545

EDITORIAL 567

REVIEW 569

Index, Title-page, and Contents to Volume III.

The "Philippine Journal of Science" is issued as follows:

Section A. General Science, $2, United States currency, per year.
Section B. Medical Sciences, $3, United States currency, per year.
Section C. Botany, $2, United States currency, per year.

The entire "Journal" $5, United States currency, per year.
Single numbers, 50 cents, United States currency.
Authors receive 100 copies of their papers free.
The numbers in each section will appear as rapidly as material is available. Each section will be separately paged and indexed. Subscriptions may be sent to the DIRECTOR OF PRINTING, Manila, P. I.

FOREIGN AGENTS.

THE MACMILLAN COMPANY, 64-66 Fifth Avenue, New York City.
Messrs. WM. WESLEY & SON, 28 Essex Street, Strand, London, W. C.
Mr. MARTINUS NIJHOFF, Nobelstraat 18, The Hague, Holland.
Messrs. MAYER & MÜLLER, Prinz Louis Ferdinandstrasse 2, Berlin, N. W.
Messrs. KELLY & WALSH, LIMITED, 32 Raffles Place, Singapore, S. S.
Messrs. A. W. & J. FERGUSON, 19 Baillie Street, Colombo, Ceylon.

(Entered at the post-office at Manila, P. I., as second-class matter.)

A limited number of unbound copies of previous volumes are available which may be secured from the Director of Printing at the following rates: Volume I, 1906 (not divided into sections), $5, United States currency; Supplement to Volume I, $2.50, United States currency; Volume I (without supplement), $3, United States currency; all later volumes at current subscription prices.